

- (13) Cyclization of the bisallene by a  $6\pi$ -electron process followed by prototropic rearrangement to restore the benzene sextet is an alternative possibility which cannot, at present, be excluded.
- (14) Hauptmann has observed the formation of a 2,4-disubstituted 6,7-di-*tert*-butyl-3-thiabicyclo[3.2.0]hepta-1,4-diene on treatment of 1,5-di-*tert*-butyl-3-bromopenta-1,4-diyne with sodium sulfide. This reaction is also suggested to proceed through a bisallene intermediate which then undergoes intramolecular dimerization. See H. Hauptmann, *Tetrahedron Lett.*, 3589 (1974).
- (15) For a related example in which the benzene rings do not become involved in the cyclization see H. A. Staab and B. Draeger, *Chem. Ber.*, **105**, 2320 (1972).
- (16) Attempts to trap the diradical intermediate with olefins have so far been unsuccessful.

Peter J. Garratt,\* Soon Bin Neoh

Department of Chemistry, University College London  
London WC1H 0AJ, England


Received January 17, 1975

### Quaternary Ammonium Enolates as Synthetic Intermediates. Regiospecific Alkylation Reaction of Ketones

Sir:

For the regiospecific alkylation of ketones<sup>1</sup> specific enolate anions have been generated by the aid of various precursors among which silyl enol ethers are the most versatile.<sup>2</sup> With all these specific enolates, some problems remain unresolved: concomitant formation of dialkylation products and loss of regiospecificity.<sup>3</sup> This is attributable to the low reactivities of lithium enolates toward alkyl halides. Substitution of the counterion from lithium to quaternary ammonium<sup>4</sup> seems to activate the enolate anion and to make the regiospecific monoalkylation more feasible.

We have examined the reactions of silyl enol ethers with benzyltrimethylammonium fluoride (BTAF) in the presence of alkyl halides under the expectation that BTAF would be a reagent to cleave the silicon-oxygen bond of silyl enol ethers to afford quaternary ammonium enolates because of high reactivity of fluoride anion in aprotic media<sup>5</sup> and the high value of silicon-fluorine bond energy.<sup>6</sup>



The regiospecific benzylation of 2-methylcyclohexanone via the trimethylsilyl enol ether **2** is illustrative. BTAF<sup>7</sup> (219 mg, 1.30 mmol) and 4A molecular sieves (1.0 g) were suspended in 4 ml of tetrahydrofuran (THF) and stirred overnight under an argon atmosphere.<sup>8</sup> A solution of the silyl enol ether **2** (213 mg, 1.16 mmol), and benzyl bromide (171 mg, 1.00 mmol) in 2 ml of THF was added to the suspension, and the reaction mixture was stirred for 1 hr at room temperature and then at 50° for 30 min. Filtration and removal of solvent afforded a pale yellow oil which by gas chromatographic (GLPC) analysis (XE-60) proved to contain little by-products. The crude reaction mixture was chromatographed on silica gel to give *cis*-6-benzyl-2-methylcyclohexanone (19 mg, 9%), and the *trans* isomer (161 mg, 80%).<sup>3a,b</sup> The NMR spectra of both compounds absolutely lacked the characteristic signals of 2-benzyl-2-methylcyclohexanone.<sup>3a,b</sup>




Table I. Alkylation of Ketones via Quaternary Ammonium Enolates

Silyl enol ether	RX	Molar ratio, silyl enol ether:RX <sup>a</sup>	React condn <sup>b</sup>	Yield, <i>c,d</i> %
1	Benzyl bromide	1.0:1.0	A	63
1	Methyl iodide	1.0:20	B <sup>e</sup>	79 <sup>f</sup>
2	Benzyl bromide	1.2:1.0	A	89
2	Cinnamyl bromide	1.0:1.0	A	78
2	Methyl bromoacetate	1.2:1.0	B	56
2	Butyl iodide	1.0:10	B <sup>e</sup>	40
3	Benzyl bromide	2.0:1.0	A	59
3	Benzyl bromide	1.0:1.1	C	37
3	Methyl bromoacetate	1.2:1.0	A	42
3	Butyl iodide	1.0:3.0	A	37
4	Benzyl bromide	1.0:2.2	A	67
4	Allyl bromide	1.0:2.0	B	58

<sup>a</sup> BTAF was used in a little excess (1.1 equiv) to silyl enol ethers.

<sup>b</sup> The reaction conditions are not necessarily optimum: (A) 1 hr at room temperature, 0.5 hr at 50°; (B) 14 hr at room temperature; (C) 2 hr at room temperature. <sup>c</sup> Isolation yield. <sup>d</sup> All compounds were characterized by NMR, ir, and elementary analysis. <sup>e</sup> No solvent was used. <sup>f</sup> Yield was determined by GLPC (QF-1) using tetralin as an internal standard.

The results are summarized in Table I. In every crude reaction mixture obtained after filtration of the precipitate were found the regiospecifically monoalkylated ketone, the unreacted alkyl halide, and the ketone resulting from simple hydrolysis of the starting silyl enol ether. Practically, neither dialkylation nor other by-products were detected.

The regiospecificity is lost to some extent in the alkylation reaction of the lithium enolate corresponding to the silyl enol ether **2**.<sup>3a,b,c</sup> On the other hand, both of the quaternary ammonium enolates formed from the silyl enol ether **2** and **3** underwent regiospecific alkylation. In addition to reactive alkyl halides, relatively less reactive alkylating agents, butyl iodide and methyl bromoacetate, also alkylated 2-methylcyclohexanone regiospecifically,<sup>9</sup> while the lithium enolate corresponding to the silyl enol ether **2** fails to give the desired product on butylation.<sup>10</sup>

With regard to the regiospecificity and the yields of monoalkylation products, the quaternary ammonium enolate described here is useful especially for the alkylation at the less highly substituted position of cyclohexanone derivatives, whereas the lithium enolate is suited for that at the highly substituted position. They appear to compensate each other.

The other features of this alkylation procedure, the preferential attack of the fluoride anion on silicon, the mild reaction conditions,<sup>11</sup> and the simplicity of the procedure, should also be noted.

### References and Notes

- (1) For a review, see H. O. House, "Modern Synthetic Reactions", 2nd ed, W. A. Benjamin, New York, N.Y., 1972, Chapter 9.
- (2) G. Stork and P. F. Hudrik, *J. Am. Chem. Soc.*, **90**, 4462, 4464 (1968); H. O. House, L. J. Czuba, M. Gall, and H. D. Olmstead, *J. Org. Chem.*, **34**, 2324 (1969).
- (3) (a) H. O. House, M. Gall, and H. D. Olmstead, *J. Org. Chem.*, **36**, 2361 (1971); (b) M. Gall and H. O. House, *Org. Synth.*, **52**, 39 (1972); (c) I. J. Borowitz, E. W. R. Casper, R. K. Crouch, and K. C. Yee, *J. Org. Chem.*, **37**, 3873 (1972). The reactions of lithium enolates formed from enol phosphorylated derivatives are reported. The method described therein, i.e., adding a lithium enolate to an alkyl halide, is essential for the monoalkylation of lithium enolates. (d) A highly reactive alkyl halide,  $\gamma$ -iododiglate, alkylates lithium enolates regiospecifically in high yields: P. L. Stotter and K. A. Hill, *J. Am. Chem. Soc.*, **96**, 6524 (1974).
- (4) For the general property of quaternary ammonium salts, see M. L. Bender, "Mechanisms of Homogeneous Catalysis from Protons to Proteins", Wiley-Interscience, New York, N.Y., 1971, pp 205-206; see also D. J. Sam and H. E. Simmons, *J. Am. Chem. Soc.*, **96**, 2252 (1974).
- (5) C. L. Liotta and H. P. Harris, *J. Am. Chem. Soc.*, **96**, 2250 (1974).
- (6) L. Pauling, "The Nature of Chemical Bond", 3rd ed, Cornell University Press, New York, N.Y., 1960, Chapter 3.
- (7) BTAF was prepared by neutralizing commercial Triton B (40% metha-

- nolic solution) by aqueous 47% HF, dried at 100°, <1 mm for 24 hr, finely pulverized, dried again for 24 hr, and stored over P<sub>2</sub>O<sub>5</sub>.
- (8) Thus highly hygroscopic BTAF can be manipulated in air, otherwise it should be manipulated in a drybox.
  - (9) The homogeneity of the products was examined by GLPC (QF-1), NMR, and by the aid of equilibration under basic conditions; see, for example, G. H. Posner, J. J. Sterling, C. E. Whitten, C. M. Lentz, and D. J. Brunelle, *J. Am. Chem. Soc.*, **97**, 107 (1975).
  - (10) 2-Butyl-2-methylcyclohexanone is the sole alkylation product, even when the lithium enolate corresponding to **2** is alkylated with butyl iodide, ref 3c.
  - (11) Using tetrabutylammonium fluoride, Corey succeeded in the removal of trialkylsilyl groups in the presence of various functional groups: E. J. Corey and B. B. Snider, *J. Am. Chem. Soc.*, **94**, 2549 (1972); E. J. Corey and A. Venkateswarlu, *ibid.*, **94**, 6190 (1972).

Isao Kuwajima,\* Eiichi Nakamura

Laboratory of Organic Chemistry, Tokyo Institute of  
Technology, Ookayama, Meguro-ku, Tokyo, Japan

Received February 24, 1975

### Observation of Photoinduced CIDNP at Low Radiation Levels. Proton CIDNP from Laser Irradiation of Di-*tert*-butyl Ketone

Sir:

We have observed proton CIDNP (chemically induced dynamic nuclear polarization) during laser irradiation of a solution of di-*tert*-butyl ketone (DTBK) in the probe of an NMR spectrometer. Average laser power at the sample was only 0.7 mW. Previously, photolytically induced CIDNP has been observed only with high power broadband arc lamp excitation sources.<sup>1</sup>

For this study, a Chromatix Model 1050 tunable uv-visible laser system was used.<sup>2</sup> The radiation was directed into the quartz sample tube from below by a low-loss mirror. Calculated beam diameter at the sample (5 ft from the laser) was 3.6 mm. The Varian HA-100 NMR spectrometer was in its normal configuration, except that the variable temperature attachment was removed from the bottom of the V-4333 probe. CIDNP occurred throughout the laser tuning range of 285–305 nm. Best signal levels were obtained with 0.03–0.07 *M* solutions of di-*tert*-butyl ketone in carbon tetrachloride. This contrasts with sample concentrations of 0.1–0.3 *M* used with 1 KW arc sources.<sup>3,4</sup>

Strong CIDNP signals were observed for the aldehydic proton of pivaldehyde (9.35 ppm, A), chloroform (7.24 ppm, A), isobutylene (4.60 ppm, A; 1.70 ppm, A), and *tert*-butyl chloride (1.59 ppm, E). The net polarizations of the products are consistent with those predicted by Kaptein's rules<sup>5</sup> for a triplet precursor pivaloyl-*tert*-butyl radical pair. Arc source photolysis of di-*tert*-butyl ketone in perfluoromethylcyclohexane doped with carbon tetrachloride yielded identical results.<sup>3</sup>

The observation of CIDNP at such low photon fluxes ( $\approx 10^{15}$  photons/sec) is surprising. It is commonly stated that  $10^{17}$ – $10^{18}$  photons/sec are required for practical photochemical results,<sup>6</sup> i.e., the formation of detectable photo-product in a reasonable time. The success of this experiment can be attributed to a proper combination of several factors.

The HA-100 spectrometer can detect  $\sim 10^{-3}$  *M* protons, or  $\sim 2.5 \times 10^{17}$  spins in the volume of 0.4 cm<sup>3</sup> used in our experiment. A photoproduct with a quantum yield of one and a CIDNP enhancement factor of  $\approx 100$  will yield an effective measurable concentration. At 313 nm the decomposition quantum yield of DTBK is 0.7.<sup>7</sup> For the products of the DTBK photoreaction, Fischer<sup>4b</sup> has reported enhancement factors of  $10^2$ – $10^3$ . Although DTBK reacts from both singlet and triplet states, the predictions for the sign of CIDNP polarizations are the same for the products which

can form by either path (*tert*-butyl chloride and isobutylene).<sup>8</sup> The decrease in yield of a particular product as a result of the several competing reactions<sup>4b,8</sup> causes less than an order of magnitude change from the maximum rate of  $\sim 10^{15}$  molecules/sec.

In anticipation of the problems associated with observing CIDNP at low photon flux levels we had determined that the sample tube could be positioned so that its bottom was only 5 mm below the receiver coil of the probe with no loss of resolution or signal-to-noise ratio. This minimum "cell thickness" and the proper substrate concentration were important parts of the experiment. At DTBK concentrations of 0.03–0.07 *M* (OD 0.9–2.3 at 296 nm) and with a laser beam diameter of 3.6 mm, most of the incident light was absorbed within the active volume of the receiver coil. We could not observe CIDNP at ketone concentrations greater than  $\sim 0.14$  *M*. Above that concentration, unpolarized photoproducts were observed building up with time. Evidently, if the optical density is too high, the polarized photoproducts are formed below the active receiver coil volume, and lose their polarization before diffusing into it.<sup>9</sup> We also observed this concentration effect in our study of C-13 CIDNP during arc source photolysis of DTBK.<sup>8</sup> In that work the total power reaching the sample was  $\sim 200$  mW (250–340 nm) and CIDNP was observed for 25% solutions (1.44 *M*). The CIDNP signal intensity decreased for more concentrated solutions.

The observation of proton CIDNP at photon fluxes as low as  $\sim 10^{15}$ /sec is thus conditional upon four factors: (a) a high quantum yield for polarized product(s), (b) CIDNP enhancement factors of 100 or more, (c) optimum sample optical density, and (d) location of the irradiated region within the active receiver coil volume.

### References and Notes

- (1) The photo-CIDNP literature was recently reviewed: H. D. Roth, *Mol. Photochem.*, **5**, 91 (1973).
- (2) The laser was operated in the burst, Q-switched mode at 75 bursts/sec, 4–6 pulses per burst. Peak power ( $P_p$ ) in the uv was about 85 W, with a pulse width at half-height ( $W_{1/2}$ ) of 70 nsec. The calculated average power ( $P_p \times W_{1/2} \times \text{pps}$ ) of 2.2 mW at 290 nm was confirmed by measurements with an Eppley thermopile. Losses in the mirror system used to direct the beam into the probe reduced the measured average power reaching the sample to 0.7 mW.
- (3) M. Tomkiewicz, A. Groen, and M. Cocivera, *Chem. Phys. Lett.*, **10**, 39 (1971); *J. Chem. Phys.*, **56**, 5850 (1972).
- (4) (a) H. Fischer and G. P. Laroff, *Chem. Phys.*, **3**, 217 (1974); (b) B. Blank, A. Henne, and H. Fischer, *Helv. Chim. Acta*, **57**, 920 (1974).
- (5) R. Kaptein, *Chem. Commun.*, 732 (1971).
- (6) C. R. Masson, V. Boekelheide, and W. A. Noyes, Jr., in "Technique of Organic Chemistry", Vol II, 2nd ed, A. Weissberger, Ed., Interscience, New York, N.Y., 1956, p 271.
- (7) N. C. Yang, E. D. Feit, M. H. Hui, N. J. Turro, and J. C. Dalton, *J. Am. Chem. Soc.*, **92**, 6974 (1970).
- (8) W. B. Moniz, C. F. Poranski, Jr., and S. A. Sojka, submitted for publication.
- (9) This consideration is not so important if the light is applied from the side directly into the region of the receiver coil (ref 4b).

W. B. Moniz,\* C. F. Poranski, Jr.

Chemistry Division, Naval Research Laboratory  
Washington, D.C. 20375

Received February 10, 1975

### A Route to Prostaglandins via a General Synthesis of 4-Hydroxycyclopentenones

Sir:

We wish to report a general method for the synthesis of hydroxycyclopentenones of type **1**. These are versatile substances especially because we have been able to effect their ready transformation into the isomeric hydroxycyclopenten-