





Scheme I<sup>a</sup>


Whereas the dictyoxepin structure has been confirmed by the x-ray crystallographic work of Clardy,<sup>1</sup> that of dictyolene rests solely upon spectral data and biogenetic analogy to dictyoxepin, especially with regard to the stereochemistry at C-7 and C-11. In this report, we describe definitive stereorational total syntheses of dictyolene (**20**) and its C-11 epimer **12** which fully support the assigned structure II.

Dienal **1**, readily prepared from *m*-toluic acid in 71% overall yield,<sup>2</sup> upon alkylation with ethyl bromoacetate via the diisopropyl enamine and subsequent hydrolysis, afforded the ester aldehyde **2** in 40% yield<sup>3</sup> (Scheme I). Cyclization of this aldehyde with perchloric acid in ethyl acetate led to the cis,anti-cis tricyclic lactone **3**, mp 69–71 °C, along with the isomeric cis,anti,trans hydroxy ester **21**. These two products, formed



as a roughly 1:1 mixture in 90% yield, could be separated by saponification and careful acidification, whereupon a mixture of lactone **3** and hydroxy acid **22** was produced. Acid **22** could then be removed through base extraction. The assignment of structure for lactone **3** follows from (1) the established course of such cyclizations,<sup>2</sup> (2) its ready formation upon acidification of the hydroxy acid salt mixture,<sup>4</sup> and (3) comparison with an independently synthesized sample.<sup>5</sup>


Alkylation of lactone **3** with 2-methyl-5-iodo-2-pentene<sup>6</sup> yielded lactone **4** while the use of methyl iodide gave rise to the



crystalline lactone **5**, mp 96–97 °C (91% yield). In each case, the stereochemical outcome conforms to approach from the least hindered face of the lactone enolate **3a** by the alkylating agent (kinetic alkylation). Furthermore, since treatment with lithium diisopropylamide failed to epimerize lactones **4** and **5**, we surmise that they represent the more stable alkyl epimers, as well. Inspection of molecular models (see conformational drawings) allows a clear choice of stereochemistry for these key lactone intermediates.

According to the assignment suggested by Erickson, Clardy, et al.,<sup>1</sup> conversion of the lactone carbonyl of **4** to a methyl grouping should lead to the C-11 epimer of dictyolene, **12**. Indeed, this was found to be the case. Reduction of lactone **4** with lithium aluminum hydride gave diol **6** which could be selectively protected with *tert*-butyldimethylsilyl chloride to give ether **7**.<sup>7</sup> Acetylation of this intermediate followed by cleavage of the silyl ether **8** with aqueous hydrochloric acid and mesylation of the resulting alcohol **9** afforded acetoxy mesylate **10**. This mesylate was reduced with sodium cyanoborohydride and sodium iodide in HMPA<sup>8</sup> to acetate **11** which, upon treatment with lithium aluminum hydride, yielded alcohol **12**. The infrared and NMR spectra of this alcohol closely resembled those of dictyolene (**11**) but clear differences could be readily discerned, particularly in the chemical shift of the C-11 methyl group.<sup>9</sup>

In our initial planning, we had hoped to convert mesylate **17**, prepared from lactone **5** using the sequence employed in the conversion of lactone **4** to mesylate **10**, to dictyolene through coupling with a 3,3-dimethylallylcopper reagent.<sup>10</sup> However, preliminary experiments with either catalyzed or stoichiometric reagents on the mesylate **23** were totally unpromising; only the reduction product **24** was formed.<sup>11</sup> Ac-



cordingly, we turned to the use of 3,3-dimethylallylnickel bromide, a reagent employed by Corey and Semmelhack in their synthesis of  $\alpha$ -santalene.<sup>12,13</sup> Reaction in dimethylformamide with the iodo acetate **18**, derived from mesylate **17** using sodium iodide in acetone,<sup>6</sup> afforded the prenylated homologue **19**. Cleavage with lithium aluminum hydride gave racemic dictyolene (**20**) whose spectral properties were identical with those of authentic material.<sup>9</sup>

**Acknowledgments.** We are indebted to the National Institutes of Health for support of this work through Research Grant CA 11089 awarded by the National Cancer Institute.

## References and Notes

- H. H. Sun, S. M. Waraszkievicz, K. L. Erickson, J. Finer, and J. Clardy, *J. Am. Chem. Soc.*, **99**, 3516 (1977).
- J. A. Marshall and P. G. M. Wuts, *J. Org. Chem.*, **42**, 1794 (1977).
- Enamines of aldehydes are readily prepared and handled when hindered secondary amines are employed. Cf. T.-L. Ho and C. M. Wong, *Synth. Commun.*, **4**, 147 (1974), and references therein.
- In related systems we have found that trans-fused  $\gamma$ -butyrolactones form with appreciably greater difficulty than the corresponding cis-fused isomers. For a possible rationale, see J. A. Marshall and N. Cohen, *J. Org. Chem.*, **30**, 3475 (1965), and references therein.

- (5) J. A. Marshall and P. G. M. Wuts, *J. Org. Chem.*, in press.  
 (6) Prepared from the known bromide using sodium iodide in acetone: M. Julia, S. Julia, and R. Guégan, *Bull. Soc. Chim. Fr.*, 1072 (1960).  
 (7) E. J. Corey and A. Venkateswarlu, *J. Am. Chem. Soc.*, **94**, 6190 (1972).  
 (8) R. O. Hutchins, B. E. Maryanoff, and C. A. Milewski, *Chem. Commun.*, 1097 (1971).  
 (9) We are indebted to Professor K. L. Erickson for a preprint of her publication and for copies of the infrared and NMR spectra of dictyolene.  
 (10) G. H. Posner, *Org. React.*, **22**, 253 (1974); G. M. Whitesides, W. F. Fisher, Jr., J. San. Filippo, Jr., R. W. Bashe, and H. O. House, *J. Am. Chem. Soc.*, **91**, 4871 (1969).  
 (11) A number of reasonable pathways may be formulated for this unusual reduction. Presumably, a  $\gamma$ -hydrogen atom of the prenylcopper reagent is transferred with the eventual formation of isoprene. We plan to examine this reaction in more detail.  
 (12) E. J. Corey and M. F. Semmelhack, *J. Am. Chem. Soc.*, **89**, 2755 (1967).  
 (13) M. F. Semmelhack, *Org. React.*, **19**, 115 (1972).

James A. Marshall,\* Peter G. M. Wuts

Department of Chemistry, Northwestern University  
 Evanston, Illinois 60201


Received September 26, 1977

**$\pi, \pi$ -Biradicaloid Hydrocarbons:  
 Spectral Characterization of Singlet and Triplet  
 1,3-Perinaphthadiyl and Observation of  
 Thermal and Photochemical 2,3-Hydrogen Shifts<sup>1</sup>**

Sir:

Although biradicaloid geometries<sup>2</sup> are likely to play a fundamental role in molecular photochemical processes,<sup>3,4</sup> very little is known from direct observations about molecules at such geometries and, in particular, about the reactivity and physical properties of their excited states. We now wish to report the preparation and spectral characterization of the lowest singlet and triplet states of a trimethylene biradical constrained to a  $\pi, \pi$  geometry by interaction with the peri positions of naphthalene, as well as the kinetics of a thermal 2,3-hydrogen shift and the observation of a photochemical 2,3-hydrogen shift in the biradical.

When I<sup>5</sup> is irradiated at 77 K in a rigid glass ( $\lambda \approx 260$  nm,  $\sim 10^{-3}$  M solution in degassed 3-methylpentane, 2-methyltetrahydrofuran, 1-pentanol, or EPA), its UV absorption, emission, and excitation spectra are gradually replaced by those of a new species stable for many hours (Figure 1). Simultaneously, the characteristic lines of a triplet appear in the ESR spectrum ( $D/hc = 0.026 \pm 0.001$  cm<sup>-1</sup>,  $E/hc < 0.002$  cm<sup>-1</sup>). Each member of the two  $\Delta m = 1$  pairs of observed lines of the species produced from Ia, as well as the  $\Delta m = 2$  line, is



split into a triplet. The coupling constant is approximately the same for both pairs,  $a_H = 26 \pm 2$  G. The lines of the species produced from Ib are similarly split into very poorly resolved doublets. Temperature variation of the ESR intensity (93–123 K, 1-pentanol) is strong, and standard analysis shows that the triplet state lies  $640 \pm 40$  cal above the ground state. The




Figure 1. Absorption (—), fluorescence (---), and fluorescence excitation (···) spectra of II (3-MP, 77 K). The excitation spectrum is on two unrelated scales.

UV-vis absorption spectrum shows only a very slight decrease in intensity when temperature is raised from 77 to 110 K, while the shapes of the fluorescence and fluorescence excitation spectra show no significant change. All of the bands in the optical spectra appear to belong to only one species (fluorescence excitation follows absorption). Thus, they are either all due to light absorption by the singlet ground state, whose fraction in the  $S_0$ - $T_1$  mixture should only vary between 95 and 86% over this temperature range, or else are all due to a completely unrelated species. The latter alternative is made extremely unlikely by the observation that at higher temperatures both the ESR and the UV signals exhibit a first-order decay in time and that all of the rate constants fall on the same straight line in an Arrhenius plot (119–133K, annealed 1-pentanol,  $\Delta E_{act} = 4.5 \pm 0.6$  kcal/mol,  $A = 10^{4.5 \pm 1}$  s<sup>-1</sup>). Our failure to observe the UV-vis absorption of the  $T_1$  state is understandable, since it comprises only a minor fraction of the mixture and its calculated (PPP<sup>6</sup>) transition energies and intensities lead us to suspect strong overlap with the UV-vis absorption of the  $S_0$  state.

The product of the thermal transformation of the new species is II, easily identified by its UV absorption, emission, and excitation spectra. Measurement of its rate of formation by UV absorption yields the same rate constants as above. Also, UV or visible irradiation of the new species produces II (77 K). Comparison of quantitative absorption spectra of I, of the new species, and of II showed that both photochemical steps are quantitative and this permitted us to estimate the extinction coefficients of the new species given in Figure 1 (uncorrected for the fraction present in  $T_1$ ). Similarly, the thermal reaction is found to be quantitative in the thermal range given. At higher temperatures in 1-pentanol, and already at lower temperatures in less viscous glasses, thermal disappearance of the new species is more complex. The phenalenyl radical is observed in ESR and naphthalene-like absorption in UV spectra. Bimolecular reactions are presumably facile at these lower viscosities, and some return to I may be occurring as well.

On basis of the combined evidence, we postulate the planar structure III for the new species and presume that  $S_0$ - $T_1$  equilibration is rapid (cf. the related 1,8-naphthoquinodimethanes IV<sup>7</sup> ( $D/hc = 0.0218$  cm<sup>-1</sup>,  $E/hc = 0.0021$  cm<sup>-1</sup>,  $T_1$  45 cal/mol above  $S_0$ , stable at 77 but not at 87 K) and V<sup>8</sup> ( $D/hc = 0.018$  cm<sup>-1</sup>,  $E/hc < 0.003$  cm<sup>-1</sup>,  $T_1$  200 cal/mol above  $S_0$ ), as well as the related localized 1,3 biradical, VI<sup>9</sup> ( $D/hc = 0.084$  cm<sup>-1</sup>,  $E/hc = 0.0020$  cm<sup>-1</sup>), with poorly resolved hyperfine structure). The larger  $S_0$ - $T_1$  splitting in III is readily assigned to the effect of cyclic hyperconjugation with the methylene group, which raises the energy of the symmetrical nonbonding orbital and leaves that of the antisymmetrical nonbonding orbital essentially intact. Such splitting of the two orbitals should be even larger in VI (by about a factor of 1.5