

commenced. Diisobutylaluminum hydride (5 equiv) reduction of **9** at 0 °C (acid workup), followed by manganese dioxide oxidation of the resulting allylic alcohol, gave the aldehyde-cyclopentenone **10** in 65% yield. Ozonolysis of this material (−78 °C 2 h) in methylene chloride containing pyridine (1.1 equiv) and workup with dimethyl sulfide provided the lactol **11** in 70% yield.⁹ Treatment of the latter with methanol, trimethyl orthoformate, and acetyl chloride afforded the corresponding bis acetal which on reaction with methylene triphenylphosphorane in THF gave the diene **12**.¹⁰ Lastly, reduction of the diene portion of **12** using tris(triphenylphosphine)chlororhodium in benzene (30 psi, H₂),¹¹ followed by acetal hydrolysis (acetone, H₂SO₄), Jones oxidation, and diazomethane treatment, afforded the lactone ester **13** (oil, 49% from **11**) as a 2:1 mixture of methyl group epimers at C₁₄—the β-methyl isomer predominating.¹²

The above mixture of epimers could not be readily separated by chromatography and, therefore, was utilized as such for elaboration into the α-methylene lactone **14**. Several procedures for conversion of **13** into **14** were examined and by far the most convenient and efficient involved treatment of **13** with methoxymagnesium carbonate (20 equiv, 160 °C, 2 h), followed by reaction of the resulting lactone acid with 30% formalin solution containing diethylamine.¹³ Vacuum filtration chromatography of the resulting mixture gave a 55% yield of the α-methylene lactone **14** (oil), determined by ¹H NMR spectroscopy to be exclusively the β-methyl isomer at C₁₄.¹⁴

Comparison of **14** with a sample of the α-methylene lactone (oil) kindly provided by Professor S. Danishefsky conclusively demonstrated these substances to be identical.¹⁵ Since Danishefsky and co-workers have stereoselectively converted **14** into pentalenolactone (**1**) in good overall yield,^{2a} our assemblage of **14** constitutes a total synthesis of this natural product. The preparation of **14** from 3-methoxy-2-methylcyclopentenone requires 19 steps and proceeds in an overall yield of 5.3%.

Acknowledgment. This research was supported by PHS Grant CA-21496. Support from the Syntex Corporation and the Hoffmann-La Roche Foundation is gratefully acknowledged. We thank Mr. A. Lopes for help in preparing quantities of compounds **3** and **4**.

References and Notes

- (1) (a) D. G. Martin, G. Slomp, S. Mizsak, D. J. Duchamp, and C. G. Chidester, *Tetrahedron Lett.*, 4901 (1970); (b) S. Takeuchi, Y. Ogawa, and H. Yonehara, *ibid.*, 2737 (1969); (c) S. Takeuchi, J. Uzawa, H. Seto, and H. Yonehara, *ibid.*, 2943 (1977); (d) H. Seto, T. Sasaki, H. Yonehara, and J. Uzawa, *ibid.*, 923 (1978).
- (2) (a) S. Danishefsky, M. Hiram, K. Gombatz, T. Harayama, E. Berman, and P. Schuda, *J. Am. Chem. Soc.*, **100**, 6536 (1978); (b) F. Playac and C. H. Heathcock, *Tetrahedron Lett.*, 2115 (1979), have recently described yet another interesting potential route to this natural product.
- (3) The numbering for pentalenolactone used in this text follows that given in ref 1a.
- (4) A detailed preparation of this compound has been described by M. L. Quesada, R. H. Schlessinger, and W. H. Parsons, *J. Org. Chem.*, **43**, 3968 (1978).
- (5) For an account of similar alkylation procedures, see ref 4. Compound **3**, as well as all other substances described herein, has been fully characterized.
- (6) Since **4** is a 1:1 mixture of epimers and since the bicyclic product **2** is formed in >50% yield, it seems clear that epimerization is occurring at C₅ during the course of reaction—when epimerization of C₅ occurs (before or after cyclization) is not known with certainty by us. Reaction sequences for which an overall yield is reported normally were carried out using crude intermediates—only the final product of a given sequence was purified.
- (7) The transformation **2** → **7** was accomplished by (a) sodium borohydride reduction of the cyclopentanone residue and protection of the resulting secondary alcohol residue with chloromethylmethyl ether; (b) ozonolysis of the allyl moiety, followed by reductive workup (sodium borohydride) and subsequent acidification to yield the lactone.
- (8) Deprotonation of **2** in the kinetic manner affords the enolate derived by abstraction of the angular methine proton at C₈. This differing course of deprotonation (kinetic vs. thermodynamic) was demonstrated by alkylation experiments. The source of this phenomenon may well lie in the considerable strain experienced by the kinetic enolate and caused by sp² hybridization of five of the eight carbon atoms present in the pentalene skeleton of **2**.
- (9) Pyridine is critical to the success of this reaction. For a leading reference concerning the effect of pyridine on ozonolysis reactions of this type, see G. Slomp and J. L. Johnson, *J. Am. Chem. Soc.*, **80**, 915 (1958).
- (10) Reaction of either the lactone-acid or the lactone-ester analogues of **11** with this Wittig reagent was not successful. Undoubtedly, these results are due to the acidity of the C₈ methine hydrogen—hence the conversion of **11** into its bis acetal derivative.
- (11) J. W. Osborn, F. H. Jardine, J. F. Young, and G. Wilkinson, *J. Chem. Soc. A*, 1711 (1966).
- (12) This ratio was readily determined by ¹H NMR examination of the doublets representing the two methyl epimers—the β isomer exhibiting a chemical shift and coupling constant essentially identical with those of C₁₄ methyl group of the natural product. Danishefsky reports^{2a} completely stereoselective reduction of an A-ring diene system possessing a carbonyl group at C₁₁, a single bond between C₆ and C₇, as well as a β-oriented carbomethoxy group attached to C₈. It is our feeling that the hindering effect of this C₈ carbomethoxy group may well enhance the stereoselectivity of this reduction.
- (13) For a detailed description of this highly useful methylenation sequence, see W. H. Parker and F. J. Johnson, *J. Org. Chem.*, **38**, 2489 (1973).
- (14) The authors claim no credit for this clearly fortuitous experimental event—the origin of which is yet obscure.
- (15) We thank Professor Danishefsky for generous samples of both the α-methylene lactone **14** and methyl pentalenolactonate, as well as for stimulating discussion during the course of this work.

W. H. Parsons, R. H. Schlessinger,* Martha L. Quesada

Department of Chemistry, University of Rochester
Rochester, New York 14627

Received August 28, 1979

Additions and Corrections

Three-Electron Oxidations. VIII. Direct Evidence for the Synchronous Character of Three-Electron Oxidations [*J. Am. Chem. Soc.*, **96**, 6802 (1974)]; **Three-Electron Oxidations. IX. Chromic Acid Oxidation of Glycolic Acid** [*Ibid.*, **97**, 1444 (1975)]; **Three-Electron Oxidations. X. Cooxidation of Isopropyl Alcohol and Glycolic Acid** [*Ibid.*, **97**, 3762 (1975)]. By FARIZA HASAN and JAN ROČEK,* Department of Chemistry, University of Illinois at Chicago Circle, Chicago, Illinois 60680.

One of the original authors (J.R.) and S. Ramesh were unsuccessful in attempts to reproduce the synthesis of glycolic-*d*₂ acid, HOCD₂CO₂H, described in the original publications. Glycolic-*d*₂ acid prepared by two other methods gave considerably lower deuterium isotope effects than originally reported.

We are therefore forced to conclude that the unusually high values for deuterium isotope effects reported in Tables I, V, and IV, respectively, of the original set of publications were in error. Results of a full reinvestigation of the chromic acid oxidation of glycolic acid and its cooxidation with isopropyl alcohol will be reported as soon as completed.

Hydrogen Atom Exchange between Nitroxides and Hydroxylamines [*J. Am. Chem. Soc.*, **101**, 3592 (1979)]. By MARTIN A. SCHWARTZ, J. WALLACE PARCE, and HARDEN M. MCCONNELL,* Stauffer Laboratory for Physical Chemistry, Stanford University, Stanford, California 94305.


In the first sentence in the Experimental Section, 2-Methyl-2-nitro-5-pentanone should be replaced by 2-

Methyl-2-nitro-5-hexanone, and ethyl vinyl ketone should be replaced by methyl vinyl ketone.

A Stereocontrolled Synthesis of the (\pm) Djerassi-Prelog Lactic Acid [*J. Am. Chem. Soc.*, **101**, 1315 (1979)]. By GILBERT STORK* and VIJAY NAIR, Department of Chemistry, Columbia University, New York, New York 10027.

Page 1315, column 1; the second sentence in the first paragraph (line 2) should read, "Its structure was established¹⁻³ by Prelog and by Djerassi. . . ."

Structure **1** should appear as shown below:



Column 2, paragraph 3; line 11 should read, "C-6 and of the oxygen 'bridge' at C-5. . . ."


Reaction of a Metal Alkyl with Ethylene as a Model for Ziegler-Natta Polymerization. Evidence for the Olefin Insertion Mechanism [*J. Am. Chem. Soc.*, **101**, 3973 (1979)]. By ERIC R. EVITT and ROBERT G. BERGMAN,* Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, and the Department of Chemistry, University of California and the Materials and Molecular Research Division, Lawrence Berkeley Laboratory, Berkeley, California 94720.

Beginning in the third line of the second paragraph, the words "of **1** (δ 4.49, 0.61 ppm) and ethylene (5.27) were replaced by those from methane (0.22) and propene (1.58, CH₃), as well as by a new cyclopentadienyl signal (5.50, 5 H) and two new multiplets at 1.11 (2 H) and 2.09 (2 H) ppm" should be replaced by "of **1** (δ 4.46, 0.58 ppm) and ethylene (5.24) were replaced by those from methane (0.19) and propene (1.58, CH₃), as well as by a new cyclopentadienyl signal (4.48, 5 H) and two new multiplets at 1.07 (2 H) and 2.48 (2 H) ppm".

Coal Chemistry. 8. Reactions of Tetralin with Coal and with Some Carbon-14-Containing Model Compounds [*J. Am. Chem. Soc.*, **101**, 5009 (1979)]. By CLAIR J. COLLINS,* F. RAAEN, BEN M. BENJAMIN, P. H. MAUPIN, and W. HOWARD ROARK, Chemistry Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37830.

In the title, directly following the words "Coal Chemistry", the number 8 should be deleted.




In eq 1, first column, p 5011, the last structure is incorrect. Equation 1 should read:



After ref 14 add: See also R. A. Marty and P. DeMayo, *Chem. Commun.*, 127 (1971).

Optically Active Amines. 28. Spectral Observations on Chiral N-Substituted *p*-Nitroanilines [*J. Am. Chem. Soc.*, **101**, 5186 (1979)]. By HOWARD E. SMITH,* WILLIAM I. COZART, TOMAS DE PAULIS, and FU-MING CHEN, Departments of Chemistry, Vanderbilt University, Nashville, Tennessee 37235, and Tennessee State University, Nashville, Tennessee 37203.

On p 5190 the Newman projections should be:



Metallacyclopentane to Metallacyclobutane Ring Contraction [*J. Am. Chem. Soc.*, **101**, 5451 (1979)]. By STEPHAN J. MCLAIN, JOSÉ SANCHO, and RICHARD R. SCHROCK,* Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139.

The first structure in eq 4 should be:

