

Studies on the further use of intermediate **6** and **8** in synthesis as well as application of the β -metal ketone strategy to other naked (or ligand free) metal cations are in progress.

Acknowledgment. This work was supported in part by Grant-in-Aid for Special Project Research No. 57118002 provided by the Ministry of Education, Science, and Culture, Japan. We thank Shin-Etsu Chemical Industry Ltd. for the gift of trimethylchlorosilane.

Supplementary Material Available: Spectral data of products listed in Table I (7 pages). Ordering information is given on any current masthead page.

(Iminomethylidene)phosphines (RP=C=NR)

Curt Wentrup,^{*1a} Horst Briehl,^{1a} Gerd Becker,^{1a,b}
Gudrun Uhl,^{1a,b} Hans-Jürgen Wessely,^{1a,b}
André Maquestiau,^{1c} and Robert Flammang^{1c}

Department of Chemistry, University of Marburg
D-3550 Marburg, West Germany
Institute for Inorganic Chemistry
University of Stuttgart
D-7000 Stuttgart 80, West Germany
Laboratory of Organic Chemistry
State University of Mons
B-7000 Mons, Belgium
Received August 15, 1983

There has been much recent interest in compounds incorporating P=C^{2,3} and P=P⁴ double bonds. Furthermore, the stable (2,2-dimethylpropylidene)phosphine, (CH₃)₃CC≡P, featuring a PC triple bond has been prepared.⁵ However, previous attempts to isolate cumulenes possessing a two-coordinate phosphorus atom, RP=C=X, have failed,^{2,3,6} except in a single case, the preparation of the stable and sterically protected (CH₃)₃CP=C=NC(CH₃)₃.⁷ We now wish to report a general method of preparation of unstable (iminomethylidene)phosphines (monophosphorus analogues of carbodiimides), their direct observation by IR and mass spectrometry, and their initial chemical reactions.

The required starting materials, (iminomethyl)phosphines **1**, exist in equilibrium with carbamoylphosphines [(CH₃)₃SiP(R)-CON(Ph)Si(CH₃)₃] and smaller amounts of alkylidene phosphines RP=C(OSi(CH₃)₃)N(Ph)Si(CH₃)₃.⁸ In analogy with the reaction used in the preparation of (CH₃)₃CC≡P,⁵ the compounds **1** could be expected to eliminate hexamethyldisiloxane and thus give rise to the desired (iminomethylidene)phosphines, RP=C=NPh (Scheme I).

(1) (a) University of Marburg. (b) Present address: University of Stuttgart. (c) University of Mons.

(2) Appel, R.; Knoll, F.; Ruppert, I. *Angew. Chem.* **1981**, *93*, 771; *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 731.

(3) Becker, G.; Becker, W.; Mundt, O. *Phosphorus Sulfur* **1983**, *14*, 267.

(4) Yoshifuji, M.; Shima, I.; Inamoto, N.; Hirotsu, K.; Higuchi, T. *J. Am. Chem. Soc.* **1981**, *103*, 4587. Cowley, A. H.; Kilduff, J. E.; Newman, T. H.; Pakulski, M. *Ibid.* **1982**, *104*, 5820. Bertrand, G.; Couret, C.; Escudie, J.; Majid, S.; Majoral, J.-P. *Tetrahedron Lett.* **1982**, *23*, 3567. Couret, C.; Escudie, J.; Satg , J. *Ibid.* **1982**, *23*, 4941. Centinkaya, B.; Hudson, A.; Lappert, M. F.; Goldwhite, H. J. *J. Chem. Soc., Chem. Commun.* **1982**, 609. Centinkaya, B.; Hitchcock, P. B.; Lappert, M. F.; Thorne, A. J.; Goldwhite, H. *Ibid.* **1982**, 691. Niecke, E.; R ger, R.; Lysek, M.; Pohl, S.; Schoeller, W. *Angew. Chem.* **1983**, *95*, 495; *Angew. Chem., Int. Ed. Engl.* **1983**, *22*, 486; Cowley, A. H.; Kilduff, J. E.; Norman, N. C.; Pakulski, M.; Atwood, J. L.; Hunter, W. E. *J. Am. Chem. Soc.* **1983**, *105*, 4845.

(5) Becker, G.; Gresser, G.; Uhl, W. *Z. Naturforsch., B* **1981**, *36B*, 16.

(6) Appel, R.; Laubach, B. *Tetrahedron Lett.* **1980**, *21*, 2497.

(7) Kolodiaznyi, O. I. *Tetrahedron Lett.* **1982**, *23*, 4933; *J. Gen. Chem. USSR (Engl. Transl.)* **1982**, *52*, 2101.

(8) Itoh, K.; Fukui, M.; Ishi, Y. *J. Chem. Soc. C* **1969**, 2002. Becker, G.; Mundt, O. *Z. Anorg. Allg. Chem.* **1979**, *459*, 87. Issleib, K.; Schmidt, H.; Meyer, H. *J. Organomet. Chem.* **1980**, *192*, 33. Becker, G.; Wessely, H.-J., manuscript in preparation.






Figure 1. IR spectrum (-196 °C) of the product of pyrolysis of **3a** at 480 °C.

Scheme I



In the event, the treatment of **1** with a catalytic amount of solid NaOH led only to the isolation of the dimers **3**.⁹ **3b** and **3e** were also obtained by spontaneous decomposition of the adducts **2b** and

(9) (a) Satisfactory elemental analyses (C, H, N, P) were obtained for these compounds. (b) X-ray crystallographic determinations of **3c** and **3d** confirm the structure assignments. The two exocyclic phosphorus ligands are trans oriented. Full crystallographic data will be published: Becker, G.; H rer, J.; Riffel, H.; Uhl, G.; Wessely, H.-J., in preparation.




Figure 2. CID mass spectrum of m/z 211 from the pyrolysis of **3a** at 430 °C.

2e, the latter formed by the addition of phenyl isothiocyanate to the alkyl- or aryl(trimethylsilyl)phosphine.¹⁰

If the diphosphetanes **3** are, in fact, dimers of (iminomethylidene)phosphines, the monomers might be regenerated via a cycloreversion under suitable reaction conditions. This, indeed, is readily achieved by flash vacuum pyrolysis of **3**. Product formation was monitored by low-temperature IR spectroscopy, and a pyrolysis unit has also been attached to the ion source of a reversed-geometry Varian MAT 311A mass spectrometer¹¹ equipped with a collision chamber for obtaining CID (collisionally induced dissociation) spectra of initial pyrolysis products.

2,4-Bis(phenylimino)-1,3-diphenyl-1,3-diphosphetane (**3a**) (mp 139 °C) was vaporized at 139–145 °C (10^{-5} torr). When the pyrolysis temperature was increased to 400 °C, an IR band at 1853 cm^{-1} appeared in the pyrolyzate condensed at -196 °C. The maximum intensity of this band was observed at a pyrolysis temperature at 480 °C (Figure 1). On warming the pyrolyzate to -55 °C, the intensity of the 1853- cm^{-1} band started decreasing; at the same time, a band at 1560 cm^{-1} due to the starting material **3a** started increasing. This process was complete at -30 °C, and after warming to room temperature **3a** was recovered in better than 90% yield.¹²

These observations indicate that **3a** dissociates into two molecules of the (iminomethylidene)phosphine **4a** on gas-phase pyrolysis. The monomeric **4a** reverts to **3a** above -55 °C in the solid state. The strong band at 1853 cm^{-1} in the IR of **4a** is assigned to the asymmetric stretching vibration of the P=C=N moiety. For comparison, carbodiimides (RN=C=NR) absorb near 2100 cm^{-1} .

The analogous pyrolysis of **3b** (mp 220 °C) at 480 °C (sublimation temperature 110 °C, increasing to 200 °C in 35 min) gave rise to a strong IR band at 1839 cm^{-1} ascribed to **4b**. This material started redimerizing to **3b** at -25 °C, a process that was complete at 0 °C. Comparable results were obtained with the precursors **3c-e**.

The IR assignments were corroborated by using the mass spectrometry reactor. With increasing pyrolysis temperature, the

M^+ peak due to **3a** decreased, while that of **4a** increased. The CID mass spectrum of **4a** obtained at 430 °C is shown in Figure 2. The base peak at m/z 183 may be ascribed to the dibenzophospholyl cation, $C_{12}H_8P^+$, which is typical of diphenylated phosphorus compounds.¹³ In the pyrolysis of **3b** the parent peak of the precursor (m/z 382) disappeared at 490 °C, while that of **4b** (m/z 191) reached maximum intensity. Here, the spectrum is dominated by a loss of isobutene to give m/z 135, formally corresponding to $PhN=C=Ph^+$ or $PhNH-C\equiv P^+$.

We are continuing the studies of (iminomethylidene)phosphines, in particular cycloaddition reactions and attempts to obtain other phosphorus-containing cumulenes, e.g., $RP=C=O$.

Registry No. **1a**, 24103-42-2; **1b**, 87218-80-2; **1c**, 87729-47-3; **1d**, 87729-48-4; **1e**, 87729-49-5; **2b**, 87729-50-8; **2e**, 87729-51-9; **3a**, 87729-52-0; **3b**, 87218-81-3; **3c**, 87729-53-1; **3d**, 87729-54-2; **3e**, 87729-55-3; **4a**, 87729-56-4; **4b**, 87218-77-7; **4c**, 87729-57-5; **4d**, 87729-58-6; **4e**, 87729-59-7.

(13) Budzikiewicz, H.; Djerassi, C.; Williams, D. H. "Mass Spectrometry of Organic Compounds"; Holden-Day: San Francisco, 1967; p 647.

Enzyme System Generation of Singlet ($^1\Delta_g$) Molecular Oxygen Observed Directly by 1.0–1.8- μm Luminescence Spectroscopy

Ahsan U. Khan

Institute of Molecular Biophysics
and Department of Chemistry
Florida State University
Tallahassee, Florida 32306

Received July 25, 1983

Since the discovery of the chemical generation of singlet oxygen,^{1,2} numerous attempts have been made to demonstrate the generation of singlet oxygen in biological systems. Using our ultrasensitive 1.2- μm region spectrometer, we now report the observation of a strong singlet Δ oxygen luminescence emission in the IR (Figure 1) produced in the decomposition of hydrogen peroxide by the enzymes *lactoperoxidase*, *catalase*, and *chloroperoxidase*. Preliminary reports on kinetic studies of lactoperoxidase-generated singlet oxygen³ and a spectroscopic study of chloroperoxidase-generated singlet oxygen⁴ have been given recently.

The *lactoperoxidase* system consists of a 0.01 $\mu\text{M}/\text{mL}$ solution of lactoperoxidase (from milk, lyophilized; Sigma) in 0.01 M acetate buffer at pH + pD of 4.5 and 0.80 M KBr as a cofactor (Mallinckrodt, Analytical Reagent) at room temperature. The solvent is a 1:1 ratio of H_2O and D_2O . The experimental conditions are chosen to optimize singlet molecular oxygen emission; see the chemical scavenger studies of singlet molecular oxygen by Piatt et al.⁵ and by Rosen and Klebanoff.⁶ The enzyme and the hydrogen peroxide solutions are mixed in a 1:1 ratio under argon pressure into an optical cell with an overflow.

The luminescence emission spectrum of the lactoperoxidase/ H_2O_2 consists of a single emission band (Figure 1A) with a peak at 1.28 μm (1280 nm), and a full bandwidth at half-maximum of 251 cm^{-1} . There is also the suggestion of a broad underlying band which will not be discussed further here. Other singlet molecular oxygen luminescence sources have been characterized

(10) Becker, G.; Uhl, G., unpublished results.

(11) Maquestiau, A.; Van Haverbeke, Y.; De Meyer, C.; Flammang, R.; Perlaux, J. *Bull. Soc. Chim. Belg.* 1976, 85, 69.

(12) The band at 2120 cm^{-1} in Figure 1 is due to phenyl isocyanide, formed in a competing thermal fragmentation of **3a**. This material evaporates during warm-up and thus does not contaminate the final product. A yield of ca. 10% of phenyl isocyanide was obtained by distilling it into a cold trap and subsequently identifying it by comparison with an authentic sample.

(1) Khan, A. U.; Kasha, M. *J. Chem. Phys.* 1963, 39, 2105.

(2) Cf.: "Singlet Molecular Oxygen"; Schaap, A. P., Ed.; Dowden, Hutchinson and Rose: Stroudsburg, PA, 1976.

(3) Kanofsky, J. R. *J. Biol. Chem.* 1983, 258, 5991.

(4) Khan, A. U.; Gebauer, P.; Hager, L. P. *Proc. Natl. Acad. Sci. U.S.A.* 1983, 80, 5195.

(5) Piatt, J. F.; Cheema, A. S.; O'Brien, P. J. *FEBS Lett.* 1977, 74, 251.

(6) Rosen, H.; Klebanoff, S. J. *J. Biol. Chem.* 1977, 252, 4803.