

reactions. Extraneous signals in the η^5 -(CH₃)₅C₅ region indicate that competing thermolysis of **2**,^{17a} which likely involves ring methyl metalation,^{4b,13c,17b} is also operative. Interestingly, the reaction of **2** with CD₄ is significantly slower than with CH₄ and a preliminary analysis yields a kinetic isotope effect of 6 ± 2 .¹⁸ This result argues that H₃C-H bond breaking is the rate-limiting step in methane activation. That hydrolysis of the CD₄ reaction product produces pentamethylcyclopentadiene with <3% H(C-H₃)₄(CH₂D)C₅ by GC/MS indicates that CD₄ attack on a ring-metalated species is not a major reaction pathway. That the neopentane produced upon hydrolysis contains ca. $3 \pm 1\%$ neopentane-*d*₁¹⁹ suggests that eq 4 may be reversible.

These results demonstrate that it is possible to design isolable organoactinides of sufficiently high energy content that the stoichiometric (as opposed to catalytic) activation (with some selectivity) of saturated hydrocarbon molecules becomes thermodynamically favorable. Moreover and perhaps most fascinating, such activations are kinetically rather facile.

Acknowledgment. We thank the National Science Foundation (Grants CHE-8009060 and CHE-8306255) for generous support of this research. We thank G. M. Smith for sharing data on Cp₂Th[CH₂C(CH₃)₃][CH₂Si(CH₃)₃] and Dr. P. L. Watson for information in advance of publication.

(17) (a) Verified by independent thermolysis of **2** in C₆D₁₂. (b) For example, thermolysis of **2** in C₆D₁₂ followed by D₂O quenching results in appreciable quantities of D(CH₃)₄(CH₂D)C₅.^{13c}

(18) A highly accurate measurement is precluded by the considerably greater participation of the side reactions under these conditions.

(19) In addition to the expected neopentane-*d*₁, substantial quantities of neopentane-*d*₀ are also detected. Such a result is expected from the competing thermolysis of **2**, which produces a ring-metalated neopentyl compound.^{13c,17}

Synthesis of a Stable Cyclopropene Fused by a Six-Membered Ring. A Novel Approach from Fused Thiirene Sulfoxide

Wataru Ando,* Yukio Hanyu, and Toshikazu Takata


Department of Chemistry, The University of Tsukuba
Sakura-mura, Ibaraki 305, Japan

Katsuhiko Ueno

Research Institute for Polymers and Textile
Yatabe-cho, Ibaraki 305, Japan

Received December 21, 1983

Considerable attention has recently been focused on strained bicyclic alkenes, especially the [n.1.0]bicyclic system **1** (*n* = 1-4).¹



Although olefins should have a planar structure unless geometrical constraints^{2a,b} are present or the appropriate symmetry^{2c} is lacking, Wagner^{3a} and Pople^{3b} have shown by ab initio calculations that

(1) Greenberg, A.; Liebman, J. F. "Strained Organic Molecules"; Academic Press: New York, 1978; pp 97-98, 348.

(2) (a) Greenhouse, R.; Borden, W. T.; Hirotsu, K.; Clardy, J. *J. Am. Chem. Soc.* **1977**, *99*, 1664. (b) Review: Liebman, J. F.; Greenberg, A. *Chem. Rev.* **1976**, *76*, 311. (c) Mislow, K.; Benjamin, W. A. "Introduction to Stereochemistry"; Academic Press: New York, 1966; pp 11-13.





Figure 1. ORTEP drawing of the molecular structure of bicyclo[4.1.0]hex-1(6)-ene **8**. Bond lengths (Å): C₁-C₂ 1.521, C₁-C₃ 1.526, C₂-C₃ 1.286, C₂-C₄ 1.491, C₃-C₅ 1.478. Bond angles (°): C₂-C₁-C₃ 49.92, C₁-C₂-C₃ 64.82, C₁-C₃-C₂ 65.26, C₂-C₃-C₅ 130.23, C₃-C₂-C₄ 129.16.

all bicyclic alkenes **1** should have nonplanar structures. Considerable efforts have been directed toward the preparation of bicyclic alkenes.^{1,4} Evidence for intermediacy of bicyclo[4.1.0]alkene⁵ **1d**,^{6,7} as well as **1a**,⁸ **1b**,⁹ and **1c**,⁷ produced by dehydrohalogenation and dehalogenation of the corresponding halides, was presented as trapped Diels-Alder adducts by Gassman⁶ and Wiberg.⁷ The dimethyl derivative **2** was spectroscopically identified below -35 °C by Closs and Boll.¹⁰ However, experimental structural studies have never been reported for **1** owing to their instability.

We wish to report an attractive route to a thermally stable bicyclo[4.1.0]hept-1(6)-ene derivative via [2 + 3] cycloaddition of fused thiirene sulfoxide **3** with diphenyldiazomethane and its X-ray crystal analysis.

Fused thiirene sulfoxide **3**, recently prepared in our laboratory, was shown to be quite reactive.¹¹ It was treated with a 5-fold amount of diphenyldiazomethane in CH₂Cl₂ at room temperature. After standing for several hours, 3*H*-pyrazole **5** (54%)^{12,13} was obtained by chromatographic purification.



Photolysis (≥ 365 nm)¹⁴ of **5** in benzene at room temperature

(3) (a) Wagner, H.-U.; Szeimies, G.; Chandrasekhar, J.; Schleyer, P. v. R.; Pople, J. A.; Binkley, J. S. *J. Am. Chem. Soc.* **1978**, *100*, 1210. (b) Hehre, W. J.; Pople, J. A. *Ibid.* **1975**, *97*, 6941.

(4) Spanget-Larsen, J.; Gleiter, R. *Tetrahedron* **1983**, *39*, 3345.

(5) As a special case, benzocyclopropenes are stable compounds; see a review: Billups, W. E. *Acc. Chem. Res.* **1978**, *11*, 245.

(6) (a) Gassman, P. G.; Valcho, J. J.; Proehl, G. S. *J. Am. Chem. Soc.* **1979**, *101*, 231. (b) Gassman, P. G.; Valcho, J. J.; Proehl, G. S.; Cooper, C. F. *Ibid.* **1980**, *102*, 6519.

(7) Wiberg, K. B.; Bonneville, G. *Tetrahedron Lett.* **1982**, *23*, 5385.

(8) Szeimies, G.; Harnisch, J.; Baumgartl, O. *J. Am. Chem. Soc.* **1977**, *99*, 5183. Szeimies-Seebach, U.; Szeimies, G. *Ibid.* **1978**, *100*, 3966.


(9) Harnisch, J.; Baumgartl, O.; Szeimies, G.; Van Meerssche, M.; Germain, G.; Declercq, J.-P. *J. Am. Chem. Soc.* **1979**, *101*, 3370.

(10) Closs, G. L.; Boll, W. A.; Heyn, H.; Dev, V. *J. Am. Chem. Soc.* **1968**, *90*, 173.

(11) (a) Ando, W.; Hanyu, Y.; Takata, T.; Ueno, K. *J. Am. Chem. Soc.* **1982**, *104*, 4981. (b) Ando, W.; Hanyu, Y.; Takata, T.; Sakurai, T.; Kobayashi, K. *Tetrahedron Lett.*, in press.

(12) Spectral and elemental analysis data of **5**: mp 142.0-143.0 °C from hexane-benzene; ¹H NMR δ (CDCl₃) 1.34 (s, 6 H), 2.08 (s, 6 H), 3.10 (s, 3 H), 7.64-7.35 (m, 10 H); ¹³C NMR δ (CDCl₃) 157.4, 153.7, 153.4, 150.2, 133.3, 129.0, 128.3, 107.2, 62.4, 59.1, 25.1, 25.0, 23.6; UV (nm, hexane) λ_{max} 356 (ϵ 203) (N=N); MS, *m/e* 415 (M⁺). Anal. Calcd for C₂₅H₂₅N₃O₂: C, 69.40; H, 6.02; N, 16.87. Found: C, 69.22; H, 6.15; N, 16.79.

resulted in the quantitative release of nitrogen and gave a 63% yield of the fused cyclopropene **8**¹⁵ in addition to a 37% yield of the indene **9**.^{16,17}



The desired product **8** was separated by recrystallization from hexane-benzene at room temperature. Surprisingly, the fused cyclopropene **8** is a stable colorless crystalline compound (mp 143.5-144.0 °C). Weak absorption, presumably due to C-C double bond of the cyclopropene ring, appeared at 1805 cm⁻¹ (shoulder).¹⁵ The structure of **8** was determined by X-ray crystal structure analysis (Figure 1).¹⁸

Inspection of the X-ray crystal analysis data revealed the nonplanar structure of **8**, and the angle between the two rings was 162.4°, which is 7.4° larger than that estimated for **1c**.^{3a,19} The bond angles of C₃-C₂-C₄ and C₂-C₃-C₅ (129.2° and 130.2°, respectively) are ca. 20° smaller than those in nonfused cyclopropene (149.9°),²⁰ as expected. The ¹H NMR spectrum¹¹ of **5** shows resonances at 1.75 (s, CH₃ × 4), 3.03 (s, NCH₃), and 7.24 ppm (s, Ph) characteristic of its C₂ symmetry. The ¹³C NMR spectrum indicates that the olefinic carbon resonance appears at 155.5 ppm.¹⁵ Surprisingly, this value is ca. 30 ppm lower than those found for tetramethyl- (118.9 ppm) and 3,3-dimethylcyclopropenes (125.0 ppm) etc.²¹ This large downfield shift should be due to the highly strained framework of **8** by ring fusion, as found in the case of the thiirene sulfoxide **3**.^{11a}

The stability of **8** is clearly dependent on the α-substitution of tetramethyl groups, which might fix the bicyclic ring system.

(13) Compound **3** reacted with ethyl diazoacetate to afford 2*H*-pyrazole, via an additional 1,3-H shift, as reported in the reaction of diphenyl thiirene sulfoxide with phenyldiazomethane: Carpino, L. A.; Chen, H.-W. *J. Am. Chem. Soc.* **1979**, *101*, 390.

(14) A methanol solution of phenanthrene (5 g/L) was used as a filtered solution (path length 1 cm).

(15) **8**: colorless crystals, mp 143.5-144.0 °C from hexane-benzene; ¹H NMR δ (CDCl₃) 7.24 (brs, 10 H), 3.03 (s, 3 H), 1.75 (s, 12 H); ¹³C NMR δ (CDCl₃) 153.0, 144.2, 135.4, 128.2, 128.0, 126.6, 63.3, 58.4, 24.6, 24.3; IR (cm⁻¹, KBr) 1805 (ν_{C=C}); MS, *m/e* 387 (M⁺). Anal. Calcd for C₂₅H₂₅N₃O₂: C, 74.39; H, 6.50; N, 10.84. Found: C, 74.31; H, 6.44; N, 10.92.

(16) **9**: oil, ¹H NMR δ (CDCl₃) 7.28-7.66 (s, 9 H), 3.98 (s, 1 H), 3.06 (s, 3 H), 1.99 (s, 3 H), 1.93 (s, 3 H), 1.28 (s, 3 H), 1.09 (s, 3 H); ¹³C NMR δ (CDCl₃) 155.3, 152.7, 148.4, 145.9, 139.9, 138.7, 134.9, 129.7, 128.6, 127.8, 127.7, 125.3, 123.5, 120.8, 77.3, 65.3, 62.7, 56.2, 28.1, 24.9, 21.2, 21.0; MS, *m/e* 387 (M⁺).

(17) The formation of **8** and **9** is explained by the common intermediate, vinyl carbene **7**, derived from the diazo compound **6**. During the photoirradiation the reaction solution initially became light red, which corresponded to **6**, and then N₂ gas was evolved. Indene **9** may be produced from **8** via the known photoequilibrium between **7** and **8**: Halton, B.; Kulig, M.; Battiste, M. A.; Perreten, J.; Gibson, D. M.; Griffin, G. W. *J. Am. Chem. Soc.* **1971**, *93*, 2327.

(18) The crystal has monoclinic space group *p*2₁/*c* with *a* = 10.381 (2) Å, *b* = 8.719 (1) Å, *c* = 23.717 (5) Å, and β = 98.30 (2)° with *Z* = 4. Intensity data were collected on a four circle diffractometer with graphite monochromated Cu Kα radiation (3° < θ < 120°). 3712 unique reflections measured of which 2670 had intensities greater than 3σ |F₀| and were used for structure analysis. The structure was refined to a value of 0.086. For the detailed crystallographic data, supplementary material is available.

(19) Unfortunately, the angle for **1d** is not calculated,^{3a} but is considered to be slightly larger than 155° given for **1c**.^{3a}

(20) Kasai, P. H.; Meyers, R. J.; Eggers, D. F., Jr.; Wiberg, K. B. *J. Chem. Phys.* **1959**, *30*, 512.

(21) Bachbuch, M.; Grishin, Y. K.; Formanovskii, A. A. *Dokl. Acad. Nauk SSSR*, **1978**, *243*, 1171.

Above all, the nonplanar structure of bicyclo[4.1.0]alkene suggested by the calculations is confirmed by the present result.

Supplementary Material Available: Listings of atomic positional and thermal parameters, bond lengths, and bond angles for compound **8** (18 pages). Ordering information is given on any current masthead page.

Fourier Transform Infrared Photoacoustic Spectroscopy: A Novel Conformational Probe. Demonstration of α-Helical Conformation of Poly(γ-benzyl glutamate)

V. Renugopalakrishnan*

Laboratory for the Study of Skeletal Disorders and Rehabilitation
Childrens Hospital Medical Center
Harvard Medical School, Boston, Massachusetts 02115

Rajendra S. Bhatnagar

Laboratory of Connective Tissue Biochemistry
604-HSW, School of Dentistry
University of California
San Francisco, California 94143

Received November 7, 1983

Fourier transform infrared photoacoustic spectroscopy (FT-IR PAS) has emerged as a novel technique for studying a wide range of problems in chemistry and biology.¹ The determination of secondary structures of biopolymers from observed vibrational frequencies is one of the long-range goals of molecular spectroscopy.² Biopolymers, in general, have been difficult to be investigated by conventional infrared spectroscopy due to difficulties in uniformly dispersing them into an alkali halide matrix. Incorporation into alkali halide matrix may result in structural alterations during the pelleting process³ and hydration of the sample as well. FT-IR PAS offers an alternative method to investigate biopolymers per se in less than milligram quantities by totally eliminating artifactual effects introduced by incorporation into alkali halide matrix. FT-IR PAS represents a major advance in infrared spectroscopy which has not been extensively utilized in chemistry and biology. In this communication, the first report in the literature of an application of FT-IR PAS to the determination of molecular conformations, we are presenting the results of the application of this novel conformational probe to poly(γ-benzyl glutamate). Poly(γ-benzyl glutamate) has been shown to prefer α-helical structure in its higher molecular weight fractions by X-ray diffraction,⁴ conventional IR,⁵ and Raman⁶ spectroscopic methods. Abe and Krimm⁷ and Nevskaya and

* Current address: Laboratory of Skeletal Disorders and Rehabilitation, Orthopaedic Research, Enders-1220, Childrens Hospital Medical Center, Boston, MA 02115.

(1) (a) Vidrine, D. W. *Appl. Spectrosc.* **1980**, *34*, 314. (b) Krishnan, K. *Ibid.* **1981**, *35*, 549. (c) Kinney, J. B.; Staley, R. H.; Reichel, C. L.; Wrighton, M. S. *J. Am. Chem. Soc.* **1981**, *103*, 4273. (d) MacClelland, J. F. *Anal. Chem.* **1983**, *55*, 89A. (e) Rockley, M. G.; Davies, D. M.; Richardson, H. H. *Science (Washington, D.C.)* **1980**, *210*, 918.

(2) (a) Lord, R. C. *Appl. Spectrosc.* **1977**, *31*, 187. (b) Tu, A. T. "Raman Spectroscopy in Biology: Principles and Applications"; Wiley: New York, 1982. (c) Krimm, S. *Biopolymers* **1983**, *22*, 217.

(3) (a) Baker, A. W. *J. Phys. Chem.* **1957**, *61*, 450. (b) Milkey, R. G. *Anal. Chem.* **1958**, *30*, 1931. (c) for structural alterations of alkali halide matrix at high pressures, see: Knittle, E.; Jeanloz, R. *Science (Washington, D.C.)* **1984**, *223*, 53.

(4) Elliot, A.; Fraser, R. D. B.; McRae, T. P. *J. Mol. Biol.* **1965**, *11*, 821.

(5) (a) Miyawazawa, T.; Blout, E. R. *J. Am. Chem. Soc.* **1961**, *83*, 712. (b) Tsuboi, M. *J. Polym. Sci.* **1962**, *59*, 139. (c) Tomita, K.; Rich, A.; de Loze, C.; Blout, E. R. *J. Mol. Biol.* **1962**, *4*, 83. (d) Masuda, Y.; Miyazawa, T. *Makromol. Chem.* **1967**, *103*, 261.

(6) (a) Koenig, J. L.; Sutton, P. L. *Biopolymers* **1971**, *10*, 89. (b) Chen, M. C.; Lord, R. C. *J. Am. Chem. Soc.* **1974**, *96*, 4750. (c) Fasman, G. D.; Itoh, K.; Liu, C. S.; Lord, R. C. *Biopolymers* **1978**, *17*, 1729.