

**Figure 3.** Cyclic voltammograms of  $4^+$  (—) and  $\text{Cu}(7)_2^+$  (-----) in DMF;  $0.1 \text{ M } \text{N}(\text{C}_4\text{H}_9)_4^+\text{ClO}_4^-$ ; room temperature; mercury cathode; scan rate =  $10 \text{ mV/s}$ .

preparation of a formally copper(0) complex of exceptional stability.

**Acknowledgment** is made to the Centre National de la Recherche Scientifique for financial support. We also thank Dr. J. J. André and M. Bernard for ESR measurements.

### Formation of Ferryltetraphenylporphyrin by Laser Irradiation

Krzysztof Bajdor and Kazuo Nakamoto\*

Todd Wehr Chemistry Building, Marquette University  
Milwaukee, Wisconsin 53233

Received February 17, 1984


The crucial step in the reaction cycle of cytochrome P-450 is the formation of the ferryl group via the cleavage of dioxygen bonded to iron protoporphyrin. The main purpose of this communication is to report the first observation of the resonance Raman (RR) spectra of ferryltetraphenylporphyrin,  $\text{OFe}(\text{TPP})$ , which was formed unexpectedly during our measurement of the RR spectra of  $\text{Fe}(\text{TPP})\text{O}_2$  in pure  $\text{O}_2$  matrices at  $\sim 15 \text{ K}$ .

A stable, "base-bound" complex,  $\text{Fe}(\text{TPP})(\text{pip})_2$  (pip = piperidine), was placed in a miniature oven under the cold tip of our matrix isolation Raman apparatus.<sup>2</sup> The oven was heated at  $\sim 180 \text{ }^\circ\text{C}$  under vacuum until the vacuum gauge indicated complete dissociation of the base from the complex. The "base-free"  $\text{Fe}(\text{TPP})$  thus obtained was vaporized by heating the oven at  $\sim 225 \text{ }^\circ\text{C}$  by laser beam and cocondensed with pure oxygen on the inclined surface of the cold tip which was cooled to  $\sim 15 \text{ K}$  by a CTI Model 21 closed-cycle helium refrigerator. As our previous matrix-isolation IR studies<sup>3</sup> indicate, such cocondensation reactions produce five-coordinate, "base-free"  $\text{Fe}(\text{TPP})\text{O}_2$ . Resonance Raman spectra of the cocondensation


(1) For example, see: Alexander, L. S.; Goff, H. M. *J. Chem. Educ.* **1982**, *59*, 179.

(2) Scheuermann, W.; Nakamoto, K. *Appl. Spectrosc.* **1978**, *32*, 251, 302.

(3) Nakamoto, K.; Watanabe, T.; Ama, T.; Urban, M. W. *J. Am. Chem. Soc.* **1982**, *104*, 3744. Watanabe, T.; Ama, T.; Nakamoto, K. *J. Phys. Chem.* **1984**, *88*, 440.



**Figure 1.** Resonance Raman spectra of  $\text{Fe}(\text{TPP})$  cocondensed with dioxygen at  $\sim 15 \text{ K}$  ( $406.7\text{-nm}$  excitation,  $1\text{--}2 \text{ mW}$ ). (A)  $^{56}\text{Fe}(\text{TPP})$  with  $^{16}\text{O}_2$  (solid line) and  $^{54}\text{Fe}(\text{TPP})$  with  $^{16}\text{O}_2$  (broken line). (B)  $^{56}\text{Fe}(\text{TPP})$  with  $^{18}\text{O}_2$  (solid line) and  $^{54}\text{Fe}(\text{TPP})$  with  $^{18}\text{O}_2$  (broken line). (C)  $^{56}\text{Fe}(\text{TPP})$  with isotopically mixed dioxygen ( $^{16}\text{O}_2/^{16}\text{O}^{18}\text{O}/^{18}\text{O}_2 = 1/2/1$ ). The solid and broken lines indicate that spectra after 20 and 3 min of laser irradiation, respectively.



**Figure 2.** A plot of relative intensity of the  $852\text{-cm}^{-1}$  band of  $^{16}\text{O}^{56}\text{Fe}(\text{TPP})$  vs. time of laser irradiation.

products were measured using a Spex Model 1401 double monochromator in conjunction with a Spectra-Physics Model 164-01 Kr ion laser. Throughout this investigation, the samples were excited with  $1\text{--}2 \text{ mW}$  of the  $406.7\text{-nm}$  radiation (Soret excitation).

The solid line of Figure 1A shows the RR spectrum of  $^{56}\text{Fe}(\text{TPP})$  ( $^{56}\text{Fe}$ : Fe in natural abundance, 92% pure  $^{56}\text{Fe}$ ) cocondensed with  $^{16}\text{O}_2$ . It shows a strong signal at  $852 \text{ cm}^{-1}$  in addition to the normal  $\text{Fe}(\text{TPP})$  bands. The intensity of this band is time dependent, reaching the maximum after about 20 min. This intensity remains constant as long as the matrix temperature is kept at  $\sim 15 \text{ K}$ . The intensities of all other bands remain almost unchanged during this period. A plot of the relative intensity of the  $852\text{-cm}^{-1}$  band vs. the time of laser irradiation is shown in Figure 2. It is clear that the above photolysis follows the first-order kinetics.

The  $852\text{-cm}^{-1}$  band was shifted to  $818 \text{ cm}^{-1}$  in an  $^{18}\text{O}_2$  matrix (Figure 1, trace B). Similar experiments with an isotopically scrambled oxygen ( $^{16}\text{O}_2/^{16}\text{O}^{18}\text{O}/^{18}\text{O}_2 = 1/2/1$ ) produced two bands at  $852$  and  $818 \text{ cm}^{-1}$  as shown in trace C. (The broken line in trace C indicates the spectrum obtained after 3 min of laser irradiation). These results indicate quite clearly that the  $852\text{-cm}^{-1}$  and  $818\text{-cm}^{-1}$  bands are due to the  $\nu(^{16}\text{O}\text{--Fe})$  and  $\nu(^{18}\text{O}\text{--Fe})$ ,

respectively, of ferryltetraphenylporphyrin,  $\text{OFe}(\text{TPP})$ , which is formed by the cleavage of dioxygen of  $\text{Fe}(\text{TPP})\text{O}_2$  via laser photolysis. The observed isotopic shift ( $852 - 818 = 34 \text{ cm}^{-1}$ ) is in good agreement with that expected for a perturbed  $\text{FeO}$  molecule ( $38 \text{ cm}^{-1}$ ). These bands cannot be attributed to the  $\mu$ -oxo dimer,  $(\text{Fe}(\text{TPP}))_2\text{O}$ , since no strong Raman bands are seen near  $360 \text{ cm}^{-1}$ .<sup>4</sup>

Further support for our conclusion is provided by  $^{54}\text{Fe}$ - $^{56}\text{Fe}$  isotope substitution. As are shown by the broken lines of Figure 1, A and B, the bands at  $852$  and  $818 \text{ cm}^{-1}$  of  $^{16}\text{O}^{54}\text{Fe}(\text{TPP})$  and  $^{18}\text{O}^{54}\text{Fe}(\text{TPP})$ , respectively, are shifted by  $4.0 \text{ cm}^{-1}$  to higher frequencies by the substitution, and these values are again in good agreement with those expected for a perturbed  $\text{FeO}$  molecule ( $3.5 \text{ cm}^{-1}$ ).

A simple diatomic approximation gives a force constant of  $5.32 \text{ mdyn}/\text{\AA}$  for the above ferryl group. This is much larger than that of the  $\text{Fe}-\text{O}$  bonds in  $(\text{Fe}(\text{TPP}))_2\text{O}$  ( $3.8 \text{ mdyn}/\text{\AA}$ )<sup>4</sup> and in oxyhemoglobin ( $3.09 \text{ mdyn}/\text{\AA}$ ).<sup>5,6</sup> In this respect, the formulas such as  $\text{PFe}^{\text{IV}}=\text{O}^{2-}$  or  $\text{PFe}^{\text{V}}=\text{O}^{2-}$  describe the ferryl group better than  $\text{PFe}^{\text{III}}-\text{O}^-$  or  $\text{PFe}^{\text{IV}}-\text{O}^-$  (P: porphyrin).

According to recent ab initio MO calculations,<sup>7</sup> the negative charge and polarization of the dioxygen greatly increase upon coordination to an iron porphyrin (i.e.,  $\text{Fe}-\text{O}_1(-0.46e)-\text{O}_2(-0.19e)$ ). This trend will be accelerated by the donation of the second electron from NADH to the iron center and the presence of a cysteinyl sulfur ( $\text{S}^-$ ) at the trans position to the dioxygen in cytochrome P-450. It is then not surprising that the  $\text{O}-\text{O}$  bond cleavage occurs quite easily under biological conditions. We are now conducting experiments to answer the question of whether we can mimic the hydroxylation reaction of cytochrome P-450 in a matrix environment.

**Acknowledgment.** This work was supported by a National Science Foundation Grant (PCM-8114676).

(4) Burke, J. M.; Kincaid, J. R.; Spiro, T. G. *J. Am. Chem. Soc.* **1978**, *100*, 6077.

(5) Duff, L. L.; Appelman, E. H.; Shriver, D. F.; Klotz, I. M. *Biochem. Biophys. Res. Commun.* **1979**, *90*, 1098.

(6) Duff-Weddell, L. L. Ph.D. Thesis, Northwestern University, Evanston, IL, 1981.


(7) Nozawa, T.; Hatano, M.; Nagashima, U.; Obara, S.; Kashiwagi, H. *Bull. Chem. Soc. Jpn.* **1983**, *56*, 1721.

### Unique, One-Step, Double Isomerization (2E,4Z $\rightleftharpoons$ 2Z,4E) of 6-Oxo-2,4-heptadienoic Acid Catalyzed by Maleylacetone Cis-Trans Isomerase

Anthony L. Feliu, Karl J. Smith, and Stanley Seltzer\*

Chemistry Department  
Brookhaven National Laboratory  
Upton, New York 11973  
Received January 3, 1984

Maleylacetone cis-trans isomerase, which catalyzes the reactions shown in eq 1,<sup>1</sup> requires glutathione (GSH) as a coenzyme.<sup>2</sup>




Previous studies indicate that GSH binds to the enzyme along the backbone of the tripeptide, pointing its SH group away from

(1) Seltzer, S. *Enzymes* **1973**, *6*, 381-406.


(2) Seltzer, S. *J. Biol. Chem.* **1973**, *248*, 215-222.

Scheme I<sup>a</sup>



<sup>a</sup> (a)  $\text{NaH}/\text{THF}$ ; (b)  $(\text{C}_6\text{H}_5)_3\text{P} = \text{CHCOCH}_3$ ; (c)  $(\text{C}_6\text{H}_5)_3\text{P} = \text{CHCO}_2\text{CH}_3/\text{THF}$ ; (d)  $t\text{-C}_4\text{H}_9\text{OH}/1 \text{ N NaOH}$  (1:10),  $0^\circ\text{C}$ , 4 min; (e)  $\text{I}_2/\text{THF}$ , reflux.

the enzyme's surface.<sup>3</sup> In the enzymatic and nonenzymatic reaction, reversible nucleophilic addition of GSH to C2 of **1** forming a dienediol intermediate (**3**) thereby allows internal rotation about the C2-C3 bond; ketonization with expulsion of GSH provides **2** (eq 2).<sup>4</sup>



To examine the structural requirements of the enzyme, 6-oxo-2,4-heptadienoic acids (**4**) and methyl esters **5** were synthesized (Scheme I) to provide the four possible cis-trans isomeric skeletons, only one of which was reported previously. Base-catalyzed hydrolysis of the esters **5-EZ** and **5-EE** yielded the corresponding acids. Repeated attempts to hydrolyze methyl 6-oxo-2(Z),4(Z)-heptadienoate (**5-ZZ**), however, led only to the **4-ZE** acid. The routes of synthesis (Scheme I) and the NMR spectra<sup>7</sup> establish the structures of these new compounds.

As expected, **4-ZE** acid is isomerized to **4-EE** by the enzyme and obligatory GSH.<sup>8</sup> The  $K_M$  for **4-ZE** is  $3.2 \times 10^{-3} \text{ M}$  vs.  $8.4 \times 10^{-4} \text{ M}$  for maleylacetone;<sup>3,9</sup>  $k_{\text{cat}}$  is  $\sim 0.36$  times that for maleylacetone.

(3) Morrison, W. S.; Wong, G.; Seltzer, S. *Biochemistry* **1976**, *15*, 4228-4233.

(4) Seltzer, S.; Lin, M. *J. Am. Chem. Soc.* **1979**, *101*, 3091-3097.

(5) Doerr, I. L.; Willette, R. E. *J. Org. Chem.* **1973**, *38*, 3878.

(6) Burness, D. M. "Organic Syntheses"; Wiley: New York, 1973; Collect. Vol. V, p 403. Fakstorp, J.; Raleigh, D.; Schniepp, L. E. *J. Am. Chem. Soc.* **1950**, *72*, 869. Clauson-Kaas, H.; Limborg, F. *Acta Chem. Scand.* **1947**, *1*, 619.

(7) See supplementary material.

(8) A highly purified preparation of the vibrio 01 maleylacetone isomerase (specific activity 23 units/mg) was obtained by affinity chromatography on GS-agarose.<sup>3</sup>

(9) The kinetics of isomerization were followed by HPLC analysis on a C-18 column (1% acetic acid, 5% acetonitrile, 94% water or 2.5 mM tetra-*n*-butyl ammonium phosphate in 93.3% 0.01 M phosphate buffer, pH 7.4, 6.7% acetonitrile).