

to calculate maximum and minimum measurements of an enriched abundance, i.e., (M + 2) normalized against 100% abundance for M and corrected for the natural abundance of (M + 2) in the mixture. Thus, for each conversion we had eventually a maximum and minimum measure of the enrichment in a sample. Maximum and minimum KIE were calculated from these data and are expressed as an average with its error for each run in the tables.

**Acknowledgment.** We thank Dr. Frank W. Crow, Midwest Center for Mass Spectrometry (an NSF Regional Facility), Department of Chemistry, University of Nebraska, Lincoln, NE, for mass spectrometric analyses of the isotope abundances in the commercial acetone and the synthesized, labeled 2,2'-dimethoxyazobenzenes prior to their use in the rearrangements.

**Registry No. 1,** 787-77-9; **2,** 119-90-4; **2** (bis(trifluoroacetyl) deriva-

tive), 92241-72-0; **2-<sup>15</sup>N<sub>2</sub>,** 92241-73-1; **3-<sup>15</sup>N<sub>2</sub>,** 92241-77-5; **3-<sup>13</sup>C<sub>2</sub>,** 92241-86-6; *trans*-**3-<sup>2</sup>H<sub>2</sub>,** 92241-92-4; *cis*-**3-<sup>2</sup>H<sub>2</sub>,** 92241-93-5; **4,** 85167-01-7; **5,** 92241-78-6; **6,** 92241-82-2; **7,** 92241-83-3; **8,** 92269-47-1; **8** (unlabeled), 37856-18-1; **9,** 5081-36-7; **9** (acid chloride), 67579-92-4; **10,** 92241-87-7; **11,** 16292-88-9; **12,** 92241-89-9; **13,** 92241-90-2; **14,** 92241-91-3; <sup>15</sup>N, 14390-96-6; <sup>13</sup>C, 14762-74-4; D<sub>2</sub>, 7782-39-0; AgClO<sub>4</sub>, 7783-93-9; PhOCOCl, 1885-14-9; Ag<sup>15</sup>NO<sub>3</sub>, 92241-74-2; *o*-HOC<sub>6</sub>H<sub>4</sub><sup>15</sup>NO<sub>2</sub>, 92241-75-3; *o*-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub><sup>15</sup>NO<sub>2</sub>, 92241-76-4; *o*-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub><sup>15</sup>NH<sub>2</sub>, 63792-03-0; (CH<sub>3</sub>)<sub>2</sub><sup>13</sup>CO, 3881-06-9; O<sub>2</sub>NC(CHO)<sub>2</sub>, 34460-99-6; *p*-HO[4-<sup>13</sup>C]C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>, 3881-07-0; *p*-HO[4-<sup>13</sup>C]C<sub>6</sub>H<sub>4</sub>NH<sub>2</sub>, 3881-08-1; [4-<sup>13</sup>C]C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub>, 55147-71-2; [4-<sup>13</sup>C]C<sub>6</sub>H<sub>5</sub>OH, 70211-36-8; *o*-CH<sub>3</sub>O[4-<sup>13</sup>C]C<sub>6</sub>H<sub>4</sub>NH<sub>3</sub>Cl, 92241-84-4; *o*-CH<sub>3</sub>O[4-<sup>13</sup>C]C<sub>6</sub>H<sub>4</sub>NH<sub>2</sub>, 92241-85-5; K<sup>15</sup>NO<sub>3</sub>, 57654-83-8; 2-[4-<sup>13</sup>C]phenyl-5-nitrobenzophenone, 92241-79-7; 2-([4-<sup>13</sup>C]-*o*-hydroxyphenyl)-5-nitrobenzophenone, 92241-80-0; 2-([4-<sup>13</sup>C]-*o*-methoxyphenyl)-5-nitrobenzophenone, 92241-81-1; 4-chloro-2-phenylquinazoline, 6484-25-9.

## Rates and Equilibria of the Reaction of 2,4,6-Triphenylthiopyrylium Ion with Piperidine and Morpholine in Me<sub>2</sub>SO. An Unusual Proton Transfer to a Nitrogen Base

Giancarlo Doddi\* and Gianfranco Ercolani\*

Contribution from the Centro C.N.R. di Studio sui Meccanismi di Reazione, c/o Dipartimento di Chimica, Università di Roma "La Sapienza", P.le Aldo Moro, 2-I-00185 Roma, Italy.

Received April 5, 1984. Revised Manuscript Received June 21, 1984

**Abstract:** The complete set of kinetic and equilibrium constants of the reaction of 2,4,6-triphenylthiopyrylium ion (**1**) with piperidine and morpholine has been obtained in Me<sub>2</sub>SO at 25 °C. The reaction involves the formation of both the corresponding 2*H*- and 4*H*-thiopyrans, which equilibrate to form only the more stable 2*H* adduct. The kinetic data are consistent with a two-step process wherein the proton transfer from the protonated 2*H*- and 4*H*-thiopyran intermediates to the amine is the rate-controlling step. The thermodynamically favored proton transfer to the neutral adducts by the solvated proton shows a rate below the diffusion limit, whereas the observed Brønsted coefficient would indicate a diffusion-controlled process. This behavior is discussed in terms of the Eigen mechanism. The factors affecting the nucleophilic addition are discussed, and a comparison is made with the previously reported reaction of **1** with primary amines.

The reaction between nucleophiles and organic cations to yield the corresponding attachment products is a fundamental process in organic chemistry, but in spite of its apparent simplicity the affecting factors are not yet fully understood.<sup>1</sup>


The heteroaromatic cations pyridinium, pyrylium, and thiopyrylium are well suited to obtain insights into these processes. In fact, these ions can be substituted on the two reactive positions, namely the C $\alpha$  and C $\gamma$  carbon atoms, with several substituents having different electronic and steric requirements; moreover, a large number of nucleophiles can react with these substrates making possible the study of the factors affecting both the reactivity and the positional selectivity.

Our interest for these reactions has prompted us to report on a detailed kinetic study of the reaction of 2,4,6-triphenylthiopyrylium ion (**1**) in Me<sub>2</sub>SO at 25 °C to yield the corresponding 2*H*- and 4*H*-thiopyrans with piperidine and morpholine, which have different nucleophilicities and identical steric requirements. The aim is to compare this reaction with the corresponding one of the butylamine and cyclohexylamine,<sup>2</sup> in order to study the effect of the structure of the amine on the addition reaction of such a reactive cation.

(1) (a) Ritchie, C. D. *Pure Appl. Chem.* **1978**, *50*, 1281. (b) Ritchie, C. D. *J. Am. Chem. Soc.* **1983**, *105*, 7313.

(2) Doddi, G.; Ercolani, G. *J. Org. Chem.* **1984**, *49*, 1806.

Scheme I



### Results

The kinetic study of the reaction of **1** with piperidine/piperidinium and morpholine/morpholinium buffers, carried out at 410 nm, where only **1** absorbs, and at 355 nm, where both **1** and the 2*H* adduct absorb (Figure 1), showed the presence of two relaxation processes, whose separation increased on increasing the amine/ammonium ratio. Whenever the condition  $\tau_1^{-1} \gg \tau_2^{-1}$  was fulfilled ( $\tau_1$  and  $\tau_2$  are the relaxation times of the first and the second process, respectively), the substrate completely disappeared when the first process was over. Similarly to the primary amine reactions,<sup>2</sup> monitoring the process at 355 nm, an initial decrease




Figure 1. UV-vis spectra of 1 (---) and the corresponding 2H adducts with piperidine (—) and morpholine (-·-) in Me<sub>2</sub>SO.




Figure 2. Plot of  $\tau_1^{-1}$  vs. [piperidine] at [piperidinium] = 0.1 M.

of absorbance followed by a slower increase up to a constant value was observed. This feature, combined with the NMR evidence,<sup>3</sup> indicates the complete formation of the 2H adduct.

This pattern can be rationalized by Scheme I, in which the first process is the formation of both 4H- and 2H-thiopyrans and the second one is equilibration to the more stable 2H adduct. The  $k_p$  and  $k_{-p}$  terms refer to proton transfer and are defined as  $k_p = k_p^s + k_p^{am}[am]$  and  $k_{-p} = k_{-p}^{sH}[sH] + k_{-p}^{amH}[amH]$ , in which  $k_p^s$  and  $k_p^{am}$  are the rate constants for the deprotonation of the charged adducts by solvent and amine, respectively, and  $k_{-p}^{sH}$  and  $k_{-p}^{amH}$  are the rate constants for the protonation of neutral adducts by the solvated proton and the ammonium ion, respectively.

Taking into account that the equilibrium is completely shifted toward the 2H adduct, the application of the steady-state approximation with respect to 4H<sup>+</sup> and 2H<sup>+</sup> to Scheme I, gives the following relationships:<sup>4</sup>

$$\tau_1^{-1} + \tau_2^{-1} = \frac{k_2[am]k_{2p}}{k_{-2} + k_{2p}} + \frac{k_4[am]k_{4p}}{k_{-4} + k_{4p}} + \frac{k_{-4}k_{-4p}}{k_{-4} + k_{4p}} \quad (1)$$

$$\tau_1^{-1}\tau_2^{-1} = \frac{k_2[am]k_{2p}}{k_{-2} + k_{2p}} \frac{k_{-4}k_{-4p}}{k_{-4} + k_{4p}} \quad (2)$$

**Piperidine. Study of the Two Relaxation Processes.** As mentioned above, when the reaction is carried out with buffers having  $[am] \approx 10^{-3}$ – $10^{-2}$  M and  $[amH] = 0.1$  M, the substrate fully




Figure 3. Plot of  $\tau_2^{-1}$  vs. [piperidine] at [piperidinium] = 0.1 M.

disappears and the condition  $\tau_1^{-1} \gg \tau_2^{-1}$  is fulfilled (Table S1, A, B; Figures 2 and 3).

Under these conditions an approximate expression of  $\tau_1^{-1}$  holds:

$$\tau_1^{-1} = \left( \frac{k_2k_{2p}}{k_{-2} + k_{2p}} + \frac{k_4k_{4p}}{k_{-4} + k_{4p}} \right) [am] \quad (3)$$

Introducing eq 3 into eq 2 affords:

$$\tau_2^{-1} = \left( \frac{k_2k_{2p}}{k_{-2} + k_{2p}} \frac{k_{-4}k_{-4p}}{k_{-4} + k_{4p}} \right) / \left( \frac{k_2k_{2p}}{k_{-2} + k_{2p}} + \frac{k_4k_{4p}}{k_{-4} + k_{4p}} \right) \quad (4)$$

A plot (not shown) of  $\tau_1^{-1}/[am]$  vs.  $[am]$  is linear at the lower amine concentrations, with an intercept that is negligible within the experimental errors, whereas at the higher ones a downward curvature is observed. This behavior indicates that at the higher amine concentrations one of the following relationships, at least, is fulfilled— $k_{-2} \approx k_{2p}$ ,  $k_{-4} \approx k_{4p}$ —whereas at the lower ones eq 5 holds:

$$k_{-2} \gg k_{2p} \text{ and } k_{-4} \gg k_{4p} \quad (5)$$

Moreover, since the intercept is negligible:

$$k_{2p}^{am} [am] \gg k_{2p}^s \text{ and } k_{4p}^{am} [am] \gg k_{4p}^s \quad (6)$$

and eq 3 becomes:

$$\tau_1^{-1} = \left( \frac{k_2k_{2p}^{am}}{k_{-2} + k_{2p}^{am}[am]} + \frac{k_4k_{4p}^{am}}{k_{-4} + k_{4p}^{am}[am]} \right) [am]^2 \quad (7)$$

that at the lower amine concentrations reduces to:

$$\tau_1^{-1} = \left( \frac{k_2k_{2p}^{am}}{k_{-2}} + \frac{k_4k_{4p}^{am}}{k_{-4}} \right) [am]^2 \quad (8)$$

Two sets of experiments carried out at two different amine concentrations with buffers having lower  $[am]/[amH]$  ratios than the previous ones, i.e., under conditions in which the third term of eq 1 cannot be neglected, give two parallel straight lines in the  $(\tau_1^{-1} + \tau_2^{-1})$  vs.  $[amH]$  plot (Table S1 C, D; Figure 4). This feature indicates that:

$$k_{-4p}^{amH}[amH] \gg k_{-4p}^{sH}[sH] \quad (9)$$

and since at these amine concentrations both eq 5 and eq 6 hold, eq 1 becomes

$$\tau_1^{-1} + \tau_2^{-1} = \left( \frac{k_2k_{2p}^{am}}{k_{-2}} + \frac{k_4k_{4p}^{am}}{k_{-4}} \right) [am]^2 + k_{-4p}^{amH} [amH] \quad (10)$$

(3) Cordischi, V. C.; Doddi, G.; Stegel, F. *J. Org. Chem.* **1982**, *47*, 3496.

(4) For an exhaustive treatment of the relaxation processes, see: Bernasconi, C. F. "Relaxation Kinetics"; Academic Press: New York, 1976.




Figure 4. Plot of  $(\tau_1^{-1} + \tau_2^{-1})$  vs. [piperidinium] at [piperidine] =  $3.51 \times 10^{-4}$  M (●) and [piperidine] =  $6.15 \times 10^{-4}$  M (○).

$k_{-4p}^{amH}$  being the slope of the plots of Figure 4.

A simplified expression of eq 4 can be written by taking account of eq 6 and 9:


$$\tau_2^{-1} = \left( \frac{k_2 k_{2p}^{am} / k_{-2} + k_{2p}^{am} [am]}{k_{-2} + k_{2p}^{am} [am]} \frac{k_{-4} k_{4p}^{amH} [amH]}{k_{-4} + k_{4p}^{amH} [amH]} \right) / \left( \frac{k_2 k_{2p}^{am}}{k_{-2} + k_{2p}^{am} [am]} + \frac{k_4 k_{4p}^{am}}{k_{-4} + k_{4p}^{am} [am]} \right) \quad (11)$$

The  $k_2$ ,  $k_4$ ,  $k_{2p}^{am}/k_{-2}$ , and  $k_{4p}^{am}/k_{-4}$  terms are evaluated by the  $\tau_1^{-1}$  and  $\tau_2^{-1}$  data by using an iterative method (see Appendix). We were compelled to use this method because of the need to have  $\tau_1^{-1}$  data at amine concentrations in which the reaction is too fast for the stopped flow method.

In Figures 2 and 3 are reported both the experimental values of  $\tau_1^{-1}$  and  $\tau_2^{-1}$  and the curves calculated by eq 7 and 11.

The  $k_{-2p}^{amH}$  term can be determined by the relationship  $k_{-2}^{amH} = k_2 k_{2p}^{am} / k_{-2} K_{2T}$ , in which  $K_{2T}$  is the equilibrium constant for the  $2H$  adduct formation from **1** (see Experimental Section). By the relationship  $K_{4T} = k_4 k_{4p}^{am} / k_{-4} k_{-4p}^{amH}$ ,  $K_{4T}$  can also be evaluated.

The  $k_{-2}$  and  $k_{-4}$  terms are experimentally inaccessible, but their estimation can be obtained by the relationships  $k_{-2} = k_2 K_2^{am} / K_{2T}$  and  $k_{-4} = k_4 K_4^{am} / K_{4T}$ , where  $K_2^{am} = k_{2p}^{am} / k_{-2p}^{amH}$  is the equilibrium constant for the proton transfer reaction between the  $2H^+$  adduct and the amine and  $K_4^{am} = k_{4p}^{am} / k_{-4p}^{amH}$  is the corresponding one for the  $4H^+$  adduct.  $K_2^{am}$  and  $K_4^{am}$  could be estimated by assuming an additive effect of the thiopyranil group on the acidity of the ammonium moiety of the charged adducts, i.e.,  $K_2^{but} = K_2^{pip}$  and  $K_4^{but} = K_4^{pip}$  ( $K_2^{but}$  and  $K_4^{but}$  are referred to the butylamine reaction and were determined in our previous work),<sup>2</sup> but probably the assumption that  $K_2^{pip}/K_2^{but}$  and  $K_4^{pip}/K_4^{but}$  coincide with  $K^{pip}/K^{but}$  (1.65) reported for the deprotonation of 3,6-bis(dimethylamino)xanthylum adduct (**2**) in  $Me_2SO$ <sup>5</sup> is a better one. The equilibrium constants for the formation of  $2H^+$  and  $4H^+$  adducts,  $K_2 = k_2/k_{-2}$ , and  $K_4 = k_4/k_{-4}$ , can now be evaluated.



**pH-Jump Experiments.** When a solution of the  $2H$  adduct  $\approx 10^{-5}$  M is subjected to a pH-jump by mixing it with a  $Me_2SO$





Figure 5. Plot of  $k_{obsd2}$  (●) and  $k_{obsd4}$  (○) vs. [*p*-toluenesulfonic acid].

solution of *p*-toluenesulfonic acid,<sup>6</sup> the complete formation of **1** is observed. Considering that the equilibrium is completely shifted toward **1** owing to the protonation of the forming amine, the reaction scheme becomes:



As shown in Figure 5, the  $k_{obsd2}$  is linearly related to the strong acid concentration (Table S2) with 0 intercept. The rate law is consistent with  $2H^+$  as a steady-state intermediate:  $k_{obsd2} = k_{-2} k_{-2p}^{sH} [sH] / (k_{-2} + k_{2p}^{sH})$ . Since  $k_{-2} \gg k_{2p}^{sH}$ , as inferred from relationship 5, the equation reduces to  $k_{obsd2} = k_{-2p}^{sH} [sH]$ , i.e., the protonation of the  $2H$ -thiopyran by the solvated proton is the rate-determining step. Therefore  $k_{-2p}^{sH}$  coincides with the slope of  $k_{obsd2}$  plot.

The  $k_{obsd4}$  term was determined by performing the pH-jump immediately after the solution of **1** and amine was prepared, before any significant equilibration between the two adducts had occurred. Only one relaxation process was detected, indicating that the decompositions of the two adducts to yield **1** occur at similar rates. As already reported,<sup>2</sup>  $k_{obsd4}$  can be obtained by the following equation:

$$\exp(-k_{obsd4}t) = \frac{\Delta OD - \Delta OD_2^\circ \exp(-k_{-2p}^{sH} [sH]t)}{\Delta OD_4^\circ} \quad (12)$$

Where  $\Delta OD_2^\circ$  and  $\Delta OD_4^\circ$  are the amplitudes related to the decomposition of the  $2H$  and  $4H$  adducts and are given by the following relationships:

$$\Delta OD_2^\circ = \Delta OD^\circ \left[ \frac{k_2 k_{2p}^{am}}{k_{-2}} / \left( \frac{k_2 k_{2p}^{am}}{k_{-2}} + \frac{k_4 k_{4p}^{am}}{k_{-4}} \right) \right] \quad (13)$$

$$\Delta OD_4^\circ = \Delta OD^\circ \left[ \frac{k_4 k_{4p}^{am}}{k_{-4}} / \left( \frac{k_2 k_{2p}^{am}}{k_{-2}} + \frac{k_4 k_{4p}^{am}}{k_{-4}} \right) \right] \quad (14)$$

$\Delta OD^\circ$  being the amplitude of the entire process.

As shown in Figure 5 and Table S2, also  $k_{obsd4}$  shows a linear dependence on the concentration of the acid, and therefore  $k_{-4p}^{sH}$  can be obtained by the corresponding slope.

The  $k_{2p}^{sH}$  and  $k_{4p}^{sH}$  terms can now be evaluated by the relationships  $k_{2p}^{sH}/k_{-2p}^{sH} = K_{2a} = K_a K_2^{am}$  and  $k_{4p}^{sH}/k_{-4p}^{sH} = K_{4a} = K_a K_4^{am}$ , where  $K_{2a}$  and  $K_{4a}$  are the acid dissociation constants of  $2H$  and  $4H$  adducts, respectively, and  $K_a$  is the acid dissociation constant of the conjugated acid of the amine.

(6) *p*-Toluenesulfonic acid is completely ionized in  $Me_2SO$  over the concentration range from  $10^{-4}$  to  $10^{-2}$  M: Ritchie, C. D.; Uschold, R. E. *J. Am. Chem. Soc.* **1967**, *89*, 1721.

**Table I.** Kinetic and Equilibrium Constants for the Reaction of **1** with Piperidine, Morpholine, Butylamine, and Cyclohexylamine in Me<sub>2</sub>SO at 25 °C<sup>a</sup>

	piperidine ( $K_a = 2.2 \times 10^{-11}$ M) <sup>b</sup>	morpholine ( $K_a = 1.15 \times 10^{-9}$ M)	butylamine <sup>c</sup> ( $K_a = 1.3 \times 10^{-11}$ M)	cyclohexylamine <sup>c</sup> ( $K_a = 1.3 \times 10^{-11}$ M)
$k_2$ , <sup>d</sup> s <sup>-1</sup> M <sup>-1</sup>	$4.1 \times 10^4$	$9.0 \times 10^3$ <sup>f</sup>	$3.7 \times 10^3$	$7.0 \times 10^2$
$k_{-2}$ , s <sup>-1</sup>	$1.7 \times 10^4$ <sup>e</sup>	$1.7 \times 10^5$ <sup>g</sup>	$5.4 \times 10^2$	$>7 \times 10^2$
$K_2$ , M <sup>-1</sup>	$2.5$ <sup>e</sup>	$5.2 \times 10^{-2}$ <sup>g</sup>	7.0	
$k_{2p}^{am}$ , s <sup>-1</sup> M <sup>-1</sup>	$7.2 \times 10^4$ <sup>e</sup>	$5.8 \times 10^4$ <sup>g</sup>		
$k_{-2p}^{amH}$ , s <sup>-1</sup> M <sup>-1</sup>	$1.9 \times 10^{-2}$	$1.0 \times 10^{-2}$		
$K_2^{am}$	$3.8 \times 10^6$ <sup>e</sup>	$5.8 \times 10^6$ <sup>g</sup>	$2.3 \times 10^6$	
$k_{2p}^s$ , s <sup>-1</sup>	$4.1$ <sup>e</sup>	$3.4 \times 10^2$ <sup>g</sup>		
$k_{-2p}^{sH}$ , s <sup>-1</sup> M <sup>-1</sup>	$4.9 \times 10^4$	$5.2 \times 10^4$		
$K_{2a}$ , M	$8.4 \times 10^{-5}$ <sup>e</sup>	$6.6 \times 10^{-3}$ <sup>g</sup>	$3.0 \times 10^{-5}$	
$K_{2T}$ , M <sup>-1</sup>	$9.5 \times 10^6$	$3.0 \times 10^5$	$1.6 \times 10^7$	$4.8 \times 10^6$
$k_4$ , s <sup>-1</sup> M <sup>-1</sup>	$1.1 \times 10^5$	$2.5 \times 10^4$ <sup>f</sup>	$1.4 \times 10^4$	$2.7 \times 10^3$
$k_{-4}$ , s <sup>-1</sup>	$2.9 \times 10^4$ <sup>e</sup>	$7.0 \times 10^5$ <sup>g</sup>	$6.5 \times 10^2$	$>7 \times 10^2$
$K_4$ , M <sup>-1</sup>	$4.0$ <sup>e</sup>	$3.6 \times 10^{-2}$ <sup>g</sup>	22	
$k_{4p}^{am}$ , s <sup>-1</sup> M <sup>-1</sup>	$5.0 \times 10^5$ <sup>e</sup>	$5.0 \times 10^5$ <sup>g</sup>		
$k_{-4p}^{amH}$ , s <sup>-1</sup> M <sup>-1</sup>	10	6.9		
$K_4^{am}$	$4.8 \times 10^4$ <sup>e</sup>	$7.2 \times 10^4$ <sup>g</sup>	$2.9 \times 10^4$	
$k_{4p}^s$ , s <sup>-1</sup>	$0.1$ <sup>e</sup>	$6.8$ <sup>g</sup>		
$k_{-4p}^{sH}$ , s <sup>-1</sup> M <sup>-1</sup>	$9.5 \times 10^4$	$8.1 \times 10^4$		
$K_{4a}$ , M	$1.1 \times 10^{-6}$ <sup>e</sup>	$8.3 \times 10^{-5}$ <sup>g</sup>	$4 \times 10^{-7}$	
$K_{4T}$ , M <sup>-1</sup>	$1.9 \times 10^5$	$2.6 \times 10^3$	$6 \times 10^5$	$2.5 \times 10^4$

<sup>a</sup>  $\mu = 0.1$  M. <sup>b</sup> From ref 25. <sup>c</sup> From ref 2. <sup>d</sup> Uncorrected for the statistical factor. <sup>e</sup> Estimated by the assumption  $K_i^{pip}/K_i^{but} = 1.65$ , as described in the text. <sup>f</sup> Estimated by the assumption  $(k_2/k_4)_{mor} = (k_2/k_4)_{pip}$ . <sup>g</sup> Estimated by the assumption  $K_i^{mor}/K_i^{but} = 2.50$ , as described in the text.

In Table I are collected the kinetic and equilibrium constants evaluated or estimated for the piperidine reaction, together with the corresponding data for the morpholine reaction. For a comparison in this table are reported also the data for the butylamine and cyclohexylamine reaction.<sup>2</sup>

**Morpholine.** The morpholine reaction behaves similarly to the piperidine one. At variance with this reaction the proton transfer step is rate controlling also at the higher accessible amine concentrations, as indicated from both the linear dependence of  $\tau_1^{-1}/[am]$  vs.  $[am]$  (with a negligible intercept within the experimental errors) and the independence of  $\tau_2^{-1}$  on the amine concentration, (Table S3A). Under these conditions eq 11 is reduced to:

$$\tau_2^{-1} = \left( \frac{k_2 k_{2p}^{am}}{k_{-2}} k_{-4p}^{amH} [amH] \right) / \left( \frac{k_2 k_{2p}^{am}}{k_{-2}} + \frac{k_4 k_{4p}^{am}}{k_{-4}} \right) \quad (15)$$

According to eq 8 the slope of the plot of  $\tau_1^{-1}/[am]$  vs.  $[am]$  coincides with  $([k_2 k_{2p}^{am}/k_{-2}] + [k_4 k_{4p}^{am}/k_{-4}])$ . Introducing into eq 15 this term and  $k_{-4p}^{amH}$  as obtained as previously reported for the piperidine reaction (Table S3 B, C),  $k_2 k_{2p}^{am}/k_{-2}$  and hence  $k_4 k_{4p}^{am}/k_{-4}$  can be evaluated.

The  $k_2$  and  $k_4$  terms were evaluated by experiments carried out in the presence of acetate buffers, in order to make the deprotonation step fast with respect to the nucleophilic attack. Plots of  $\tau_1^{-1}$  vs.  $[am]$  are linear with slopes that are independent of the concentrations of the acetate ion (Table S4). Under these conditions the slopes coincide with  $(k_2 + k_4)$ . With the assumption that the  $k_2/k_4$  ratio coincides with that observed for the piperidine reaction,  $k_2$  and  $k_4$  can now be evaluated.<sup>7</sup>

The equilibrium constants  $K_2^{mor}$  and  $K_4^{mor}$  were obtained as previously indicated for piperidine, by the ratio  $K^{mor}/K^{but} = 2.5$ .

The remaining kinetic and equilibrium constants reported in Table I are obtained as previously indicated for the piperidine reaction (Table S5).

## Discussion

**Nucleophilic Attack Step.** The order of reactivity for the attack on the  $\alpha$  and  $\gamma$  positions, piperidine > morpholine  $\approx$  butylamine > cyclohexylamine (Table I), is closely related to the one observed in many attachment reactions of amines to organic cations,<sup>5,8</sup> activated alkenes and benzenes,<sup>9</sup> and in nucleophilic aromatic

substitutions of heteroaromatic cations or neutral activated compounds.<sup>10</sup> The higher reactivity of secondary amines with respect to the less hindered primary ones indicates that severe steric strain to the approach of the nucleophile is not involved.<sup>11</sup> However, the lower reactivity of cyclohexylamine with respect to butylamine indicates that this effect is not quite negligible.

Two-point Brønsted plots of  $\log K_i$  ( $i = 2, 4$ ) vs.  $pK_a$  of the conjugated acid of the amines yield  $\beta_{eq}$ . Both  $\beta_{eq2}$  (0.98) and  $\beta_{eq4}$  (1.2) indicate that the difference of carbon basicity between piperidine and morpholine is comparable to their difference of proton basicity.

The  $\beta_{nuc2}^n$  (0.4) and  $\beta_{nuc4}^n$  (0.32), obtained by the slopes of  $\log k_i$  vs.  $\log K_i$  plots, indicate that in the transition state leading to  $2H^+$  and  $4H^+$  adducts, about 30–40% of the positive charge is localized on the attacking amino group and hence the forming C–N bond is less than half complete.

A comparison of  $k_4/k_2$  ratio (2.8) with that observed for the less reactive primary amines (3.8)<sup>2</sup> and for the more reactive methoxide ion in methanol (1.4)<sup>12</sup> shows that the decrease of selectivity with the increase of reactivity ( $\approx 100\times$ ) is quite negligible, as observed for nucleophilic reactions of organic cations.<sup>1a</sup>

A comparison of the  $k_{-i}$  terms of the primary and secondary amines (Table I) indicates that in consequence of the greater steric strain of the secondary amine moieties, the latter are better leaving groups than the former. Moreover the nucleofugicity of morpholine is enhanced by the electronic effects responsible of its lower basicity.

The most important feature that distinguishes the secondary amine reaction from that of primary amines is the different rate-controlling step. In the latter reaction the rate-determining

(8) (a) Bunton, C. F.; Huang, S. K. *J. Am. Chem. Soc.* **1974**, *96*, 515. (b) Doddi, G.; Illuminati, G.; Mecozzi, M.; Nunziante, P. *J. Org. Chem.* **1983**, *48*, 5268.

(9) (a) Bernasconi, C. F.; Carrè, D. *J. Am. Chem. Soc.* **1979**, *101*, 2698. (b) Bernasconi, C. F.; Murray, C. J.; Carrè, D. *J. Am. Chem. Soc.* **1983**, *105*, 4349. (c) Buncel, E.; Crampton, M. R.; Strauss, M. J.; Terrier, F. "Electron Deficient Aromatic- and Heteroaromatic-Base Interactions"; Elsevier: Amsterdam, 1984. (d) Buncel, E.; Eggimann, W. *J. Am. Chem. Soc.* **1977**, *99*, 5958. (e) Bernasconi, C. F.; Müller, M. C.; Schmid, P. *J. Org. Chem.* **1979**, *44*, 3189. (f) Crampton, M. R.; Gibson, B. *J. Chem. Soc., Perkin Trans. 2* **1981**, 533.

(10) (a) Aveta, R.; Doddi, G.; Illuminati, G. *J. Am. Chem. Soc.* **1983**, *105*, 5661. (b) Miller, J. "Aromatic Nucleophilic Substitution"; Elsevier: Amsterdam, 1968.

(11) The amine addition to the hindered 1,1-dinitro-2,2-diphenylethylene shows the reverse order primary > secondary: Bernasconi, C. F.; Carrè, D. J.; Fox, J. P. In "Techniques and Applications of Fast Reactions in Solution"; Getting, W. J.; Wyn-Jones, E., Eds.; Reidel: Dordrecht, Holland, 1979; p 453.

(12) Doddi, G.; Illuminati, G.; Insam, N.; Stegel, F. *J. Org. Chem.* **1979**, *44*, 4496.

(7) The corresponding ratio for the reaction of butylamine and cyclohexylamine is independent of the different reactivity of the amines.<sup>2</sup>

step is the nucleophilic attack to yield the  $2H^+$  and  $4H^+$  adducts, whereas in the former the rate-controlling step is the deprotonation of the charged adducts to yield the neutral ones, even though with piperidine the rates of the two steps become comparable at the higher amine concentrations. This different behavior depends both on the increase of  $k_{-i}$  and mainly on a decrease of the  $k_{ip}^{am}$  terms for the secondary amine reaction (Table I). In fact, since for the primary amine reaction  $k_{ip}^{am}[am] > 10k_{-i}$ ,<sup>2</sup> we can estimate that  $k_{ip}^{am} > 10^6 M^{-1} s^{-1}$ , i.e., more than  $10\times$  that observed for the secondary amines, in spite of the comparable  $K_i^{am}$ .

In the following section we account for the factors that can affect such proton transfer processes.


**Rate-Limiting Proton Transfer.** A very significant result is the finding that the strongly favored deprotonation of the charged adducts by amines ( $K_i^{am}$ ) and the protonation of neutral adducts by the solvated proton ( $1/K_{ia}$ ) are activated rather than encounter controlled ( $k_{ip}^{am}$ ,  $k_{-ip}^{sH}$ ).

It is well-known that proton transfer between nitrogen atoms in water is diffusion controlled, provided that the reaction is thermodynamically favored by 2 or more  $pK_a$  units.<sup>13</sup> Moreover, Bianchin and Delpuech reported a rate constant for proton transfer between  $NH_4^+$  and  $NH_3$  in  $Me_2SO$  ( $1.17 \times 10^9 M^{-1} s^{-1}$ ) that coincides with that observed in aqueous solution ( $1.21 \times 10^9 M^{-1} s^{-1}$ ).<sup>14</sup> This result shows that a fast proton transfer can also occur in an aprotic solvent. However, recent reports on thermodynamically favored proton transfer between bases and charged amine adducts of 1,3,5-trinitrobenzene,<sup>9e,f,15</sup> 3,6-bis(dimethyl-amino)xanthylium ion,<sup>5</sup> and activated alkenes,<sup>9a,b</sup> and acid such as 1-*cis*-2-*cis*-6-trimethylpiperidinium ion,<sup>16</sup> in  $Me_2SO$  or 50% aqueous  $Me_2SO$  solutions, indicate that the rates of these processes are below the diffusion limit. This behavior was attributed to steric interactions hindering the approach of the base to the crowded adduct, and actually an increasing of bulkiness of base can depress the rates of these processes without any relationship with its basicity.<sup>9a</sup>

The importance of the steric effects is also shown by the reaction of **2** with morpholine in  $Me_2SO$ , in which the steric hindrance on the ammonium nitrogen should be lower than the  $4H^+$ -thiopyran, having a rate  $3\times$  higher than  $k_{4p}^{mor}$ , in spite of a three order of magnitude lower acidity.<sup>5</sup> Similar conclusions are drawn from the comparison between  $k_{2p}^{am}$  and  $k_{4p}^{am}$  showing that the more acidic  $2H^+$  adduct ( $\approx 100\times$ ) is deprotonated by the amine more slowly than the  $4H^+$  isomer. Moreover our finding  $k_{2p}^{mor} \approx k_{2p}^{pip}$  and  $k_{4p}^{mor} \approx k_{4p}^{pip}$  clearly shows that if the steric situation between the acid and the base is unchanged, the rate of the proton transfer is not affected.

Steric effects seem to be operating also in the protonation of the  $2H$  and  $4H$  adducts by the solvated proton. A Brønsted plot of  $\log k_{-ip}^{sH}$  vs.  $\log K_{ia}$  is linear with  $\beta \approx 0$ , whereas the thermodynamically unfavored reverse reaction has  $\alpha \approx 1$ . Brønsted slope is commonly interpreted as indicating the degree of proton transfer in the transition state.<sup>17</sup> It follows that in the transition state the proton is released to a very small extent to the neutral adducts. This behavior resembles that shown by diffusion-controlled reactions of "normal" acids,<sup>18</sup> whereas the  $k_{-ip}^{sH}$  terms are largely below the diffusion limit.

A plausible explanation involves restructuring of the solvation shell of the proton. Since the structure of this reagent in  $Me_2SO$  is  $[Me_2SO \cdots H \cdots Me_2SO]^+$ ,<sup>19</sup> the proton cannot be transferred along a Grotthus chain, and a molecule of the solvent must be replaced by the sterically crowded adduct in order for the proton transfer to take place. The Eigen scheme for this proton transfer process is



Our findings indicate that the rate-controlling step is the conversion of the encounter complex **b** into the hydrogen-bonded complex **c**.

Kreevoy reported on a slow protonation of substituted tri-benzylamines by the solvated proton in  $Me_2SO$  that shows  $\beta = 0.59$ .<sup>20</sup> Tribenzylamines should be less hindered than the  $2H$ - and  $4H$ -thiopyrans, and the **b**-**c** process is expected to be favored, making the proton transfer process **c**-**d** concurring to the overall rate.

Even though Brønsted coefficients suggesting diffusion-controlled processes are reported for proton transfer reactions of carbon acids whose rates are below the diffusion limit,<sup>21</sup> this is a remarkable finding for a nitrogen base.

### Experimental Section

**Materials.** 2,4,6-Triphenylthiopyrylium perchlorate was available from our previous work.<sup>2</sup>

$Me_2SO$  was dried as previously described.<sup>2</sup> The amines were refluxed over sodium for 2 h and distilled under argon.

Piperidinium and morpholinium perchlorates were prepared by the respective amines and aqueous perchloric acid (70%) and, after removal of the water, recrystallized twice from benzene-acetone 3:1 and 2:1, respectively. (**Warning:** Organic perchlorates are potentially explosive.)

Tetramethylammonium acetate was prepared by neutralization of a 25% aqueous solution of tetramethylammonium hydroxide (FLUKA) with acetic acid and by crystallization from acetone after removal of the water. All the salts were dried by warming at 80 °C under vacuum.

**Spectra.** The UV-vis spectra of the substrate and the corresponding  $2H$  adducts with piperidine and morpholine (Figure 1) were recorded on a Cary 219 spectrophotometer.

**Rate Measurements.** All kinetic experiments were carried out on a Durrum 110 stopped-flow spectrophotometer at 25 °C under pseudo-first-order conditions. The ionic strength was maintained at 0.1 M with  $KClO_4$ . The reaction solutions were freshly prepared and handled under argon. The two relaxation processes were monitored at 410 nm, where only **1** absorbs ( $[1] \approx 10^{-5} M$ ), and at 355 nm, where both **1** and the corresponding  $2H$  adduct absorb.<sup>22</sup>

**Equilibrium Measurements.** The  $pK_a$  of morpholine (8.9) in  $Me_2SO$  at  $\mu = 0.1 M$  was determined spectrophotometrically as previously described.<sup>2</sup>

The equilibrium constant  $K_{2T}$ , referred to the equilibrium **1** + 2amine  $\rightleftharpoons 2H$  + ammonium (see Scheme I), was determined spectrophotometrically as previously described<sup>2</sup> (for the morpholine reaction a 1-cm quartz cell was used).

### Appendix

Assuming at first that during the first process the formation of the  $2H$  adduct is negligible with respect to the  $4H$  isomer,<sup>23</sup> eq 7 reduces to:

$$\tau_1^{-1} \approx \frac{k_4 k_{4p}^{am} [am]^2}{k_{-4} + k_{4p}^{am} [am]} \quad (16)$$

that rearranged gives:

$$\frac{[am]}{\tau_1^{-1}} \approx \frac{k_{-4}}{k_4 k_{4p}^{am} [am]} + \frac{1}{k_4} \quad (17)$$

The plot of  $[am]/\tau_1^{-1}$  vs.  $1/[am]$  is linear with intercept  $\approx 1/k_4$  and slope  $\approx k_{-4}/k_4 k_{4p}^{am}$ , and hence  $k_4$  and  $k_{4p}^{am}/k_{-4}$  can be

(13) Ahrens, M.-L.; Maass, G. *Angew. Chem., Int. Ed. Engl.* **1968**, *7*, 818.

(14) Bianchin, B.; Delpuech, J.-J. *Bull. Soc. Chim. Fr.* **1973**, 34.

(15) Crampton, M. R.; Greenhalgh, C. *J. Chem. Soc., Perkin Trans. 2* **1983**, 1175.

(16) Delpuech, J.-J.; Bianchin, B. *J. Am. Chem. Soc.* **1979**, *101*, 383.

(17) Kresge, A. J. In "Proton-Transfer Reactions"; Caldin, E., Gold, V., Eds.; Chapman and Hall: London, 1975.

(18) Eigen, M. *Angew. Chem., Int. Ed. Engl.* **1964**, *3*, 1.

(19) (a) Kreevoy, M. M.; Williams, J. M. *J. Am. Chem. Soc.* **1967**, *89*, 5499. (b) Potts, R. A. *Inorg. Chem.* **1970**, *9*, 1284.

(20) Kreevoy, M. M.; Wang, Y. *J. Phys. Chem.* **1977**, *81*, 1924.

(21) Hibbert, F.; Long, F. A.; Walters, E. A. *J. Am. Chem. Soc.* **1971**, *93*, 2829.

(22) The molar extinction coefficients of  $4H$ -thiopyrans are generally negligible with respect to  $2H$ -isomers, at wavelengths higher than 300 nm. Suld, G.; Price, C. C. *J. Am. Chem. Soc.* **1962**, *84*, 2090.

(23) This assumption is supported by the absorbance value at 355 nm when the first process is over, that indicates a concentration of the  $2H$  adduct lower than the  $4H$  isomer.

estimated. Rearranging eq 11 affords:

$$\frac{\tau_2^{-1}[k_4k_{4p}^{\text{am}}[\text{am}]^2/(k_{-4} + k_{4p}^{\text{am}}[\text{am}])]}{[k_{-4}k_{-4p}^{\text{amH}}[\text{amH}]/(k_{-4} + k_{4p}^{\text{am}}[\text{am}]) - \tau_2^{-1}} = \frac{k_2k_{2p}^{\text{am}}[\text{am}]^2}{k_{-2} + k_{2p}^{\text{am}}[\text{am}]} \quad (18)$$

The left-hand term of eq 18 that for the sake of simplicity we indicate as  $k_2^*$ , can be evaluated by introducing the approximate value of  $k_4$  and  $k_{4p}^{\text{am}}/k_{-4}$  just obtained.

Applying the same mathematical treatment of eq 16 to eq 18 affords:

$$\frac{[\text{am}]}{k_2^*} = \frac{k_{-2}}{k_2k_{2p}^{\text{am}}[\text{am}]} + \frac{1}{k_2} \quad (19)$$

From the plot of  $[\text{am}]/k_2^* \text{ vs. } 1/[\text{am}]$ , approximate values of  $k_2$  and  $k_{2p}^{\text{am}}/k_{-2}$  can be obtained.

With the introduction of these terms into eq 7, better values of  $[k_4k_{4p}^{\text{am}}[\text{am}]^2/(k_{-4} + k_{4p}^{\text{am}}[\text{am}])]$  are obtained, and so on, until constant values of  $k_2$ ,  $k_4$ ,  $k_{2p}^{\text{am}}/k_{-2}$ , and  $k_{4p}^{\text{am}}/k_{-4}$  are obtained.

Actually this iterative procedure was carried out with a personal computer by using a non-linear least-squares method.<sup>24</sup>

**Registry No.** 1-CO<sub>4</sub><sup>-</sup>, 2930-37-2; 2-HClO<sub>4</sub>, 92314-76-6; 2H (X = CH<sub>2</sub>), 92314-68-6; 2H (X = O), 87691-76-7; 2H-HClO<sub>4</sub> (X = CH<sub>2</sub>), 92314-69-7; 2H-HClO<sub>4</sub> (X = O), 92314-70-0; 4H (X = CH<sub>2</sub>), 92314-71-1; 4H (X = O), 92314-73-3; 4H-HClO<sub>4</sub> (X = CH<sub>2</sub>), 92314-72-2; 4H-HClO<sub>4</sub> (X = O), 92314-74-4; Me<sub>4</sub>N<sup>+</sup>OH<sup>-</sup>, 75-59-2; CH<sub>3</sub>C(O)O<sup>-</sup>Me<sub>4</sub>N<sup>+</sup>, 1058-12-1; piperidine, 110-89-4; piperidinium perchlorate, 57367-18-7; morpholine, 110-91-8; morpholinium perchlorate, 35175-75-8.

**Supplementary Material Available:** Tables S1-S5 summarizing all the kinetic measurements (6 pages). Ordering information is given on any current masthead page.

(24) Wentworth, W. E. *J. Chem. Educ.* **1965**, *42*, 96, 162.

(25) Courtot-Coupez, J.; Le Demez, M. *Bull. Soc. Chim. Fr.* **1969**, 1033.

## Kinetics of the 1,2-Migration of Carbon-Centered Groups in 2-Substituted 2,2-Dimethylethyl Radicals<sup>1</sup>

D. A. Lindsay, J. Lusztyk,<sup>2</sup> and K. U. Ingold\*


*Contribution from the Division of Chemistry, National Research Council of Canada,*

*Ottawa, Ontario, Canada K1A 0R6. Received March 22, 1984.*

*Revised Manuscript Received June 18, 1984*

**Abstract:** The rearrangements  $\text{RCMe}_2\text{CH}_2\cdot \rightarrow \text{RCH}_2\dot{\text{C}}\text{Me}_2$  ( $k_r$ ) (R = Ph, Me<sub>3</sub>CC≡C, Me<sub>3</sub>CC=O, and N≡C) have been studied over a range of temperatures by product analyses with use of the common competing reaction  $\text{RCMe}_2\text{CH}_2\cdot + \text{CCl}_4 \rightarrow \text{RCMe}_2\text{CH}_2\text{Cl} + \text{CCl}_3\cdot$  ( $k_{\text{Cl}}$ ). For R = H<sub>2</sub>C=CH the rearrangement was so fast that only the rearranged chloride, RCH<sub>2</sub>CMe<sub>2</sub>Cl, was produced. All these rearrangements occur via a 3-membered cyclic intermediate radical (or transition state). Various considerations led to the following Arrhenius equation for chlorine abstraction:  $\log(k_{\text{Cl}}/\text{M}^{-1}\text{s}^{-1}) = (8.14 \pm 0.42) - (5.52 \pm 0.63)/\theta$ , where  $\theta = 2.3RT$  kcal/mol, and this equation is used to calculate Arrhenius parameters for migration of all but the H<sub>2</sub>C=CH group. Comparison of these parameters with those already available from kinetic EPR measurements leads to a choice of preferred Arrhenius parameters for all five rearrangements. The cyano group had an unexpectedly low mobility while the pivaloyl group underwent a surprisingly rapid 1,2-shift. Migratory aptitudes increase along the series R = N≡C < Me<sub>3</sub>CC≡C < Ph < Me<sub>3</sub>CC=O < H<sub>2</sub>C=CH, with  $k_r$  at 25 °C = 0.9, 93, 762,  $1.7 \times 10^5$ , and  $1.0 \times 10^7$  s<sup>-1</sup>, respectively. The preferred pre-exponential factors all lie in the range  $10^{10.9}$ – $10^{12.0}$  s<sup>-1</sup>, while the activation energies vary from 16.4 kcal/mol for R = N≡C to 5.7 kcal/mol for R = H<sub>2</sub>C=CH. These results are discussed in relation to the results of thermochemical kinetic calculations and to earlier work on the 1,2-migration of unsaturated groups in radicals.

In free radical chemistry the 1,2-migration of an unsaturated group has excited interest<sup>3</sup> ever since the discovery by Urry and Kharasch<sup>4</sup> in 1944 of the neophyl rearrangement, i.e., **1a** → **2a**.



**a**, R = C<sub>6</sub>H<sub>5</sub>; **b**, R = Me<sub>3</sub>CC≡C;

**c**, R = CH<sub>2</sub>=CH; **d**, R = Me<sub>3</sub>CC=O; **e**, R = C≡N

In the early 1970's kinetic EPR spectroscopic methods were developed for measuring the rates of unimolecular radical reac-

tions.<sup>5-9</sup> This advance was important because once the Arrhenius parameters for a radical rearrangement have been determined the rearrangement can serve as a "clock" to measure rate constants for appropriate radical-molecule reactions.<sup>10</sup> We have used the EPR technique to measure Arrhenius parameters for the neophyl rearrangement<sup>11</sup> and for some related reactions;<sup>11</sup> for a homo-propargylic rearrangement,<sup>12</sup> **1b** → **2b**; and for a homoallylic rearrangement,<sup>13-16</sup> **1c** → **2c**. To the best of our knowledge there

(5) Watts, G. B.; Griller, D.; Ingold, K. U. *J. Am. Chem. Soc.* **1972**, *94*, 8784-8789.

(6) Griller, D.; Roberts, B. P. *J. Chem. Soc., Perkin Trans. 2* **1972**, 747-751.

(7) Davies, A. G.; Griller, D.; Roberts, B. P. *J. Chem. Soc., Perkin Trans. 2* **1972**, 993-998.

(8) Lal, D.; Griller, D.; Husband, S.; Ingold, K. U. *J. Am. Chem. Soc.* **1974**, *96*, 6355-6357.

(9) Griller, D.; Ingold, K. U. *Acc. Chem. Res.* **1980**, *13*, 193-200.

(10) Griller, D.; Ingold, K. U. *Acc. Chem. Res.* **1980**, *13*, 317-323.

(11) Maillard, B.; Ingold, K. U. *J. Am. Chem. Soc.* **1976**, *98*, 1224-1226, 4692.

(12) Ingold, K. U.; Warkentin, J. *Can. J. Chem.* **1980**, *58*, 348-352.

(1) Issued as NRCC No. 23760.

(2) N.R.C.C. Research Associate 1982-1984.

(3) For recent reviews of radical rearrangements see: (a) Beckwith, A. L. J.; Ingold, K. U. In "Rearrangements in Ground and Excited States"; de Mayo, P., Ed.; Academic Press: New York, 1980; pp 161-310. (b) Beckwith, A. L. J. *Tetrahedron* **1981**, *37*, 3073-3100. (c) Surzur, J. M. In "Reactive Intermediates"; Abramovitch, R. A., Ed.; Plenum Press: New York, 1982; pp 121-295.

(4) Urry, W. H.; Kharasch, M. S. *J. Am. Chem. Soc.* **1944**, *66*, 1438-1440.