

Thirdly, we predict a variation in $r_0(\text{CH})$ with internal rotation angle of the order of 0.004–0.006 Å, in both compounds. While the absence of a fully quantitative treatment of the potential function precludes an exact prediction, the general resemblance of the $\nu_{\text{as}}(\text{CH}_3)$ and $\nu(\text{CHD}_2)$ splittings to those in $(\text{MeBO})_3$, where a variation of the above order is confirmed by ab initio calculations,³⁸ provides additional support for the proposed bond length fluctuation in the $\text{MeMn}(\text{CO})_5$ and $\text{MeRe}(\text{CO})_5$ compounds.

Conclusions

1. The infrared spectra of $\text{MeMn}(\text{CO})_5$ and $\text{MeRe}(\text{CO})_5$ species in the CH and CD stretching regions can be qualitatively interpreted in terms of a CH stretching force constant which varies with internal rotation angle of an essentially freely rotating methyl group.

(38) Although described earlier as 4-31G calculations,¹² these were in fact STO-3G ones, scaled up according to STO-3G/4-31G differences in MeNO_2 and MeBF_2 (P. D. Mallinson, private communication).

2. The ground-state barrier to internal rotation is probably of V_{12} origin and very low, $<10 \text{ cm}^{-1}$.

3. Mean CH bond lengths of 1.095₉ and 1.098₀ are predicted for the Mn and Re compounds, respectively, with variations of the order of 0.004–0.006 Å during internal rotation.

4. The CH bonds are of average strength, that in the Re compound being slightly the weaker, and bear an inverse relationship to $D_0(\text{M}-\text{CH}_3)$, as in other MeM compounds.

5. Further development of theory is needed for a quantitative interpretation of the spectra, especially the $\nu_{\text{as}}(\text{CH}_3)$ and $\nu_{\text{as}}(\text{CD}_3)$ splittings.

Acknowledgment. We are much indebted to Dr. D. Cavagnat for detailed calculations of CH_3 and CD_3 internal rotation/torsion energy levels. We thank the S.E.R.C. for supporting this work.

Registry No. $\text{Zn}(\text{CHD}_2)_2$, 92525-33-2; $\text{CH}_3\text{Re}(\text{CO})_5$, 14524-92-6; $\text{CH}_3\text{Mn}(\text{CO})_5$, 13601-24-6; $\text{CHD}_2\text{Re}(\text{CO})_5$, 92525-30-9; $\text{CHD}_2\text{Mn}(\text{CO})_5$, 92525-31-0; $\text{CH}_2\text{DMn}(\text{CO})_5$, 20792-56-7; $\text{CD}_3\text{Mn}(\text{CO})_5$, 15653-52-8; $\text{CD}_3\text{Re}(\text{CO})_5$, 92525-32-1.

Molecular Structure of *s-cis*- and *s-trans*-Acrolein Determined by Microwave Spectroscopy

C. E. Blom,[†] G. Grassi, and A. Bauder*

Contribution from the Laboratorium für Physikalische Chemie, Eidgenössische Technische Hochschule, CH-8092 Zürich, Switzerland. Received December 21, 1983

Abstract: The rotational spectra of highly enriched single D-, ¹³C-, and ¹⁸O-substituted species of acrolein have been measured and analyzed over 12–58 GHz. The complete substitution structure has been determined for the less abundant *s-cis* conformer from the ground-state rotational constants. In addition newly assigned μ_b -type transitions for all isotopic species of the more abundant *s-trans*-acrolein have improved the structure of this conformer. Careful measurements of the Stark effect have resulted in an accurate determination of the electric dipole moment of the *s-trans* conformer. A comparison of the molecular structures of the two conformers has revealed significant differences in the central C–C bonds.

Among the simplest 1,3-conjugated double-bond systems butadiene ($\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$), acrolein ($\text{CH}_2=\text{CH}-\text{CH}=\text{O}$), and glyoxal ($\text{O}=\text{CH}-\text{CH}=\text{O}$) only acrolein possesses a non-vanishing electric dipole moment in the most abundant *s-trans* conformer.¹ Acrolein represents a crucial system for studying the effect of the change of conformation on the structure in a 1,3-conjugated double-bond system by the microwave method. Several investigations of the rotational spectra of *s-trans*-acrolein have been reported.^{2–5} They include the assignment of μ_a -type transitions of isotopically substituted species measured in natural abundance except for the deuterated species.³ Their analysis allowed the determination of a substitution structure of the planar *s-trans* conformer. Only recently the rotational spectrum of the less abundant *s-cis*-acrolein has been assigned.⁶ Its analysis established that this conformer is likewise a planar molecule.

The present contribution continues our earlier work on *s-cis*-acrolein,⁶ providing now the complete substitution structure of this conformer. Highly enriched isotopically substituted species of acrolein had to be prepared since the intensities of the rotational transitions of the *s-cis* conformer were only 2% of those of the *s-trans* conformer. In the course of this study it became clear that more accurate data for the isotopic species of *s-trans*-acrolein were needed in order to make a comparison of the structural parameters of *s-cis* and *s-trans* conformers possible. Therefore we have re-measured the rotational spectra of D-, ¹³C-, and ¹⁸O-substituted

species of *s-trans*-acrolein and have also assigned μ_b -type transitions. With the help of the newly measured transition frequencies three rotational and five quartic centrifugal distortion constants could be determined for all isotopic species, improving the previous analysis.³ In addition accurate measurements of the Stark splittings of optimally selected transitions have allowed a better determination of the electric dipole moment, especially of the μ_b component.

Experimental Details

Chemicals. Five isotopic species of acrolein were prepared by an aldol condensation from appropriate isotopic species of formaldehyde and acetaldehyde.⁷ The reactions were carried out in the gas phase at 280 °C in a reactor filled with an alkaline catalyst following the modifications by Renaud and Stephens.⁸ The catalyst was composed of silica gel 40 (70–230 mesh, Fluka AG) treated with a 10% sodium carbonate solution.⁷ The crude products from the aldol condensation were purified by gas chromatography (Perkin-Elmer F-21, 4.5-m packed column with 15% QF-1 on Chromosorb W, 40 °C).

(1) *s-trans* refers to the trans orientation of the conjugated double bonds about the C–C single bond, *s-cis* to the cis orientation.

(2) Wagner, R.; Fine, J.; Simmons, J. W.; Goldstein, J. H. *J. Chem. Phys.* **1957**, *26*, 634.

(3) Cherniak, E. A.; Costain, C. C. *J. Chem. Phys.* **1966**, *45*, 104.

(4) Benson, R. C.; Flygare, W. H. *J. Chem. Phys.* **1973**, *58*, 2366.

(5) Winniewisser, M.; Winniewisser, G.; Honda, T.; Hirota, E. *Z. Naturforsch. A* **1975**, *30A*, 1001.

(6) Blom, C. E.; Bauder, A. *Chem. Phys. Lett.* **1982**, *88*, 55.

(7) Schulz, H.; Wagner, H. *Angew. Chem.* **1950**, *62*, 105.

(8) Renaud, R. N.; Stephens, J. C. *J. Labelled Compd.* **1972**, *8*, 509.

[†] Present address: Physikalisch-Chemisches Institut der Justus-Liebig-Universität Giessen, Heinrich-Buff-Ring 58, D-6300 Giessen, F.R.G.

Specifically, the following isotopic species were synthesized with the above method:

Acrolein-1-*d* was obtained from acetaldehyde-1-*d* (98% D, Stohler Isotope Chemicals) and formaldehyde.

Acrolein-2-*d* was obtained from acetaldehyde and formaldehyde in D₂O solution.

Acrolein-1-¹³C was obtained from acetaldehyde-1-¹³C (90% ¹³C) and formaldehyde.

Acrolein-2-¹³C was obtained from acetaldehyde-2-¹³C (90% ¹³C) and formaldehyde.

Acrolein-3-¹³C was obtained from equimolar amounts of acetaldehyde and formaldehyde-¹³C (99% ¹³C, 20% solution in water, Stohler Isotope Chemicals).

Acetaldehyde-1-¹³C and -2-¹³C were prepared by first reducing acetic acid-1-¹³C or -2-¹³C (both 90% ¹³C, Stohler Isotope Chemicals) with LiAlH₄ to ethanol-1-¹³C and -2-¹³C, respectively.⁹ The ethanols were then oxidized with sodium dichromate in diluted sulfuric acid to the corresponding aldehydes.¹⁰

Acrolein-*cis*-3-*d* and acrolein-*trans*-3-*d* were prepared by the oxidation of allylic alcohol-*cis*-3-*d* or -*trans*-3-*d*, respectively, with molecular oxygen on a silver catalyst at 250–280 °C by the method of Imachi et al.¹¹ The crude products from the reaction were purified by gas chromatography (Perkin-Elmer F-21, 4.5-m packed column with 15% SE-30 on Chromosorb W, 40 °C). The isotopic purity was determined to be 60% for the *cis* isomer and 40% for the *trans* isomer.

Allylic alcohol-*cis*-3-*d* was obtained from the reduction of propargyl alcohol (2-propinol) with LiAlH₄ and subsequent hydrolysis with D₂O.¹²

Allylic alcohol-*trans*-3-*d* on the other hand was prepared by reducing 2-propinol-3-*d* with LiAlH₄ and hydrolysis with H₂O. 2-Propinol was deuterated with 5 N NaOD in D₂O for 1 h at 100 °C, giving 2-propinol-3-*d* with 80% yield. The degree of deuteration reached 80%.

Acrolein-¹⁸O was synthesized by acidic hydrolysis of acrolein dipropyl acetal with H₂¹⁸O (98.4% ¹⁸O, Ventron AG).¹³ The reaction was carried out in a microdistillation apparatus with a 5-cm-long Vigreux column. The temperature of the reaction mixture was slowly increased to 98 °C, and acrolein-¹⁸O was distilled off and collected with a yield of 75%.¹³ The raw product was dried over molecular sieve A3 and contained 90% ¹⁸O.

All samples of isotopic species of acrolein were stabilized by addition of a few crystals of hydroquinone against polymerization. They could be stored for several months at -80 °C without apparent decomposition.

Spectrometers. The microwave spectra were recorded over 12–58 GHz with conventional Stark spectrometers. Phase-stabilized backward-wave oscillators were used as microwave sources below 39 GHz. For the higher frequencies klystrons were phase-locked against the third or fourth harmonic of a stable 12–15 GHz backward-wave oscillator. The 4-m-long Stark cell was driven by a 30-kHz square wave. All measurements were made with less than 10 mtorr of pressure and at 22 ± 2 °C except for the determination of the dipole moment of *s-trans*-acrolein. Here the Stark cell was cooled to -70 °C by circulating methanol from a cryostat. The cooling reduced the interference from overlapping transitions in excited states.

For accurate frequency measurements the microwave spectrometers were operated under control of a PDP-8/E computer. Several sweeps over the same range could be summed in order to improve the signal-to-noise ratio. A parabola was fitted to the center of the absorption line and its maximum determined. The final frequencies of rotational transitions were averaged from sweeps in both directions and should be accurate to within 50 kHz.

Analysis of Rotational Spectra

Isotopic Species of *s-cis*-Acrolein. Predictions of transition frequencies for isotopically substituted species of *s-cis*-acrolein were based on the known moments of inertia of the parent species⁶ and on the isotopic shifts calculated from a model. The assignment always started with the 3(2,2)–2(2,1) and 3(2,1)–2(2,0) μ_a -type transitions which could be observed with very low Stark fields (30 V cm⁻¹). The search with low Stark fields was indispensable for the *s-cis* conformer which is present only with 4% abundance. At higher Stark fields many much stronger lines appeared which arise from various excited states of the abundant *s-trans* conformer.




Figure 1. Atomic numbering scheme, dipole moment, and molecular structure from least-squares fit (bond lengths in Å, bond angles in deg) for *s-cis*-acrolein and *s-trans*-acrolein.

This was even more pronounced for isotopic species when additional transitions from isotopic impurities were observed simultaneously. Preliminary values of the rotational constants *B* and *C* were adjusted from the two above-mentioned transition frequencies. Subsequently better predictions for *J* = 3–2 and *J* = 2–1 μ_a -type transitions were possible. Further transitions were then readily assigned even at the higher Stark fields necessary for their modulation. All three rotational constants could now be adjusted.

Low *J* members of the μ_b -type Q-branch series *J*(1, *J*–1)–*J*(0, *J*) were found within a few MHz of predicted positions based on experimental rotational constants and centrifugal distortion constants transferred from the parent species. Assignments of μ_a - and μ_b -type transitions were extended step by step to higher *J* values, finally including 33–40 transitions with *J* ≤ 11 for each isotopic species. For acrolein-*trans*-3-*d* (CDH=CH–CHO) only 14 transitions with *J* ≤ 6 were measured. In this case Δ_K and δ_K were fixed at the values of the parent species. Tables with listings of all measured transition frequencies are available as supplementary material.

Rotational constants and quartic centrifugal distortion constants in the vibrational ground state were adjusted for each of the eight singly substituted isotopic species of *s-cis*-acrolein from the measured transition frequencies. The results are collected in Table I. The centrifugal distortion constants are defined according to Watson's asymmetric reduction in a prolate I' representation.¹⁴

Isotopic Species of *s-trans*-Acrolein. In the previous investigation³ of the isotopic species of *s-trans*-acrolein only μ_a -type transitions were assigned. The rotational constants *A* were badly determined and only *B* and *C* could be adjusted. The lowest member of the μ_b -type Q-branch series, 1(1,0)–1(0,1), was predicted for each species from the published *B* and *C* and assuming an inertia defect of -0.0212 uÅ² equal to that of the parent species. These transitions could then easily be assigned, although with deviations from the predicted frequencies reaching 36 MHz. After preliminary adjustments of the rotational constants and the centrifugal distortion constants transitions with successively higher

(9) Ostwald, R.; Adams, P. T.; Tolbert, B. M. *J. Am. Chem. Soc.* **1952**, *74*, 2425.

(10) Gattermann, L. "Die Praxis des organischen Chemikers", 36th ed.; De Gruyter: Berlin, 1954.

(11) Imachi, M.; Cant, N. W.; Kuczkowski, R. L. *J. Catal.* **1982**, *75*, 404.

(12) Bates, E. B.; Jones, E. R. H.; Whiting, M. C. *J. Chem. Soc.* **1954**, 1854.

(13) Sawyer, C. B. *J. Org. Chem.* **1972**, *37*, 4225.

(14) Watson, J. K. G. In "Vibrational Spectra and Structure"; Durig, J. R., Ed.; Elsevier: New York, 1977; Vol. 6.

Table I. Rotational Constants (MHz) and Quartic Centrifugal Distortion Constants (kHz) for Isotopically Substituted Species of *s-cis*-Acrolein in the Vibrational Ground State^a

	¹³ CH ₂ CH- CHO	CH ₂ ¹³ CHCHO	CH ₂ CH ¹³ CHO	CH ₂ CHCH ¹⁸ O
A	22574.762 (18)	22485.459 (22)	22565.122 (21)	22339.531 (14)
B	6073.707 (3)	6198.712 (4)	6200.653 (4)	5973.952 (3)
C	4786.881 (2)	4860.028 (3)	4864.975 (3)	4714.284 (2)
Δ _J	4.93 (4)	4.97 (5)	4.95 (8)	4.78 (5)
Δ _{JK}	-28.8 (4)	-27.3 (4)	-28.5 (3)	-27.8 (3)
Δ _K	105 (5)	106 (5)	110 (4)	106 (3)
δ _J	1.419 (11)	1.495 (15)	1.454 (11)	1.369 (11)
δ _K	10.4 (5)	9.2 (8)	11.4 (5)	10.2 (6)
Δ ^b	-0.0187	-0.0186	-0.0195	-0.0181
	CH ₂ CD- CHO	CH ₂ CHCDO	<i>cis</i> - CHDCHCHO ^c	<i>trans</i> - CDHCHCHO ^d
A	20594.106 (8)	20811.966 (16)	20796.961 (20)	22634.762 (55)
B	6128.360 (2)	6118.233 (3)	6163.027 (3)	5770.070 (10)
C	4723.899 (2)	4729.863 (3)	4755.207 (3)	4598.859 (11)
Δ _J	4.62 (5)	4.62 (5)	5.28 (7)	4.46 (14)
Δ _{JK}	-21.5 (2)	-22.5 (3)	-24.5 (3)	-30.9 (17)
Δ _K	79 (2)	80 (3)	76 (4)	(109.2) ^e
δ _J	1.407 (8)	1.389 (10)	1.554 (12)	1.549 (178)
δ _K	9.6 (4)	9.3 (4)	12.7 (6)	(11.36) ^e
Δ ^b	-0.0222	-0.0367	-0.0233	-0.0216

^aNumbers in parentheses represent one standard deviation. ^bInertia defect $\Delta = I_C - I_A - I_B$ (uÅ²), conversion factor 505379.05 MHz uÅ². ^cD is located in the *cis* position with respect to the C-C single bond. ^dD is located in the *trans* position with respect to the C-C single bond. ^eFixed at the value of the parent CH₂CHCHO.

Table II. Rotational Constants (MHz) and Quartic Centrifugal Distortion Constants (kHz) for Isotopically Substituted Species of *s-trans*-Acrolein in the Vibrational Ground State^a

	¹³ CH ₂ CH- CHO	CH ₂ ¹³ CH- CHO	CH ₂ CH- ¹³ CHO	CH ₂ CH- CH ¹⁸ O
A	47255.232 (6)	46518.949 (6)	46781.044 (7)	47262.882 (6)
B	4520.7955 (6)	4642.4397 (7)	4644.7419 (7)	4428.1025 (6)
C	4126.6426 (6)	4221.7446 (6)	4225.8363 (6)	4049.3455 (6)
Δ _J	0.9922 (25)	1.0308 (21)	1.0401 (22)	0.9313 (19)
Δ _{JK}	-8.81 (8)	-8.75 (6)	-8.66 (7)	-8.28 (6)
Δ _K	362.4 (10)	368.4 (7)	349.4 (8)	357.8 (8)
δ _J	0.11169 (14)	0.12134 (12)	0.12073 (13)	0.10249 (11)
δ _K	5.423 (12)	5.740 (12)	5.616 (14)	5.295 (15)
Δ ^b	-0.0172	-0.0160	-0.0171	-0.0178
	CH ₂ CDCHO	CH ₂ CHCDO	<i>cis</i> - CHDCH- CHO ^c	<i>trans</i> - CDHCH- CHO ^d
A	39038.070 (5)	39514.424 (5)	41654.613 (5)	46660.088 (8)
B	4647.9057 (7)	4651.7010 (6)	4508.5717 (6)	4356.7777 (8)
C	4153.6536 (6)	4162.3476 (6)	4068.6004 (5)	3985.4051 (8)
Δ _J	1.0071 (22)	1.0302 (22)	1.0444 (17)	0.8582 (32)
Δ _{JK}	-3.83 (5)	-5.09 (6)	-6.37 (5)	-6.71 (10)
Δ _K	239.7 (4)	202.9 (5)	266.8 (5)	309.8 (14)
δ _J	0.13658 (16)	0.13418 (14)	0.13277 (11)	0.09183 (18)
δ _K	6.177 (16)	4.992 (8)	6.370 (12)	4.705 (19)
Δ ^b	-0.0075	-0.0168	-0.0111	-0.0220

^{a-d}See corresponding footnotes in Table I.

J were assigned and measured. Finally 63–95 transitions with $J \leq 40$ were collected for each isotopic species of *s-trans*-acrolein. The listings of the measured transition frequencies are available as supplementary material. The rotational constants and the quartic centrifugal distortion constants fitted to the measured transition frequencies are shown in Table II.

Electric Dipole Moment of *s-trans*-Acrolein. The Stark splittings were accurately observed for three M components of the transition 4(1,4)–3(1,3) and for all M components of the transition 4(0,4)–3(0,3). These transitions were selected because

Table III. Experimental and Calculated Stark Coefficients (Hz V⁻² cm²) and Electric Dipole Moment (D) of *s-trans*-Acrolein

transition	M	Δν/E ²	
		obsd ^a	obsd – calcd
4(1,4)–3(1,3)	3	49.241 (143) ^b	-0.255
	2	18.595 (51)	0.026
	0	-6.171 (8)	0.000
4(0,4)–3(0,3)	3	5.821 (30)	0.044
	2	0.635 (20)	0.005
	1	-2.532 (50)	-0.074
	0	-3.600 (60)	-0.113
		this work	ref 1
	μ _a	3.052 (4) ^b	3.06 (4)
	μ _b	0.630 (1)	0.54 (14)
	μ _c	0.0 ^c	0.0
	μ _{tot}	3.117 (4)	3.11

^aA dipole moment of 0.71521 D for OCS (ref 15) was used to calibrate the Stark cell. ^bThe uncertainty is given in parentheses as one standard deviation. ^cAssumed.

Table IV. Substitution Coordinates (Å) of *s-cis*-Acrolein in the Principal Axes System of the Parent Species CH₂=CH–CHO from the Equations of Kraitchman and from the Least-Squares Fit

nucleus	Kraitchman		least squares	
	a	b	a	b
C ₁	0.7299	0.5173	0.7287	0.5202
C ₂	-0.7472	0.5911	-0.7473	0.5909
C ₃	-1.5014	-0.5154	-1.5013	-0.5165
O ₄	1.3623	-0.5176	1.3624	-0.5162
H ₅	1.2554	1.4910	1.2564	1.4924
H ₆	-1.2014	1.5789	-1.2014	1.5793
H ₇	-0.9985	-1.4922	-0.9988	-1.4925
H ₈	-2.5813	-0.4643	-2.5812	-0.4659

they exhibited a substantial dependence on the μ_b dipole component. For each M component the slope was calculated by linear regression from the Stark shifts at ten different Stark fields. The electric field in the Stark cell was calibrated with the help of the known dipole moment of OCS of 0.71521 D.¹⁵ All measurements including the electric field calibration were made at -70 °C in order to suppress overlapping transitions from excited states. The non-vanishing components of the permanent electric dipole moment were determined in a least-squares fit of the measured Stark slopes by using weighting factors equal to the inverse squared standard deviations of the slopes. The Stark coefficients were obtained from second-order perturbation. The measured and calculated slopes and the dipole components are given in Table III.

Substitution Structures

The small values of the inertia defects Δ and their changes upon isotopic substitutions allowed the conclusion that both conformers of acrolein are planar. Changes occurred in both directions from the values of the inertia defects of the parent molecules. The magnitudes of the changes are typical for residual vibrational effects and do not indicate a stable nonplanar equilibrium structure. The analysis of excited vibrational states gave no hint of a potential barrier for the planar structure.¹⁶ In addition the accurate Stark effect measurements excluded the presence of a substantial μ_c dipole component in both conformers.

***s-cis*-Acrolein.** The Cartesian coordinates of all nuclei in *s-cis*-acrolein were determined from the differences between the moments of inertia I_a, I_b, and I_c of the singly substituted species and those of the parent species. The Cartesian coordinates refer to the principal axis system of the parent species. The three-dimensional equations of Kraitchman¹⁷ were applied to all three differences ΔI_a, ΔI_b, and ΔI_c although the molecules are planar.

(15) Muentzer, J. S. J. Chem. Phys. 1968, 48, 4544.

(16) Blom, C. E., unpublished results.

(17) (a) Kraitchman, J. Am. J. Phys. 1953, 21, 17. (b) Costain, C. C. J. Chem. Phys. 1958, 29, 864.

Table V. Structural Parameters of *s-cis*- and *s-trans*-Acrolein

<i>s-cis</i> -Acrolein			
	Kraitchman	least squares	
Bond lengths, Å			
C ₁ -C ₂	1.479 (0) ^a (2) ^b	1.478 (6) ^c	
C ₂ -C ₃	1.339 (1) (2)	1.340 (7)	
C ₁ -O ₄	1.213 (0) (2)	1.215 (6)	
C ₁ -H ₅	1.106 (3) (3)	1.106 (4)	
C ₂ -H ₆	1.087 (1) (3)	1.088 (4)	
C ₃ -H ₇	1.099 (1) (3)	1.098 (5)	
C ₃ -H ₈	1.081 (0) (1)	1.081 (3)	
Bond Angles, deg			
C ₁ C ₂ C ₃	121.4 (0) (1)	121.5 (4)	
O ₄ C ₁ C ₂	124.3 (0) (1)	124.2 (4)	
H ₅ C ₁ C ₂	115.5 (2) (2)	115.8 (4)	
C ₃ C ₂ H ₆	121.0 (0) (2)	121.1 (4)	
C ₂ C ₃ H ₇	118.5 (1) (2)	118.5 (3)	
C ₂ C ₃ H ₈	121.6 (1) (4)	121.5 (6)	
O ₄ C ₁ H ₅	120.2 (2) (2)	120.1 (4)	
C ₁ C ₂ H ₆	117.6 (0) (2)	117.4 (4)	
H ₇ C ₃ H ₈	120.0 (1) (4)	120.0 (5)	
<i>s-trans</i> -Acrolein			
	Kraitchman		least squares
	ref 2	this work	
Bond Lengths, Å			
C ₁ -C ₂	1.470 (3) ^d	1.468 (3) ^a (2) ^b	1.468 (4) ^c
C ₂ -C ₃	1.345 (3)	1.341 (4) (5)	1.340 (4)
C ₁ -O ₄	1.219 (5)	1.215 (2) (3)	1.214 (4)
C ₁ -H ₅	1.108 (3)	1.113 (3) (3)	1.113 (6)
C ₂ -H ₆	1.084 (3)	1.084 (4) (3)	1.084 (5)
C ₃ -H ₇	1.086 (5)	1.089 (7) (10)	1.090 (4)
C ₃ -H ₈	1.086 (5)	1.081 (3) (7)	1.080 (3)
Bond Angles, deg			
C ₁ C ₂ C ₃	119.8 (2)	120.3 (3) (4)	120.4 (5)
O ₄ C ₁ C ₂	123.3 (3)	123.9 (3) (3)	124.0 (6)
H ₅ C ₁ C ₂	115.1 (2)	114.7 (2) (4)	114.7 (5)
C ₃ C ₂ H ₆	122.8 (2)	122.4 (5) (5)	122.4 (4)
C ₂ C ₃ H ₇	120.0 (2)	119.8 (3) (4)	119.7 (3)
C ₂ C ₃ H ₈	121.4 (2)	122.2 (6) (9)	122.2 (5)
O ₄ C ₁ H ₅		121.3 (3) (5)	121.3 (6)
C ₁ C ₂ H ₆		117.3 (4) (3)	117.2 (5)
H ₇ C ₃ H ₈		118.0 (3) (5)	118.1 (6)

^aσ, uncertainty from eq 3 and 4, see text; (0) signifies an uncertainty of less than 0.5 in the last digit. ^bσ', uncertainty from eq 4 and 5, see text. ^cThree times the standard deviation from least-squares refinement. ^dUncertainties quoted in ref 2.

The resulting *c* coordinates were disregarded and assumed to be zero in accordance with strict planarity. The results for the *a* and *b* coordinates listed in Table IV were numerically equal to the values derived from formulas given by Rudolph¹⁸ which use all three moments of inertia in a balanced way for planar molecules. Bond lengths and bond angles derived from the Cartesian coordinates are shown in Table V.

As an alternative, bond lengths and bond angles were adjusted directly to the differences Δ*I*_a, Δ*I*_b, and Δ*I*_c for the isotopically substituted species in an iterative least-squares fit with the program GEOM.¹⁹ Center-of-mass and product-of-inertia conditions are automatically fulfilled for this procedure. The results are compared in Table V to the molecular structure from the substitution coordinates according to the equations of Kraitchman. Conversely, calculated coordinates from the least-squares structure are included in Table IV.

***s-trans*-Acrolein.** The substitution structure as well as the least-squares structure of *s-trans*-acrolein were derived in exactly the same manner as described above for the *s-cis* conformer. The results for the Cartesian coordinates are given in Table VI, and the corresponding bond lengths and bond angles are included in

Table VI. Substitution Coordinates (Å) of *s-trans*-Acrolein in the Principal Axes System of the Parent Species CH₂=CH-CHO from the Equations of Kraitchman and from the Least-Squares Fit

nucleus	Kraitchman					
	ref 2		this work		least squares	
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>
C ₁	0.590	-0.369	0.5908	-0.3646	0.5910	-0.3646
C ₂	-0.635	0.445	-0.6356	0.4422	-0.6353	0.4422
C ₃	-1.837	-0.152	-1.8372	-0.1523	-1.8372	-0.1508
O ₄	1.711	0.111	1.7112	0.1048	1.7113	0.1034
H ₅	0.427	-1.465	0.4243	-1.4649	0.4243	-1.4649
H ₆	-0.517	1.523	-0.5215	1.5205	-0.5200	1.5197
H ₇	-1.902	-1.242	-1.9029	-1.2392	-1.9026	-1.2386
H ₈	-2.758	0.418	-2.7584	0.4124	-2.7586	0.4136

Table V. Inspection of Table VI reveals that the two nuclei O₄ and C₃ are located within 0.2 Å of the principal axis *a*. Considerably larger errors must be accepted for their position than for the remaining nuclei in *s-trans*-acrolein.

Error Propagation in the Substitution Structures. Vibrational contributions to the rotational constants and experimental errors will especially affect the substitution coordinates when nuclei are close to principal axes. Taking differences Δ*I*_α between the moments of inertia of isotopically substituted species *I*_α' and those of the parent species *I*_α compensates the vibrational effects only to first order, and higher terms are not negligible for small coordinates. The variation of the inertia defect Δ in planar molecules upon isotopic substitution is a typical case showing such effects.

s-cis-Acrolein is a favorable molecule for structure determination. None of the nuclei is closer than 0.46 Å to the principal axes *a* or *b*. The situation in *s-trans*-acrolein is more critical, as pointed out above, since two nuclei are close to the principal axis *a*. Consideration of uncertainties in the structural parameters is indispensable in order to discuss structural differences between the two conformers.

The uncertainties in the substitution structure are mainly caused by the vibrational contributions to the ground-state moments of inertia. The experimental errors are almost negligible in our case. A recent discussion of the influence of molecular vibrations on the substitution coordinates has been given by van Eijck.²⁰ In the following we will not discuss the status of the substitution structure nor how close it will be to the equilibrium geometry. Instead we will try to assess the reliability of the substitution structures of *s-cis*- and *s-trans*-acrolein.

For planar molecules the isotope shift

$$\Delta P_z = P_z' - P_z \quad (1)$$

between the planar moment of inertia of the substituted species

$$P_z' = (I_a' + I_b' - I_c')/2 = -\Delta'/2 \quad (2a)$$

and the planar moment of the parent species

$$P_z = (I_a + I_b - I_c)/2 = -\Delta/2 \quad (2b)$$

is a measure of the internal consistency of the substitution coordinates. The isotope shift Δ*P*_z would vanish if the vibrational contributions to the moments of inertia were the same for substituted and parent molecule. The uncertainty σ(*x*) of the substitution coordinate *x* is estimated from the isotope shift Δ*P*_z of the planar moment of inertia²⁰

$$\sigma(x) = \Delta P_z / 2\Delta m|x| \quad (3)$$

for a specific substitution for which Δ*m* is the corresponding mass change. The uncertainties σ(*R*) of the bond lengths and bond angles follow from the uncertainties σ(*x*)

$$\sigma^2(R_i) = \sum_{j=1}^{3N} T_{ij}^2 \sigma^2(x_j) \quad (4)$$

with the help of the linearized transformation coefficients *T*_{ij} = ∂*R*_i/∂*x*_j between internal and Cartesian coordinates. *N* denotes

(18) Rudolph, H. D. *J. Mol. Spectrosc.* **1981**, *89*, 460.

(19) Nösberger, P.; Bauder, A.; Günthard, Hs. H. *Chem. Phys.* **1973**, *1*, 418.

(20) van Eijck, B. P. *J. Mol. Spectrosc.* **1982**, *91*, 348.

the number of nuclei in the molecule. The uncertainties $\sigma(R_i)$ are listed in Table V together with the bond lengths and bond angles determined from the Kraitchman substitution coordinates. They are very small for *s-cis*-acrolein but much larger for *s-trans*-acrolein. This reflects the larger isotope shifts ΔP_2 , as exemplified by Δ (cf. Table I and Table II) and the small coordinates for *s-trans*-acrolein as compared to those of the *s-cis* conformer.

The near cancellation of vibrational contributions to ΔP_2 for *s-cis*-acrolein might be accidental. Instead of using individual values, ΔP_2 average values $\langle \Delta P_2 \rangle^{1/2}$ for specific types of substitutions were introduced. Typical values for $\langle \Delta P_2 \rangle^{1/2}$ have been derived by averaging over many cases of substitutions in various molecules. The modified uncertainty $\sigma'(x)$ which is based on such an average value $\langle \Delta P_2 \rangle^{1/2}$ is given by

$$\sigma'(x) = \langle \Delta P_2 \rangle^{1/2} / 2\Delta m|x| = K/|x| \quad (5)$$

with $K = 0.0031 \text{ \AA}^2$ for H-D, $K = 0.0008 \text{ \AA}^2$ for ^{12}C - ^{13}C , and $K = 0.0006 \text{ \AA}^2$ for ^{16}O - ^{18}O substitutions.²⁰ A certain disadvantage of average K values is that the properties of individual molecules are not considered. It was recognized by van Eijck²⁰ that much smaller values of $\langle \Delta P_2 \rangle^{1/2}$ were found for rigid unsaturated ring molecules. The modified uncertainties $\sigma'(R_i)$ of the bond lengths and bond angles as calculated from eq 5 and 4 are also listed in Table V. Whereas $\sigma(R_i)$ and $\sigma'(R_i)$ are similar for *s-trans*-acrolein, $\sigma'(R_i)$ is always larger than $\sigma(R_i)$ for *s-cis*-acrolein.

For the least-squares method used to calculate the structure from the isotope shifts of the moments of inertia, the uncertainties are stated as three times the standard deviations which follow from the residuals $\Delta I_a(\text{exptl}) - \Delta I_a(\text{calcd})$ and the variance-covariance matrix. The standard deviations, however, are not very meaningful since the residuals do not represent statistical errors but systematic deviations due to vibrational effects. We feel that the least-squares method yields the most balanced estimate for the molecular structures because the center-of-mass and product-of-inertia conditions were taken into account. They are shown in Figure 1 with the dipole moments of *s-cis*-acrolein⁶ and of *s-trans*-acrolein included. The uncertainties, however, are more reasonably determined from eq 5 and 4.

Discussion and Conclusions

Inspection of Table V reveals small changes between the bond lengths and bond angles of *s-trans*-acrolein determined in this work and in the previous investigation of Cherniak and Costain.³ They used ΔI_b and ΔI_c in the two-dimensional equations of Kraitchman^{17a} since only the rotational constants B and C were available for the isotopic species at that time. This is equivalent to the neglect of variations of the inertia defect upon isotopic substitution. This work also shows that the inertia defect varies between -0.0075 and -0.0220 u\AA^2 for the isotopic species (cf. Table II) as compared to the value of -0.0212 u\AA^2 for the parent species. The larger part of the revision of the molecular structure resulted from these variations of the inertia defect rather than from inaccuracy of the previous values of the rotational constants. Further differences must be attributed to the fact that Cherniak and Costain³ assumed equal bond lengths for $\text{C}_3\text{-H}_7$ and $\text{C}_3\text{-H}_8$ in order to overcome the large uncertainty in the $b(\text{C}_3)$ coordinate. This assumption shifted the position of C_3 away from the principal axis a by 0.005 \AA . Such a procedure no longer seems justified in view of the corresponding results for *s-cis*-acrolein where different bond lengths for $\text{C}_3\text{-H}_7$ and $\text{C}_3\text{-H}_8$ were found.

The molecular structures of *s-cis*- and *s-trans*-acrolein are listed in Table V and are depicted in Figure 1 with the atomic numbering scheme. The results from the three-dimensional Kraitchman equations^{17a} and from the least-squares fit are nearly equal in both conformers. Valuable information about the effect of the conformation on the molecular geometry can be obtained from a comparison of the structural parameters. The $\text{C}=\text{O}$ and $\text{C}=\text{C}$ double bonds and the CCO angle are equal within the uncertainties for both conformers. The central $\text{C}-\text{C}$ single bond and the CCC angle increase by $0.010 (4) \text{ \AA}$ and $1.1 (4)^\circ$, respectively, in going from *s-trans*- to *s-cis*-acrolein. The methylene groups of both conformers show notable features. The $\text{C}_3\text{-H}_7$ bond is longer than the $\text{C}_3\text{-H}_8$ bond by $0.010 (12)$ and $0.017 (3) \text{ \AA}$ for *s-trans*- and *s-cis*-acrolein, respectively. This effect resembles the σ effect in substituted ethylenes.^{21,22} The $\text{C}_3\text{-H}_7$ bond of *s-cis*-acrolein is remarkably long.

Ab initio calculations of *s-cis* and *s-trans* conformers of 1,3-butadiene, acrolein, and glyoxal were reported by George et al.²³ They compared the results among the three isoelectronic molecules using 4-31G and (7,3) basis sets with full geometry optimization. Their calculated structures *s-cis*- and *s-trans*-acrolein are in reasonable agreement with our experimental data. The $\text{C}-\text{C}$ and $\text{C}-\text{H}$ bond lengths of the substitution structures are systematically longer by $0.003\text{--}0.022 \text{ \AA}$ than for the ab initio structures. The angles are usually consistent within 1° except for the angles $\text{H}_7\text{C}_3\text{H}_8$ and $\text{H}_7\text{C}_3\text{C}_2$ which show differences of more than 2° . George et al.²³ rationalized some unexpected deviations of *s-cis*-acrolein from the calculated trends in the series of the isoelectronic molecules in terms of a specific attractive interaction between the oppositely charged ends of acrolein.

Acknowledgment. Financial support by the Schweizerischer Nationalfonds (Project No. 2.407-0.82) is gratefully acknowledged. We are indebted to M. Rodler for kindly volunteering in the measurements of acrolein-*3-d* and for helpful discussions on the isotopic syntheses. We wish to thank Prof. R. L. Kuczkowski for a preprint¹¹ and for additional information on the preparation of acrolein-*3-d* and Dr. B. P. Winnewisser for critical reading of the manuscript. We are grateful to the Computer Center at the ETH Zurich for a generous grant of computer time.

Registry No. $^{13}\text{CH}_2\text{CHCHO}$, 10317-31-4; $\text{CH}_2^{13}\text{CHCHO}$, 10317-32-5; $\text{CH}_2\text{CH}^{13}\text{CHO}$, 10317-33-6; $\text{CH}_2\text{CHCH}^{18}\text{O}$, 10317-34-7; $\text{CH}_2\text{C-DCHO}$, 10317-37-0; CH_2CHCDO , 10317-38-1; *cis*-CHDCHCHO, 10317-35-8; acrolein, 107-02-8.

Supplementary Material Available: Listings of measured rotational transition frequencies of the *s-cis* and *s-trans* conformers of $^{13}\text{CH}_2=\text{CH}-\text{CHO}$, $\text{CH}_2=^{13}\text{CH}-\text{CHO}$, $\text{CH}_2=\text{CH}-^{13}\text{CH}-\text{O}$, $\text{CH}_2=\text{CH}-\text{CH}^{18}\text{O}$, $\text{CH}_2=\text{CD}-\text{CHO}$, $\text{CH}_2=\text{CH}-\text{CDO}$, *cis*-CHD=CH-CHO, and *trans*-CDH=CH-CHO in the vibrational ground state (Tables VII-XXII) (24 pages). Ordering information is given on any current masthead page.

(21) (a) Vinyl fluoride: Lide, D. R.; Christensen, D. *Spectrochim. Acta* **1961**, *17*, 665. (b) Vinyl chloride: Kivelson, D.; Wilson, E. B.; Lide, D. R. *J. Chem. Phys.* **1960**, *32*, 205. (c) Propylene: Lide, D. R.; Christensen, D. *J. Chem. Phys.* **1961**, *35*, 1374. (d) Nitroethylene: Nösberger, P.; Bauder, A.; Günthard, Hs. H. *Chem. Phys.* **1975**, *8*, 245. (e) Vinyl alcohol: Rodler, M.; Bauder, A. *J. Am. Chem. Soc.* **1984**, *106*, 4025-4028.

(22) McKean, D. C. *Spectrochim. Acta, Part A* **1975**, *31A*, 1167.

(23) George, P.; Bock, C. W.; Trachtman, M. *J. Mol. Struct.* **1980**, *69*, 183.