

for Developmental Scientific Research No. 59850142 from the Ministry of Education, Science and Culture.

Supplementary Material Available: Listings of positional and thermal parameters and mean square displacement tensors of atoms (2 pages). Ordering information is given on any current masthead page.

Hexakis(μ_3 -thio)pentakis[(η^5 -cyclopentadienyl)-titanium], [(η^5 -C₅H₅)Ti]₅(μ_3 -S)₆: Preparation and Molecular and Electronic Structure[†]


Frank Bottomley,* Gabriel O. Egharevba, and Peter S. White

Department of Chemistry
University of New Brunswick
Fredericton, New Brunswick, Canada E3B 6E2

Received February 12, 1985

In earlier work we have described [(CpCr)₄(μ_3 -O)₄] and [(CpV)₅(μ_3 -O)₆] (Cp = η^5 -C₅H₅),¹ which together with [(CpTi)₆(μ_3 -O)₈] described by Caulton and co-workers² provide a series of novel oxygen-containing clusters which obey Euler's theorem. A theoretical study of these clusters showed that there were 12 cluster orbitals occupied by 2 ((CpTi)₆O₈), 8 ((CpV)₅O₆), or 12 ((CpCr)₄O₄) electrons.³ According to this analysis a wide variety of more or less distorted octahedral (CpM)₆A₈, trigonal-bipyramidal (CpM)₅A₆, and tetrahedral (CpM)₄A₄ clusters (M = first-row transition metal and A = μ_3 -atom from groups 15 or 16) should be obtainable. In the tetrahedral case several thio derivatives [(CpM)₄S₄]ⁿ⁺ (M = Cr, Fe, Co) are known.⁴ We have been seeking other members of these series, particularly nontetrahedral derivatives, and report here the preparation and structure of the first trigonal-bipyramidal thio cluster, [(CpTi)₅(μ_3 -S)₆].⁵

In their report on (CpTi)₆O₈ Caulton and co-workers suggested that the cluster was actually obtained from Cp₂Ti(CO)₂ and H₂O.² We have confirmed this suggestion and find that the reaction between Cp₂Ti(CO)₂ and H₂O (3:4 mol ratio) in toluene at 80 °C is quantitative and gives (CpTi)₆O₈ as the *only* titanium-containing product according to the equation:



This remarkable specificity suggested that a similar reaction with H₂S would take place. It does, but the product is (CpTi)₅(μ_3 -S)₆ (73% yield based on the Cp₂Ti(CO)₂ used) not (CpTi)₆S₈, thus illustrating our contention that the clusters obeying Euler's theorem

[†] In this paper the periodic group notation is in accord with recent actions by IUPAC and ACS nomenclature committees. A and B notation is eliminated because of wide confusion. Groups IA and IIA become groups 1 and 2. The d-transition elements comprise groups 3 through 12, and the p-block elements comprise groups 13 through 18. (Note that the former Roman number designation is preserved in the last digit of the new numbering: e.g., III → 3 and 13.)

(1) Bottomley, F.; Paez, D. E.; White, P. S. *J. Am. Chem. Soc.* **1982**, *104*, 5651.

(2) Huffman, J. C.; Stone, J. G.; Krusell, W. C.; Caulton, K. G. *J. Am. Chem. Soc.* **1977**, *99*, 5829.

(3) Bottomley, F.; Grein, F. *Inorg. Chem.* **1982**, *12*, 4170.

(4) Simon, G. L.; Dahl, L. F. *J. Am. Chem. Soc.* **1973**, *95*, 2164. Simon, G. L.; Dahl, L. F. *J. Am. Chem. Soc.* **1973**, *95*, 2175. Trinh-Toan; Teo, B. K.; Ferguson, J. A.; Meyer, T. J.; Dahl, L. F. *J. Am. Chem. Soc.* **1977**, *99*, 408. Pasyanski, A. A.; Eremenko, I. L.; Orzaksakhatov, B.; Kalinnikov, V. T.; Aleksandrov, G. G.; Struchkov, Yu. T. *J. Organomet. Chem.* **1981**, *216*, 211.

(5) [Cu₅(μ_2 -S-*t*-Bu)₆]⁺ contains a trigonal bipyramid of Cu atoms enclosed within an open octahedron of μ_2 -S-*t*-Bu ligands: Dance, I. G. *J. Chem. Soc., Chem. Commun.* **1976**, 68.




Figure 1. ORTEP drawing of (CpTi)₅S₆. Hydrogen atoms are omitted for clarity. Ti-Ti distances: Ti(1)-Ti(2) 3.152 (5); Ti(1)-Ti(3) 3.148 (5); Ti(1)-Ti(4) 3.214 (5); Ti(1)-Ti(5) 3.145 (5); Ti(2)-Ti(3) 3.173 (5); Ti(2)-Ti(4) 3.169 (5); Ti(3)-Ti(4) 3.172 (5); Ti(3)-Ti(5) 3.161 (5); Ti(4)-Ti(5) 3.076 (5) Å. Ti-S distances: Ti(1)-S(3) 2.468 (6); Ti(1)-S(4) 2.485 (7); Ti(1)-S(5) 2.480 (7); Ti(1)-S(6) 2.463 (7); Ti(2)-S(2) 2.274 (6); Ti(2)-S(3) 2.308 (6); Ti(2)-S(4) 2.277 (7); Ti(3)-S(1) 2.443 (7); Ti(3)-S(2) 2.506 (7); Ti(3)-S(3) 2.461 (6); Ti(3)-S(5) 2.444 (7); Ti(4)-S(1) 2.458 (6); Ti(4)-S(2) 2.487 (7); Ti(4)-S(4) 2.488 (6); Ti(4)-S(6) 2.504 (8); Ti(5)-S(1) 2.286 (7); Ti(5)-S(5) 2.268 (7); Ti(5)-S(6) 2.263 (8) Å.





Figure 2. ESR spectrum of (CpTi)₅S₆ at 20 °C.

are closely related. Analysis of the gaseous and solid products of the reaction between Cp₂Ti(CO)₂ (6.4 mmol) and gaseous H₂S (7.7 mmol) in toluene at 80 °C for 72 h suggests that the stoichiometry of the reaction producing (CpTi)₅S₆ is



though other sulfur-containing products are obtained. The new cluster is dark green-brown, moderately air sensitive, very soluble in toluene and forms large crystals when the toluene solutions are layered with hexane.⁶

(6) Analyses of the bulk material (powder) shows it contains one molecule of toluene of crystallization per (CpTi)₅S₆ unit; this toluene is not present in the crystal examined by X-ray diffraction. Analytical results (by Beller Laboratorium, Göttingen, W. Germany) of samples from two distinct preparations: Calcd for (CpTi)₅S₆·C₆H₆·CH₃ (C₃₃H₃₃S₆Ti₅): C, 45.2; H, 3.9; S, 22.6; Ti, 28.2%. Found: C, 45.7, 45.6; H, 4.2, 4.3; S, 15.2, 10.8; Ti 27.7, 28.7%. The problem of incorrect sulfur analyses in cyclopentadienyl metal complexes has been discussed by other workers.⁷

The crystal structure determination⁸ shows that (CpTi)₅S₆ contains sulfur bridging the triangular faces of a distorted trigonal bipyramid of titanium atoms (Figure 1). The distortion takes the form of a displacement of one of the equatorial titanium atoms (Ti(4) in Figure 1) away from one other equatorial titanium (Ti(1)) and toward an axial titanium (Ti(5)). The distance Ti(4)-Ti(1) is 3.214 (5) Å, Ti(4)-Ti(5) is 3.076 (5) Å, and the average for all other Ti-Ti distances is 3.160 (5) Å with a range from 3.145 (5) to 3.173 (5) Å. All the Ti-Ti distances are longer than those of (CpTi)₆O₈ (average Ti-Ti 2.891 (1) Å²) because of the larger covalent radius of sulfur (1.02 Å) vs. oxygen (0.73 Å); (CpV)₅O₆ is an essentially perfect trigonal bipyramid with V-V distances ranging from 2.738 (3) to 2.762 (2) Å.¹

The new cluster is paramagnetic, as would be expected for a species with an odd number of electrons. The ESR spectrum (toluene solution) shows a single line at $g = 1.993$ with weak satellites attributable to hyperfine interaction with the Ti isotopes (Figure 2). The spectrum at -80 °C is essentially identical with that at 20 °C, though the satellites are not observed at the lower temperature. The bulk magnetic susceptibility per mole at 20 °C is 6.525×10^{-4} . This corresponds to a magnetic moment of 1.25 μ_B , which is low compared to the spin-only value for one electron of 1.73 μ_B due mainly to the lack of a diamagnetic correction for any of the ligands or titanium atoms.⁹ However, both the narrow line ESR spectrum and the bulk magnetic moment indicate the presence of one unpaired electron in (CpTi)₅S₆.

Both the distortion of (CpTi)₅S₆ and its one unpaired electron can be understood in terms of the theoretical model presented previously.³ The new cluster has three electrons to be accommodated in the 12 cluster orbitals. In a perfect trigonal-bipyramidal structure (D_{3h} symmetry) the electron configuration would be $(1a'_1)^2(1e')^1$. Therefore Jahn-Teller distortion of the D_{3h} structure is expected. Since both the $1a'_1$ and $1e'$ orbitals are almost completely localized on the equatorial titanium atoms, the distortion must involve these atoms, as is observed. The distortion is such that (CpTi)₅S₆ has no symmetry within experimental error.

Since the three cluster electrons are localized on the equatorial titanium atoms, one may naively consider each of these atoms to be Ti(III) (d^1) and the axial titanium atoms to be Ti(IV) (d^0). This assignment of oxidation states is in agreement with the observed Ti-S distances. The equatorial Ti-S distances average 2.474 (7) Å with a range from 2.443 (7) to 2.506 (7) Å; the axial Ti-S distances are much shorter, averaging 2.279 (7) Å with a range from 2.263 (8) to 2.308 (6) Å. The Ti-S distances may be compared to those in Cp*₂TiS₃ (Cp* = η^5 -C₅(CH₃)₅) and Cp₂TiS₅, 2.413 (4) Å^{10a} and 2.420 and 2.446 Å.^{10b} The Ti(4)-S distances are identical with those of the other equatorial titanium atoms (Ti(1)-S and Ti(3)-S); the trigonal pyramid of sulfur atoms is distorted with the trigonal bipyramid of titanium atoms in order to retain the Ti-S bonding. This is in accord with the theoretical model in which the 12 cluster orbitals of (CpM)₅A₆ are essentially pure metal orbitals and are nonbonding with respect to the M-A interaction.

Analogous reactions to those between Cp₂Ti(CO)₂ and H₂O or H₂S do not occur with Cp₂Zr(CO)₂ presumably because of the instability of the Zr(III) oxidation state. On reaction with H₂O,

Cp₂Zr(CO)₂ gives only ZrO₂ and with H₂S the product is the known⁷ complex (Cp₂ZrS)₂.

Acknowledgment. We thank the Natural Sciences and Engineering Research Council of Canada and the donors of the Petroleum Research Fund, administered by the American Chemical Society, for financial support of this work. G.O.E. thanks the University of Ife (Nigeria) for the grant of a study leave. Colin Mailer and Rod McGregor are thanked for assistance with the ESR spectra.

Supplementary Material Available: Tables of atomic coordinates, thermal parameters, and bond lengths (7 pages). Ordering information is given on any current masthead page.

Excited-State Chemistry of Tetrakis(μ -pyrophosphito)diplatinum(II). Photoinduced Addition of Aryl Bromides and Iodides to the Binuclear Complex and the Photoinduced Catalytic Conversion of Isopropyl Alcohol into Acetone and Hydrogen

D. Max Roundhill

Department of Chemistry, Tulane University
New Orleans, Louisiana 70118

Received September 13, 1984

Aqueous solutions of the binuclear platinum(II) complex Pt₂(μ -P₂O₅H₂)₄⁴⁺ show an intense absorption band at 367 nm (ϵ_{\max} 3.45×10^4) and a corresponding emission at 514 nm.¹ This emission has been identified as a long-lived phosphorescence ($\tau = 9.8$ (2) μ s) originating from a triplet excited state.² Using a simplified MO treatment, the respective ground- and excited-state configurations of these binuclear d^8 complexes are represented as $(d_{z^2})^2(d_{x^2-y^2})^2$ and $(d_{z^2})^2(d_{x^2-y^2})^1(p_z)^1$. The excited state Pt₂(μ -P₂O₅H₂)₄⁴⁺ undergoes efficient energy quenching with SO₂.³ Quenching is also observed with electron-transfer reagents, and from experiments using different oxidants the electrode potential for Pt₂(μ -P₂O₅H₂)₄⁴⁺/Pt₂(μ -P₂O₅H₂)₄³⁺ has been estimated to be < -1 V.⁴ This long microsecond lifetime for Pt₂(μ -P₂O₅H₂)₄⁴⁺ makes it conceptually feasible to carry out reaction chemistry with this triplet excited state, and its strong reducing properties makes it a useful reagent for reaction with organic halides. This paper reports the first organometallic reactions of Pt₂(μ -P₂O₅H₂)₄⁴⁺ and relates its redox behavior and radical character to the Gray MO model.

The complex Pt₂(μ -P₂O₅H₂)₄⁴⁺ undergoes thermal addition of halogens X₂ and methyl iodide to give the axially disubstituted Pt(III)-Pt(III) complexes Pt₂(μ -P₂O₅H₂)₄X₂⁴⁺ and Pt₂(μ -P₂O₅H₂)₄MeI⁴⁺.⁵ Thermal addition of the higher homologues RI (R = Et, *n*-Pr, *i*-Pr, *n*-pent) also gives Pt₂(μ -P₂O₅H₂)₄RI⁴⁺, but the complex is contaminated with Pt₂(μ -P₂O₅H₂)₄I₂⁴⁺.⁶ Aryl halides do not undergo thermal addition. Aryl bromides or iodides

(7) Shaver, A.; McCall, J. M. *Organometallics* 1984, 3, 1823.

(8) Data for (C₅H₅)Ti₅S₆ ($M = 757.4$) were obtained from a wedge-shaped chip cut from a large irregular crystal and were measured on a Picker FACS-1 diffractometer. Space group $P2_1/n$, $a = 16.978$ (5) Å, $b = 17.008$ (9) Å, $c = 10.122$ (3) Å, $\beta = 92.44$ (2)°, $Z = 4$, $D_c = 1.72$, 1383 observed ($I > 3\sigma(I)$) reflections out of 3828 with $2\theta \leq 45^\circ$. The data were corrected for absorption and all atoms except H were refined anisotropically to a conventional R of 0.074, $R_w = 0.077$. The structure was solved by MULTAN80 and refined by SHELX. Tables of atomic positions, thermal parameters, and bond lengths are available as supplementary material. Attempts to obtain crystals that diffract better are continuing.

(9) Correction for the diamagnetism of 5C₅H₅⁻ ($-50.9 \times 10^{-6} \times 5$), 6S²⁻ ($-38 \times 10^{-6} \times 6$), 2Ti⁴⁺ ($-5 \times 10^{-6} \times 2$), and 3Ti³⁺ ($-9.2 \times 10^{-6} \times 3$) gives a moment of 1.66 μ_B . See: Weiss, A.; Witte, H. "Magnetochemie"; Verlag Chemie: Weinheim, 1973; pp 93-95.

(10) (a) Bird, P. H.; McCall, J. M.; Shaver, A.; Siriwardane, U. *Angew. Chem., Int. Ed. Engl.* 1982, 21, 384. (b) Epstein, E. E.; Bernal, I. *J. Chem. Soc., Chem. Commun.* 1970, 410.

(1) Sperline, R. P.; Dickson, M. K.; Roundhill, D. M. *J. Chem. Soc., Chem. Commun.* 1977, 62-63.

(2) Che, C.-M.; Butler, L. G.; Gray, H. B. *J. Am. Chem. Soc.* 1981, 103, 7796-7797. Fordyce, W. A.; Brummer, J. G.; Crosby, G. A. *J. Am. Chem. Soc.* 1981, 103, 7061-7064. Markert, J. T.; Clements, D. P.; Carson, M. R.; Nagle, J. K. *Chem. Phys. Lett.* 1983, 97, 175-179. Bar, L.; Gliemann, G. *Chem. Phys. Lett.* 1984, 108, 14-17.

(3) Alexander, K. A.; Stein, P.; Hedden, D.; Roundhill, D. M. *Polyhedron* 1983, 2, 1389-1392.

(4) Heuer, W. B.; Totten, M. D.; Rodman, G. S.; Hebert, E. J.; Tracy, H. J.; Nagle, J. K. *J. Am. Chem. Soc.* 1984, 106, 1163-1164. Vogler, A.; Kunkely, H. *Angew. Chem., Int. Ed. Engl.* 1984, 23, 316-317.

(5) Che, C.-M.; Schaefer, W. P.; Gray, H. B.; Dickson, M. K.; Stein, P.; Roundhill, D. M. *J. Am. Chem. Soc.* 1982, 104, 4253-4255.

(6) UV-vis data for Pt₂(μ -P₂O₅H₂)₄RI⁴⁺: λ_{\max} 352 (R = Et); 355 nm (R = *i*-Pr, *n*-Pr, *n*-pent). UV-vis data for Pt₂(μ -P₂O₅H₂)₄X₂⁴⁺: λ_{\max} 338 (X = I), 305 nm (X = Br): from ref 5.