

oxygen (step  $k_2$ ) should respond to both sterics and electronic changes for R. The higher stability of the phenyl derivative may also be accommodated with this mechanism, since the planar phenyl ring is constrained to lie in the equatorial plane of the bent sandwich structure,<sup>15</sup> thus discouraging formation of reactive  $[\text{Cp}^*_2\text{Hf}(\text{C}_6\text{H}_5)(\eta^2\text{-OOCMe}_3)]$ . Although the range of solvent polarities is severely limited by the high reactivity of these compounds, no significant difference was noted for the rate of rearrangement of  $\text{Cp}^*_2\text{Hf}(\text{CH}_2\text{CH}_3)(\text{OOCMe}_3)$  among the solvents listed (eq 6), suggesting little polar character for the transition state of step  $k_2$ .

The crowded ligand environment surrounding a less reactive, third-row transition metal is very likely responsible for the thermal stability exhibited by members of this series of complexes. These same features appear to dictate the mechanism(s) for rearrangement to the very stable bis(alkoxide) derivatives.

**Acknowledgment.** This work was supported by the National Science Foundation (Grant CHE-8303735), by the USDOE Office of Energy Research, Office of Basic Energy Sciences (Grant DE-FG03-85ER13431), by Du Pont, and by Shell Companies Foundation, which are gratefully acknowledged. National Science Foundation Grant CHE-8219039 to the divisional X-ray diffraction facilities is also gratefully acknowledged.

**Supplementary Material Available:** Tables of analytical, NMR, and IR spectroscopic data for all isolated compounds, atomic coordinates, thermal parameters, bond lengths & angles, least-squares planes, and torsion angles and experimental procedures and crystal structure determination (12 pages); structure factor tables for  $\text{Cp}^*_2\text{Hf}(\text{C}_2\text{H}_5)(\text{OOC}_4\text{H}_9)$  (10 pages). Ordering information is given on any current masthead page.

(15) Although the crystal was disordered, a partial X-ray structure determination for  $\text{Cp}^*_2\text{Hf}(\text{C}_6\text{H}_5)(\text{OOCMe}_3)$  revealed this arrangement for the phenyl group. The barrier for aryl rotation is sufficiently large that NMR spectra indicate this static structure in solution at ambient temperature.

## Oxygenate Formation from Electrophilic Attack of a Rh/Zr "A-Frame" Containing a $\mu_2\text{-}\eta^2$ -Acetyl Bridge

Gregory S. Ferguson and Peter T. Wolczanski\*

Department of Chemistry, Baker Laboratory  
Cornell University, Ithaca, New York 14853


Received July 28, 1986

The current economic climate has shifted the emphasis in Fischer-Tropsch (F-T) research from hydrocarbons to oxygenates, useful precursors to a variety of chemicals. While  $\text{C}_1$  chemistry, principally MeOH based, has reached the commercial stage,<sup>1</sup> the selective formation of higher oxygenates remains a challenging problem.<sup>2</sup> Features critical to such selectivity are exhibited by supported-Rh catalysts:<sup>3</sup> (1) CO adsorption and dissociation are suppressed by strong metal support interactions (SMSI);<sup>4-6</sup> (2)

Rh engenders facile CO insertions;<sup>7</sup> (3) IR studies are indicative of  $\text{CO}_{\text{ads}}$ ,<sup>8,9</sup> acyl, and acetate<sup>10,11</sup> species that interact with the support; (4) MeOH and  $\text{C}_2$  products are formed from different sites.<sup>5,12,13</sup> Mechanistic interpretations converge on a key migration of surface Me to  $\text{CO}_{\text{ads}}$  occurring at interfacial Rh/support sites.<sup>5,10,12,14-16</sup>

Organometallic species have proven valuable in modeling<sup>17-19</sup> spectroscopic and reactivity properties of hydrocarbon F-T catalysts.<sup>20</sup> Considering highly  $\text{C}_2$ -oxygenate-selective Rh/ZrO<sub>2</sub> as representative<sup>21,22</sup> of heterogeneous oxygenate catalysts, the synthesis of a "homogenous Rh/Zr interface" was deemed appropriate. Herein is described the preparation of an "A-frame"<sup>23</sup> complex which produces oxygenates when subjected to electrophiles.

The addition of 2 equiv of  $\text{HOCH}_2\text{Ph}_2\text{P}^{24}$  to  $\text{Cp}^*\text{ZrMe}_3$  (**1**,  $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$ )<sup>25</sup> afforded white crystalline  $\text{Cp}^*\text{MeZr}(\text{OCH}_2\text{Ph}_2\text{P})_2$  (**2**)<sup>26</sup> in good yield (89%, eq 1). Treatment of **1** with 0.5 equiv



(4) (a) *Metal-Support and Metal Additive Effects in Catalysis*; Imelik, B., et al., Eds.; Elsevier: Amsterdam, 1982. (b) Tauster, S. J.; Fung, S. C.; Garten, R. L. *J. Am. Chem. Soc.* **1978**, *100*, 170-175. (c) Tauster, S. J.; Fung, S. C.; Baker, R. T. K.; Horsley, J. A. *Science (Washington, D.C.)* **1981**, *211*, 1121-1125 and references therein.

(5) van der Lee, G.; Schuller, B.; Post, H.; Favre, T. L. F.; Ponec, V. *J. Catal.* **1986**, *98*, 522-529.

(6) Takeuchi, A.; Katzer, J. R. *J. Phys. Chem.* **1982**, *86*, 2438-2441.

(7) Chuang, S. C.; Tian, Y. H.; Goodwin, J. G., Jr.; Wender, I. *J. Catal.* **1985**, *96*, 396-407.

(8) (a) Sachtler, W. M. H.; Shriver, D. F.; Hollenberg, W. B.; Lang, A. F. *J. Catal.* **1985**, *92*, 429-431. (b) Horwitz, C. P.; Shriver, D. F. *Adv. Organomet. Chem.* **1984**, *23*, 219-305.

(9) Ichikawa, M.; Sekizawa, S.; Shikakura, K.; Kawai, M. *J. Mol. Catal.* **1981**, *11*, 167-179.

(10) Orita, H.; Naito, S.; Tamaru, K. *J. Catal.* **1984**, *90*, 183-193.

(11) Fukushima, T.; Arakawa, H.; Ichikawa, M. *J. Chem. Soc., Chem. Commun.* **1985**, 729-731.

(12) Ichikawa, M.; Fukushima, T. *J. Chem. Soc., Chem. Commun.* **1985**, 321-323.

(13) Takeuchi, A.; Katzer, J. R. *J. Catal.* **1983**, *82*, 351-354.

(14) van der Lee, G.; Ponec, V. *J. Catal.* **1986**, *99*, 511-512.

(15) Sachtler, W. M. H.; Shriver, D. F.; Ichikawa, M. *J. Catal.* **1986**, *99*, 513-514.

(16) (a) Mims, C. A.; McCandlish, L. E. *J. Am. Chem. Soc.* **1985**, *107*, 696-697. (b) Zhang, X.; Biloen, P. *J. Catal.* **1986**, *98*, 468-476. (c) Vannice, M. A.; Sudhaker, C. *J. Phys. Chem.* **1984**, *88*, 2429-2432.

(17) (a) Herrmann, W. A. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 117-130. (b) Muettterties, E. L.; Stein, J. *Chem. Rev.* **1979**, *79*, 479-490. (c) Muettterties, E. L.; Rhodin, T. N.; Band, E.; Brucker, C. F.; Pretzer, W. R. *Ibid.* **1979**, *79*, 91-137. (d) Gladysz, J. A. *Adv. Organomet. Chem.* **1982**, *20*, 1-38. (e) Bradley, J. S. *Ibid.* **1983**, *22*, 1-58. (f) Wolczanski, P. T.; Bercaw, J. E. *Acc. Chem. Res.* **1980**, *13*, 121-127.

(18) Casey, C. P.; Palermo, R. E.; Rheingold, A. L. *J. Am. Chem. Soc.* **1986**, *108*, 549-550 and references therein.

(19) For other  $\mu_2\text{-}\eta^2\text{-O}(\text{R})\text{C}$  units, see: Marsella, J. A.; Huffman, J. C.; Caulton, K. G.; Longato, B.; Norton, J. R. *J. Am. Chem. Soc.* **1982**, *104*, 6360-6368 and references therein.

(20) (a) Falbe, J. *Chemical Feedstocks from Coal*; Wiley: New York, 1981. (b) Rofer-DePoorter, C. K. *Chem. Rev.* **1981**, *81*, 447-474. (c) Biloen, P.; Sachtler, W. M. H. *Adv. Catal.* **1981**, *30*, 165-216. (d) Bell, A. T. *Catal. Rev.—Sci. Eng.* **1981**, *23*, 203-232.

(21) Ichikawa, M. *Bull. Chem. Soc. Jpn.* **1978**, *51*, 2273-2277.

(22) Dall'Agnol, C.; Gervasini, A.; Morazzoni, F.; Pinna, F.; Strukul, G.; Zanderighi, L. *J. Catal.* **1985**, *96*, 106-114.

(23) (a) Puddephatt, R. *J. Chem. Soc. Rev.* **1983**, *12*, 99-127. (b) Balch, A. In *Homogeneous Catalysis with Metal Phosphine Complexes*; Pignolet, L. H., Ed.; Plenum: New York, 1983.

(24) Hellmann, v. H.; Bader, J.; Brikner, H.; Schumacher, O. *Justus Liebig's Ann. Chem.* **1962**, *659*, 49-63.


(25) (a) Ferguson, G. S.; Wolczanski, P. T. *Organometallics* **1985**, *4*, 1601-1605. (b) Wolczanski, P. T.; Bercaw, J. E. *Organometallics* **1982**, *1*, 793-799.

(1) (a) Haggin, J. *Chem. Eng. News* **1986**, *64*, 7-13. (b) Klier, K. *Adv. Catal.* **1982**, *31*, 243-313. (c) Chang, C. D. *Catal. Rev.—Sci. Eng.* **1983**, *25*, 1-118.


(2) (a) Poels, E. K.; Ponec, V. *Catalysis*; Specialist Periodic Reports, Vol. 6; Chemical Society: London, 1983; p. 196. (b) Dombek, B. D. *Adv. Catal.* **1983**, *32*, 325-416.

(3) For leading references, see: Ichikawa, M.; Lang, A. J.; Shriver, D. F.; Sachtler, W. M. H. *J. Am. Chem. Soc.* **1985**, *107*, 7216-7218.

Scheme I



of  $[(\text{OC})_2\text{RhCl}]_2$  (**3**) provided the  $\mu_2$ - $\eta^2$ -acetyl<sup>18,19</sup> "A-frame"  $\text{Cp}^*\text{Zr}(\mu\text{-OCH}_2\text{Ph}_2\text{P})_2(\mu_2\text{-}\eta^2\text{-O}=\text{C}(\text{CH}_3))(\mu\text{-Cl})\text{RhCO}$  (**4**), as yellow crystals (75%, eq 2). NMR and IR spectra<sup>27</sup> are consistent



with the structure indicated in Scheme I. Inequivalent methylene protons and a clean  $^{31}\text{P}\{\text{H}\}$  doublet ( $\delta$  22.92,  $J_{\text{RHP}} = 149$  Hz) characterized the *trans*- $\text{PPh}_2\text{CH}_2\text{O}$  bridges. Coupling constants [ $^{13}\text{C}\{\text{H}\}$   $\delta$  195.18 (RhCO,  $J_{\text{CP}} = 15$ ,  $J_{\text{CRh}} = 52$ ,  $J_{\text{CC}} = 29$  Hz), 309.22 ( $(\text{CH}_3)\text{C}=\text{O}$ ,  $J_{\text{CP}} = 11$ ,  $J_{\text{CRh}} = 30$ ,  $J_{\text{CC}} = 29$  Hz)] consistent with a *tbp* Rh core were obtained through  $^{13}\text{CO}$  exchange of the terminal CO to give **4**- $^{13}\text{C}$  and use of 50%  $^{13}\text{CO}$ -enriched **3** to yield **4**- $^{13}\text{C}_2$  (statistically 25%). The weak IR band at 1490  $\text{cm}^{-1}$  ( $\nu(^{13}\text{C}=\text{O}) = 1460$   $\text{cm}^{-1}$ ) and the  $\delta$  309.22  $^{13}\text{C}$  chemical shift are in accord with a  $\mu_2$ - $\eta^2$ -acetyl that incorporates a strong contribution from the alternate  $\mu_2$ - $\eta^2$ -oxycarbene resonance form (**4'**).<sup>8,18,19,28</sup> Inspection of molecular models indicated that the Cl is likely to be proximate to Rh. A-frame **4** was presumably produced via cleavage of **3** by **2**, chelate formation, and subsequent insertion of free or bound CO into a ZrMe or RhMe (via chloride for Me binuclear exchange)<sup>29</sup> bond accompanied by acyl rearrangement. The Rh(I) center linked to the alkoxyzirconium moiety appropriately models a heterogeneous Rh/ZrO<sub>2</sub> interface while avoiding intimate Rh/Zr contact.

The reactivity of **4** with electrophiles is illustrated in Scheme I. Exposure of **4** to 1 equiv of HCl resulted in the formation of acetaldehyde (61% by GC/MS,  $^1\text{H}$  NMR),<sup>19</sup> concomitant with  $\text{Cp}^*\text{Zr}(\mu\text{-OCH}_2\text{Ph}_2\text{P})_2(\mu\text{-Cl})_2\text{RhCO}$  (**5**, 68%).<sup>30</sup> A convenient synthesis of **5** was accomplished through reflux of a  $\text{CHCl}_3$  solution of **4**.<sup>31</sup> Treatment of **4** with methyl iodide (1 equiv) provided acetone (89%) in addition to  $\text{Cp}^*\text{Zr}(\mu\text{-OCH}_2\text{Ph}_2\text{P})_2(\mu\text{-Cl})(\mu\text{-I})\text{RhCO}$  (**6**, 85%),<sup>32</sup> which could also be prepared from NaI and **5**. When **4**- $^{13}\text{C}$  was subjected to HCl and  $\text{CH}_3\text{I}$ , the respective  $\text{CH}_3\text{CHO}$  and  $(\text{CH}_3)_2\text{CO}$ <sup>33</sup> products contained >98%  $^{12}\text{C}$  ( $^1\text{H}$  NMR, IR) in accord with direct cleavage of the bridge. The similarity in reaction conditions and products suggested that both electrophilic cleavages arise from attack at Rh (A). Tentative spectroscopic evidence for the putative intermediate resulting from HCl attack was obtained. A rhodium hydride [ $\delta$  -15.21 (dt,  $J_{\text{RHH}} = 28$ ,  $J_{\text{PH}} = 9$  Hz)], inequivalent methylene protons, a new acetyl ( $\delta$  1.05), and *t*- $\text{PPh}_2\text{CH}_2\text{O}$  bridges [ $^{31}\text{P}\{\text{H}\}$  NMR  $\delta$  14.26 ( $J_{\text{RHP}} = 98$  Hz)] implicated a structure such as  $[\text{Cp}^*\text{Zr}(\mu\text{-OCH}_2\text{Ph}_2\text{P})_2(\mu_2\text{-}\eta^2\text{-O}=\text{C}(\text{CH}_3))(\mu\text{-Cl})\text{Rh}(\text{CO})\text{H}]\text{Cl}$  (A)<sup>34</sup> or a similar nonionic formulation. Corresponding CO stretches at 2097 and 1542  $\text{cm}^{-1}$  intimated that the intermediate contains a Rh(III) center<sup>35</sup> with an intact  $\mu_2$ - $\eta^2$ -acetyl bridge.

Despite strong binding to the Zr, the  $\mu_2$ - $\eta^2$ -acetyl undergoes facile reductive elimination upon protolysis, implying that a similar pathway is reasonable for heterogeneous  $\text{H}_3\text{CCHO}$  formation.<sup>3</sup> It has been suggested that surface acetates may be penultimate intermediates on the pathway to acetaldehyde.<sup>5,10</sup> Note that the

(26) Anal. Calcd for **2**,  $\text{ZrP}_2\text{O}_7\text{C}_37\text{H}_{42}$ : C, 66.14; H, 6.30. Found: C, 65.86; H, 6.22.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ )  $\delta$  0.02 (s,  $\text{CH}_3$ , 3 H), 1.82 (s,  $\text{Cp}^*$ , 15 H), 4.77 (d,  $\text{CH}_2$ , 4 H,  $^2J_{\text{PH}} = 7.4$  Hz), 7.11 (m, Ph, 8 H), 7.53 (m, Ph, 12 H);  $^{31}\text{P}\{\text{H}\}$  NMR  $\delta$  -11.24;  $^{13}\text{C}\{\text{H}\}$  NMR  $\delta$  10.59 ( $\text{Cp}^*$ ), 27.76 ( $\text{H}_3\text{C}$ ), 71.47 (d,  $\text{CH}_2$ ,  $J_{\text{PC}} = 9.9$  Hz), 118.64 ( $\text{Cp}^*$ ), 128.54 (Ph-C(3)), 128.83 (Ph-C(4)), 133.56 (d, Ph-C(2)),  $^2J_{\text{PC}} = 17.6$  Hz), 137.68 (dd, Ph-C(1)),  $^1J_{\text{PC}} = 14.3$ ,  $J_{\text{PC}} = 2.2$  Hz).

(27) Anal. Calcd for **4**,  $\text{RhZrClP}_2\text{O}_7\text{C}_39\text{H}_{42}$ : C, 54.07; H, 4.89; Cl, 4.09. Found: C, 53.86; H, 5.04; Cl, 4.44.  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ )  $\delta$  1.47 (t,  $\text{CH}_3$ , 3 H,  $^4J_{\text{PH}} = 1.3$  Hz), 2.05 (s,  $\text{Cp}^*$ , 15 H), 4.66 (ddt(v),  $\text{CHH}$ , 2 H,  $^2J = 11.4$ ,  $^3J_{\text{RHH}} \sim 1$ ,  $J_{\text{PH}} = 3$  Hz), 5.05 (ddt(v),  $\text{CHH}$ , 2 H,  $^2J = 11.4$ ,  $^3J_{\text{RHH}} \sim 2$ ,  $J_{\text{PH}} \sim 1$  Hz), 7.39 (m, Ph, 12 H), 7.63 (m, Ph, 8 H);  $^{31}\text{P}\{\text{H}\}$  NMR  $\delta$  22.92 (d,  $J_{\text{RHP}} = 149$  Hz);  $^{13}\text{C}$  NMR  $\delta$  11.44 ( $\text{Cp}^*$ ), 40.16 ( $\text{H}_3\text{C}$ ), 72.53 (t(v),  $\text{CH}_2$ ,  $J_{\text{PC}} = 18$  Hz), 121.58 ( $\text{Cp}^*$ ), 128-134 (Ph), 195.18 (CO ( $4\text{-}^{13}\text{C}_2$ )), ddt,  $J_{\text{RHC}} = 52$ ,  $J_{\text{CC}} = 29$ ,  $^2J_{\text{CP}} = 15$  Hz), 309.22 ( $\text{C}(\text{CH}_3)\text{O}$ - ( $4\text{-}^{13}\text{C}_2$ )), ddt,  $^2J_{\text{RHC}} = 30$ ,  $^2J_{\text{CC}} = 29$ ,  $^2J_{\text{PC}} = 11$  Hz); IR ( $4/4\text{-}^{13}\text{C}_2$ ,  $\text{CD}_2\text{Cl}_2$ )  $\nu(\text{CO})$  2014/1969  $\text{cm}^{-1}$ ,  $\nu(\text{H}_3\text{CC}=\text{O})$  1490/1460  $\text{cm}^{-1}$ . Conductivity measurements were consistent with a neutral formulation.

(28) Butts, S. B.; Strauss, S. H.; Holt, E. M.; Stimson, R. E.; Alcock, N. W.; Shriver, D. F. *J. Am. Chem. Soc.* **1980**, *102*, 5093-5100.

(29) (a) Puddephatt, R. J.; Stalteri, M. A. *Organometallics* **1983**, *2*, 1400. (b) Baxter, S. M.; Wolczanski, P. T., unpublished results.

(30) Anal. Calcd for **5**,  $\text{RhZrCl}_2\text{P}_2\text{O}_7\text{C}_37\text{H}_{36}$ : C, 51.75; H, 4.58; Cl, 8.26. Found: C, 51.49; H, 4.59; Cl, 8.10.  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ )  $\delta$  2.02 (s,  $\text{Cp}^*$ , 15 H), 5.15 (s,  $\text{CH}_2$ , 4 H), 7.45 (br s, Ph, 12 H), 7.68 (m, Ph, 8 H);  $^{31}\text{P}\{\text{H}\}$  NMR  $\delta$  29.41 (d,  $J_{\text{RHP}} = 122$  Hz); IR (Nujol)  $\nu(\text{CO})$  1998  $\text{cm}^{-1}$ .

(31) In  $\text{CDCl}_3$ , trace  $\text{H}_3\text{CCHO}$  was observed; if DCl is formed, H/D exchange with protic sources (probably the walls of the tube) must be facile.

(32) Anal. Calcd for **6**,  $\text{IrRhZrClP}_2\text{O}_7\text{C}_37\text{H}_{36}$ : C, 46.77; H, 4.14; Cl, 3.73; I, 13.36. Found: C, 46.61; H, 4.27; Cl, 3.63; I, 13.21.  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ )  $\delta$  2.12 (s,  $\text{Cp}^*$ , 15 H), 5.33 (d,  $\text{CHH}$ , 2 H,  $^2J = 12$  Hz), 5.45 (dt(v),  $\text{CHH}$ , 2 H,  $^2J = 12$ ,  $J_{\text{PH}} = 1.6$  Hz), 7.29 (m, Ph, 4 H), 7.51 (m, Ph, 12 H), 7.78 (m, Ph, 4 H);  $^{31}\text{P}\{\text{H}\}$  NMR  $\delta$  23.42 (d,  $J_{\text{RHP}} = 127$  Hz); IR (Nujol)  $\nu(\text{CO})$  1994  $\text{cm}^{-1}$ .

(33) Suggs, J. W.; Wovkulich, M. J.; Cox, S. D. *Organometallics* **1985**, *4*, 1101-1107.

(34) HCl + **4**; intermediate A:  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ )  $\delta$  -15.21 (dt, RhH, 1 H,  $J_{\text{RHH}} = 28$ ,  $J_{\text{PH}} = 9$  Hz), 1.05 (t,  $\text{CH}_3$ , 3 H,  $J_{\text{PH}} < 1$  Hz), 2.10 (s,  $\text{Cp}^*$ , 15 H), 5.10 (dt(v),  $\text{CHH}$ , 2 H,  $^2J = 12$ ,  $J_{\text{PH}} = 3$  Hz), 5.60 (d,  $\text{CHH}$ , 2 H,  $^2J = 12$  Hz), 7.45 (m, Ph, 12 H), 7.79 (m, Ph, 4 H), 8.02 (m, Ph, 4 H);  $^{31}\text{P}\{\text{H}\}$  NMR  $\delta$  14.26 (d,  $J_{\text{RHP}} = 98$  Hz); IR (from  $4/4\text{-}^{13}\text{C}_2$ ,  $\text{CD}_2\text{Cl}_2$ )  $\nu(\text{CO})$  2097/2050  $\text{cm}^{-1}$ ;  $\nu(\text{H}_3\text{CC}=\text{O})$  1542/1508  $\text{cm}^{-1}$ .

(35) Tiethof, J. A.; Petersen, J. L.; Meek, D. W. *Inorg. Chem.* **1976**, *15*, 1365-1370.

$\mu_2$ : $\eta^2$ -acetyl (**4**) IR absorption<sup>18,19</sup> falls within the regime of these proposed surface species. The course of oxygenate production may utilize  $H_{ads}$  or proton transfers from Rh/support  $H_2O$  and  $ZrOH^{22}$  functionalities in effecting acyl reductive elimination. Compensation for the loss of the Zr-O link in **4** is ultimately provided by the X of the electrophile. The corresponding oxygenate release in a heterogeneous system could be accompanied by the formation of a new Zr-O bond arising from a similar carbonyl-derived intermediate, a neighboring zirconium oxide/hydroxide,<sup>22</sup> or water.

**Acknowledgment.** Support from the donors of the Petroleum Research Fund, administered by the American Chemical Society, and Cornell University is gratefully acknowledged. We thank the ARCO Foundation for a fellowship (G.S.F.) and the NIH and NSF Instrumentation Programs for support of the Cornell NMR Facility.

### Circular Dichroism of a Subnanosecond State: ( $\Delta$ )-Fe(bpy)<sub>3</sub><sup>2+</sup>\*


Steven J. Milder, Jon S. Gold, and David S. Kliger\*

Department of Chemistry, University of California  
Santa Cruz, California 95064


Received July 21, 1986

We recently presented the first time-resolved circular dichroism (CD) spectrum of a molecular excited state in a report on the CD of the luminescent (MLCT) state of ( $\Delta$ )-Ru(bpy)<sub>3</sub><sup>2+</sup>.<sup>1</sup> The small CD observed in the bpy  $\pi\pi^*$  transition at 370 nm ( $\Delta\epsilon \approx -3$ ) is consistent with a single-ligand localized model for the transferred electron in the excited state. We report here the excited-state absorption and CD spectra of ( $\Delta$ )-Fe(bpy)<sub>3</sub><sup>2+</sup>. Like Ru(bpy)<sub>3</sub><sup>2+</sup>, the ground state of Fe(bpy)<sub>3</sub><sup>2+</sup> is low-spin d<sup>6</sup>. Yet unlike Ru(bpy)<sub>3</sub><sup>2+</sup>, its lowest excited state is a metal-centered d-d state.<sup>2-5</sup> Thus, differences are expected in excited-state absorption and CD spectra of these two molecules. We found a large CD in the ligand  $\pi\pi^*$  transitions of the excited ( $\Delta$ )-Fe(bpy)<sub>3</sub><sup>2+</sup>, consistent with a metal-centered excited state. Since the excited-state lifetime of Fe(bpy)<sub>3</sub><sup>2+</sup> is 800 ps at room temperature in water,<sup>2,6</sup> this is the first report of a CD spectrum of a state that lives for less than a nanosecond.

Creutz et al. previously obtained the absorption spectrum of excited Fe(bpy)<sub>3</sub><sup>2+</sup> in the visible and near-UV regions down to approximately 310 nm and observed no significant charge-transfer transient absorptions.<sup>2</sup> Figure 1 shows the transient absorption spectrum extended to 280 nm.<sup>7</sup> Because of the short lifetime of the excited state, it was not possible to achieve saturation of the excited state with excitation energies of up to 30 mJ/pulse. We thus used the degree of bleaching observed at 488 nm, a wavelength with no significant excited-state absorption,<sup>2</sup> to correct for ground-state bleaching and produce the true excited-state spectrum. This absorption has slightly lower intensity and exhibits a small red shift relative to the ground-state  $\pi\pi^*$  absorption.



**Figure 1.** Ground-state absorption spectrum (—) and excited state absorption spectrum (●) of Fe(bpy)<sub>3</sub><sup>2+</sup> in the  $\pi\pi^*$  region. Inset: The transient difference spectrum extrapolated to complete ground-state depletion (excited-state production).



**Figure 2.** An example of the transient CD signal, representing the average of 256 measurements, seen at 297.5 nm. The signal  $(I_r - I_l)/(I_r + I_l)$  is equal to  $\Delta\epsilon(2.3cl/\delta)$ , where the concentration of ( $\Delta$ )-Fe(bpy)<sub>3</sub><sup>2+</sup> in water is  $4.0 \times 10^{-5}$  M, the path length is 0.2 cm, and the retardation in the probe beam strain plate,  $\delta$ , is 0.018 rad (see ref 8). Excitation was with a 532-nm, 30-mJ, 4-ns pulse. A maximum of 0.5 of the chromophores within the probe beam were excited at any time during the laser pulse.

Clearly, this excited-state absorption can be assigned as primarily ligand  $\pi\pi^*$ .

The excited-state CD of ( $\Delta$ )-Fe(bpy)<sub>3</sub><sup>2+</sup> was obtained by using the laser photolysis apparatus used for transient absorption work with the probe beam optics modified for CD measurements as described in ref 8. Figure 2 shows a typical signal when half of the ( $\Delta$ )-Fe(bpy)<sub>3</sub><sup>2+</sup> within the probe beam is excited at the maximum of the laser pulse. The width of the transient signal is a function of the excited-state lifetime, the excitation pulse width, and the response time of the detection system.

The actual CD signal of the excited state, rather than the change in CD upon excitation, was determined by applying a correction to the change in the signal obtained at the maximum of the laser excitation. The degree of ground-state depletion (excited-state production) at the maximum of the laser pulse was determined, and the change in the CD signal was divided by this value and then added to the ground-state CD to give the excited-state CD. The ground state racemizes quickly,<sup>9,10</sup> with a

(1) Gold, J. S.; Milder, S. J.; Lewis, J. W.; Kliger, D. S. *J. Am. Chem. Soc.* **1985**, *107*, 8285.

(2) Creutz, C.; Chou, M.; Netzel, T. L.; Okumura, M.; Sutin, N. *J. Am. Chem. Soc.* **1980**, *102*, 1309.

(3) Bergkamp, M. A.; Brunshwig, B. S.; Gutlich, P.; Netzel, T. L.; Sutin, N. *Chem. Phys. Lett.* **1981**, *81*, 147.

(4) Decurtins, S.; Felix, F.; Ferguson, J.; Gudel, H. U.; Ludi, A. *J. Am. Chem. Soc.* **1980**, *102*, 4102.

(5) Reeder, K. A.; Dose, E. V.; Wilson, L. J. *Inorg. Chem.*, **1978**, *17*, 1071.

(6) Kirk, A. D.; Hoggard, P. E.; Porter, G. B.; Rockely, M. G.; Windsor, M. W. *Chem. Phys. Lett.* **1976**, *37*, 199.

(7) Excitation was with a short pulse of a Quanta Ray DCR-2 Nd:YAG laser ( $\tau \approx 4$  ns) at 532 nm. Detection of the pulsed xenon probe beam was through a Pacific Precision 0.45-m monochromator by an EMI 9876QB photomultiplier (rise time  $\approx 1.8$  ns). The signal was recorded and analyzed on a Tektronix 7912AD/4041 data acquisition system (500-MHz bandwidth).

(8) Lewis, J. W.; Tilton, R. F.; Einterz, C. M.; Milder, S. J.; Kuntz, I. D.; Kliger, D. S. *J. Phys. Chem.* **1985**, *89*, 289.