

3.5 Hz, 1 H), 6.89 (br d,  $J = 8.8$  Hz, 2 H), 7.3-7.5 (m, 7 H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  55.02, 72.16, 73.10, 75.66, 113.55, 127.24, 127.59, 128.23, 132.41, 137.67, 158.98; IR (KBr) 3500 (m), 3350 (m), 1610 (s), 1515 (s), 1245 (s), 1120 (br s), 1075 (br s), 1025 (br s), 830 (s), 740 (s), 700 (m)  $\text{cm}^{-1}$ ; mass spectrum (70 eV),  $m/z$  (rel intens) 258 (2,  $\text{M}^+$ ), 137 (100), 109 (29), 91 (35), 77 (20); Anal. ( $\text{C}_{16}\text{H}_{18}\text{O}_3$ ) C, H.

**Equilibration of [(Benzyloxy)methyl]lithium (43) and 2-Lithio-1,3-dioxane (7).** A solution of 185 mg of **44**<sup>2b</sup> (0.485 mmol) in 1 mL of THF was cooled to  $-78$  °C, and *n*-butyllithium (0.28 mL of a 1.6 M solution in hexanes, 0.44 mmol) was added dropwise. The resulting clear, pale yellow solution was stirred at  $-78$  °C for 20 min, and a solution of **17** (183 mg, 0.485 mmol) in 1 mL of THF was added via a cannula, followed by rinsing with two 1-mL portions of THF. After the mixture was stirred 30 min further, 84 mg of *p*-anisaldehyde (0.62 mmol, 80  $\mu\text{L}$ ) was added dropwise. The resulting solution was stirred for an additional 30 min, and then the reaction was quenched at  $-78$  °C by addition of 2 mL of methanol. Capillary GC analysis revealed that carbinols **19** and **45** were formed in a ratio of 99.3:0.7, corrected for product response factors. Workup as described for the preparation of **45**, followed by flash chromatography using 45% ethyl acetate/hexanes as eluant, gave 92 mg of **19** (93% yield).

**Relative Rates of Addition of 43 and 7 to *p*-Anisaldehyde.** A solution of 127 mg (0.31 mmol) of **44** and 116 mg (0.31 mmol) of **17** in 3 mL of THF was cooled to  $-78$  °C, and *n*-butyllithium (0.39 mL of a 1.6 M solution in hexanes, 0.62 mmol) was added dropwise. The resulting pale yellow solution was stirred for 30 min, and *p*-anisaldehyde (37  $\mu\text{L}$ , 0.31 mmol) was added all at once. After stirring 30 min further, the reaction was quenched at  $-78$  °C by addition of 2 mL of methanol. Quantitative

capillary GC analysis, using octadecane as internal standard, indicated that carbinols **19** and **45** were formed in a ratio of 1.21:1 (52 and 43% yields, respectively). *p*-Anisyl-*n*-butylcarbinol (**24**) was also formed in 3% yield.

**Acknowledgment.** We thank the National Institutes of Health, the American Heart Association, and the Colorado Heart Association for generous financial support.

**Registry No.** **4**, 118418-15-8; **5**, 118418-16-9; **6**, 118418-17-0; **7**, 118418-18-1; **8**, 118418-19-2; **9**, 25604-67-5; **10**, 19798-66-4; **11**, 118418-20-5; **12**, 4544-19-8; **13**, 81381-75-1; **14**, 79411-59-9; **15**, 79411-58-8; **16**, 118418-21-6; **17**, 118418-22-7; **18**, 118418-23-8; **19**, 63457-96-5; **20**, 63458-00-4; **21**, 118418-24-9; **22**, 118418-25-0; **23**, 118418-26-1; **24**, 19523-03-6; **25**, 118418-27-2; **26**, 118418-28-3; **27**, 626-68-6; **28**, 64181-30-2; **29**, 118418-29-4; **30**, 5465-07-6; **31**, 99423-28-6; **35**, 118418-30-7; **36**, 118418-31-8; **43**, 71316-95-5; **44**, 66222-28-4; **45**, 838-66-4; *p*-anisaldehyde, 123-11-5; 4,4'-di-*tert*-butylbiphenyl, 1625-91-8; ethylene oxide, 75-21-8; trimethylsilyl trifluoromethanesulfonate, 27607-77-8; trimethyl orthoformate, 149-73-5; (phenylthio)trimethylsilane, 4551-15-9; ethylene glycol, 107-21-1; triethyl orthoformate, 122-51-0; 1,3-propanediol, 504-63-2; galvinoxyl, 2370-18-5; isopropylmagnesium chloride, 1068-55-9; tributyltin hydride, 688-73-3; cyclohexanone, 108-94-1; cyclohexanecarboxaldehyde, 2043-61-0; dimethyl sulfate, 77-78-1; 1-bromo-3-phenylpropane, 637-59-2; cyclopentene oxide, 285-67-6; oxetane, 503-30-0; 2-cyclohexen-1-one, 930-68-7; tetrakis[iodo(tri-*n*-butylphosphine)copper(I)], 59245-99-7; chlorotri-*n*-butylstannane, 1461-22-9.

## Redox Glycosidation: A New Strategy for Disaccharide Synthesis<sup>‡</sup>

Anthony G. M. Barrett,\* Barend C. B. Bezuidenhout, Alan F. Gasiiecki, Amy R. Howell, and Mark A. Russell

Contribution from the Department of Chemistry, Northwestern University, Evanston, Illinois 60208. Received November 12, 1987.

Revised Manuscript Received September 14, 1988

**Abstract:** A new procedure for the preparation of disaccharides via redox glycosidation is described. The crucial anomeric C–O bond is established by acylation not alkylation as in traditional Koenigs–Knorr chemistry. Tebbe methylenylation followed by cyclization unravels the protected nonreducing or reducing disaccharides. Thus, for example, 2,3,4,6-tetra-*O*-benzyl-D-glucopyranose (**1**) was coupled with 5-[(*tert*-butyldiphenylsilyloxy]pentanoyl chloride (**5**). The resultant  $\alpha$ -ester **2** was methylenylated with the Tebbe reagent, desilylated, and cyclized with iodine to produce the pyranosylmethyl iodide **4** in good yield. The redox glycosidation protocol was extended to a range of reducing and nonreducing disaccharide systems.

The efficient construction of oligosaccharides in a stereocontrolled manner remains a challenging area for synthetic organic chemistry. Such compounds are of considerable interest in consequence of their diverse, vital roles in many biological processes.<sup>1</sup> Classically, oligosaccharides are assembled with iterative Koenigs–Knorr<sup>2</sup> reactions to sequentially construct the glycosidic bonds. There are now many variations on this theme in which a protected glycosyl halide or related electrophile is condensed with a partially protected second sugar unit.<sup>3</sup> All these existing methods employ an alkylation strategy to elaborate the crucial glycosidic C–O bonds. In oligosaccharide synthesis it is essential that the glycosidation methods used are high yielding. Additionally, the reactions should proceed rapidly and readily control the  $\alpha$  versus  $\beta$  diastereoselectivity at each iteration irrespective of ring substituents. However, in spite of extensive studies on variants of Koenigs–Knorr chemistry spanning nearly 100 years, oligosaccharide synthesis cannot yet be considered simple routine.<sup>4</sup>

Herein we describe model studies on a new procedure for the preparation of disaccharides via redox glycosidation. In this process the two units are linked via an ester bond; subsequent reductive unravelling reveals the disaccharide entity. Thus in the process of crucial anomeric C–O bond is established via acylation,

(1) For examples, see: Kennedy, J. F.; White, C. A. In *Comprehensive Organic Chemistry*; Barton, D. H. R., Ollis, W. D., Eds.; Pergamon Press: Oxford, 1979; Vol. 5, p 755. Schaner, R. *Adv. Carbohydr. Chem. Biochem.* **1982**, *40*, 131.

(2) Igarashi, K. *Adv. Carbohydr. Chem. Biochem.* **1977**, *34*, 243.

(3) For examples of recent developments in glycosidation chemistry, see: Schmidt, R. R. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 212. Nicolaou, K. C.; Ladduwahetty, J. L. R.; Chuchulowski, A. *J. Am. Chem. Soc.* **1986**, *108*, 2466. Nicolaou, K. C.; Dolle, R. E.; Papahatjis, D. P.; Randall, J. L. *J. Am. Chem. Soc.* **1984**, *106*, 4189. Nicolaou, K. C.; Dolle, R. E.; Chuchulowski, A.; Randall, J. L. *J. Chem. Soc., Chem. Commun.* **1984**, 1153.

(4) In contrast to oligosaccharide chemistry, syntheses of oligonucleotides and polypeptides are now highly sophisticated. For examples, see: Letsinger, R. L. *Chemical Synthesis of Oligodeoxyribonucleotides: A Simplified Procedure*. In *Genetic Engineering: Principles and Methods*; Setlow, J. K., Hollaender, A., Eds.; Plenum Press: New York, 1983; Vol. 5, pp 191–209. Merrifield, B. *Science* **1986**, *232*, 341.

<sup>‡</sup>Dedicated to Professor Sir Derek H. R. Barton on the occasion of his 70th birthday.

Scheme I




Table I. Tebbe Methylenylation and Iodoetherification of Glycosyl Esters

entry	ester	(method of prepn, <sup>a</sup> % yield, $\alpha:\beta$ )	vinyl ether	(% yield)	iodo ether	(method of cyclization, <sup>b</sup> % yield, diastereoselectivity)
1	2	(A, 50, 13:1)	3	(69)	4	(A, 72, 1.4:1 <sup>c</sup> )
2	7a	(B, 44, 1:6)	7b	(86)	8	(A, 63, 1:1 <sup>c</sup> )
3	9a	(C, 52, 1:2)	9b	$\beta$ -isomer (63) <sup>d</sup>	10	(B, 79, e)
			9b	$\alpha$ -isomer (13) <sup>d</sup>	11	(B, 80, 1:1 <sup>c</sup> )
4	12a	(D, 65, 1:11)	12b	(67)	13, 14	(C, 65, 8:1)
5	15a	(E, 80, 7.4:1)	15b	(82)	16, 17	(C, 59, $\geq 40:1$ )
6	18a	(D, 71, 1:12)	18b	(72)	19, 20	(D, 69, 5:1)
7	21a	(E, 66, 18:1)	21b	(84)	22, 23	(D, 65, 2:1)
8	24a	(F, 68, e)	24b	(44)	25	(C, 54, 2:1 <sup>c</sup> )
9	26a	(F, 84, e)	26b	(83)	27	(C, 76, 2:1 <sup>c</sup> )
10	28a	(F, 98, e)	28b	(82)	29, 30	(E, 60, 3:1)

<sup>a</sup> Methods for ester preparation from the alcohol by reaction with (A)  $n\text{BuLi}$ , THF,  $-78^\circ\text{C}$ , acyl chloride (ref 8); (B)  $n\text{BuLi}$ , PhH,  $60^\circ\text{C}$ , acyl chloride (ref 8); (C)  $i\text{PrN}=\text{C}=\text{N}i\text{Pr}$ , 4-(dimethylamino)pyridine,  $\text{CH}_2\text{Cl}_2$ , acid (ref 12); (D) NaH (cat.),  $\text{CCl}_3\text{CN}$ ,  $\text{CH}_2\text{Cl}_2$ ,  $25^\circ\text{C}$  then acid,  $\text{CH}_2\text{Cl}_2$  (ref 6); (E)  $n\text{BuLi}$ , THF,  $-78^\circ\text{C}$ , 2-(acylthio)-3-nitropyridine (ref 13); (F) DCC, 4-pyrrolidinopyridine,  $\text{CH}_2\text{Cl}_2$ , acid (ref 12). <sup>b</sup> Methods used for cyclization: (A)  $\text{Bu}_4\text{NF}$ , THF followed by  $\text{I}_2$ ,  $t\text{BuOK}$ , THF,  $0^\circ\text{C}$  to  $25^\circ\text{C}$ ; (B)  $\text{Bu}_4\text{NF}$ , THF followed by  $\text{I}_2$ ,  $t\text{BuOK}$ , THF,  $-78^\circ\text{C}$  to  $-20^\circ\text{C}$ ; (C)  $\text{Bu}_4\text{NF}$ , silica,  $\text{I}_2$ ,  $t\text{BuOK}$ , THF; (D)  $\text{Bu}_4\text{NF}$ , THF followed by  $\text{I}_2$ ,  $t\text{BuOK}$ , THF,  $-78^\circ\text{C}$  to  $-40^\circ\text{C}$ ; (E)  $\text{Bu}_4\text{NF}$ , THF followed by  $\text{I}_2$ ,  $t\text{BuOK}$ , THF,  $0^\circ\text{C}$ . <sup>c</sup> Structure of major diastereoisomer not assigned. <sup>d</sup> The  $\alpha$ - and  $\beta$ -isomers were separated at the vinyl ether stage following desilylation using  $\text{Bu}_4\text{NF}$  in THF. <sup>e</sup> Ratio not determined.

not alkylation. Such a procedure benefits from the fact that acylation reactions are rapid and proceed with moderate to excellent  $\alpha$  to  $\beta$  diastereoselectivities.<sup>5,6</sup>

2,3,4,6-Tetra-*O*-benzyl-D-glucopyranose (1) was esterified with 5<sup>7</sup>, by using the lithium alkoxide method of Pfeffer<sup>8</sup> to stereoselectively (13:1) produce the  $\alpha$ -glucosyl ester 2 (50%). Subsequent methylenylation using the Tebbe reagent 6<sup>9,10</sup> gave the

anomeric vinyl ether 3 (69%). Desilylation and iodoetherification gave the model nonreducing disaccharide 4 (72%) as a mixture of diastereoisomers (1:1.4) (Scheme I). We were delighted that the delicate anomeric C–O bond of 3 was not cleaved at the intermediate iodonium ion stage. The method was extended to a range of disaccharide systems (Table I, Chart I). Four distinct procedures were used for the esterification step:<sup>11</sup> (i) carbodiimide coupling<sup>12</sup> (entries 3, 8–10), (ii)  $\beta$ -selective esterification using trichloroimidate activation<sup>6</sup> (entries 4, 6), (iii)  $\alpha$ -selective esterification by reaction of the anomeric alkoxide with the acid chloride<sup>8</sup> (entry 1) or 2-(acylthio)-3-nitropyridine<sup>13,14</sup> (entries 5, 7), and (iv)  $\beta$ -selective esterification by reaction of the anomeric alkoxide with the acid chloride at  $60^\circ\text{C}$ <sup>8</sup> (entry 2). All the esters were smoothly methylenylated, desilylated, and iodocyclized to produce the corresponding protected nonreducing (entries 1–7)

(5) Schmidt, O. T.; Herok, J. *Justus Liebigs Ann. Chem.* **1954**, 587, 63. Pederson, C.; Fletcher, H. G. *J. Am. Chem. Soc.* **1960**, 82, 3215. Fletcher, H. G. *Methods Carbohydr. Chem.* **1963**, 2, 231. Kornhauser, A.; Keglevic, D. *Carbohydr. Res.* **1969**, 11, 407. Bugiaesi, R.; Shen, T. Y. *Ibid.* **1971**, 19, 179. Keglevic, D.; Valetkovic, S.; Roglic, G.; Goles, D.; Plavsic, F. *Ibid.* **1973**, 29, 25. Ogawa, T.; Nozaki, M.; Matsui, M. *Ibid.* **1978**, 60, C7. Shoda, S.; Mukaiyama, T. *Chem. Lett.* **1982**, 861. Nicolaou, K. C.; Chucholowski, A.; Dolle, R. E.; Randall, J. L. *J. Chem. Soc., Chem. Commun.* **1984**, 1155. Smith, A. B., III; Hale, K. J.; Rivero, R. A. *Tetrahedron Lett.* **1986**, 27, 5813. Kochetkov, N. K.; Klimov, E. M.; Pogosjan, S. A.; Derivitskaya, V. A. *Izv. Akad. Nauk SSSR, Ser. Khim.* **1972**, 7, 1657.

(6) Schmidt, R. R.; Michel, J. *Angew. Chem., Int. Ed. Engl.* **1980**, 19, 731. (7) Barrett, A. G. M.; Dhanak, D. *Tetrahedron Lett.* **1987**, 28, 3327. (8) Pfeffer, P. E.; Rothman, E. S.; Moore, G. G. *J. Org. Chem.* **1976**, 41, 2926.

(9) Tebbe, F. N.; Parshall, G. W.; Reddy, G. S. *J. Am. Chem. Soc.* **1977**, 100, 3611. Pine, S. H.; Zahler, R.; Evans, D. A.; Grubbs, R. H. *Ibid.* **1980**, 102, 3270. See also: Brown-Wensley, K. A.; Buchwald, S. L.; Cannizzo, L.; Clawson, L.; Ho, S.; Meinhardt, D.; Stille, J. R.; Straus, D.; Grubbs, R. H. *Pure Appl. Chem.* **1983**, 55, 1733. For an in situ preparation of this reagent: Cannizzo, L. F.; Grubbs, R. H. *J. Org. Chem.* **1985**, 50, 2386.

(10) For examples of Tebbe methylenylation of carbohydrate systems, see: Wilcox, C. S.; Long, G. W.; Suh, H. *Tetrahedron Lett.* **1984**, 25, 395. RajanBabu, T. V.; Reddy, G. S. *J. Org. Chem.* **1986**, 51, 5458.


(11) Carboxylic acids were prepared from the corresponding aldonic acid lactones, see ref 7.

(12) Hassner, A.; Alexanian, V. *Tetrahedron Lett.* **1978**, 19, 4475. Neiser, B.; Steglich, W. *Angew. Chem., Int. Ed. Engl.* **1978**, 17, 522. Zeigler, F. E.; Burger, G. D. *Synth. Commun.* **1979**, 9, 539.

(13) Barrett, A. G. M.; Bezuidenhout, B. C. B., unpublished observations.

(14) For the preparation of 3,3'-dinitro-2,2'-dipyridyl disulfide and synthesis of 2-pyridinethiol esters, see: Matsueda, R.; Walter, R. *Int. J. Pept. Protein Res.* **1980**, 16, 392. Mukaiyama, T.; Matsueda, R.; Suzuki, M. *Tetrahedron Lett.* **1970**, 11, 1901. Mukaiyama, T.; Goto, K.; Matsueda, R.; Ueki, M. *Ibid.* **1970**, 11, 5293. Corey, E. J.; Brunelle, D. J. *Ibid.* **1976**, 17, 3409. Corey, E. J.; Nicolaou, K. C.; Melvin, L. S. *J. Am. Chem. Soc.* **1975**, 97, 653 and 654.

Chart I



and reducing (entries 8–10) disaccharides.<sup>15</sup> It is clear from these results that the protocol may be used to introduce either (iodomethyl)pyranosyl or furanosyl groups. It should be noted that the cyclizations which produced pyranosyl methyl iodides were slower than those producing tetrahydrofuran systems.

The stereochemistry of reaction requires comment. The stereochemistries of these iodoetherification reactions are clearly kinetically controlled.<sup>16</sup> Cyclizations of the simple unsubstituted vinyl ethers proceeded without any appreciable stereochemical control by D-glucose (entries 1–3) or D-galactose (entries 8, 9). However, the cyclizations of vinyl ethers bearing side-chain substituents proceeded with modest to excellent diastereoselectivities (2:1 to >40:1). It is particularly interesting to note the variation in the diastereoselectivity in entries 4–7. Presumably stereochemical control by the isopropylidene and the methoxymethyl substituents are mismatched.<sup>17</sup> The redox glycosidation of 2,3:5,6-di-O-isopropylidene- $\alpha$ -D-galactofuranose (entry 10) is a favorable auspice for application of this approach in sialic acid chemistry.<sup>18</sup>

In conclusion, it is apparent that the acylation–methylenylation–cyclization protocol provides a novel method for the elaboration of the glycosidic bond. Further studies on this protocol and other redox glycosidation technologies are under investigation.

## Experimental Section

Representative procedures are given in this section. Full experimental details are provided in the supplementary material.

**2,3,4,6-Tetra-O-methyl- $\alpha$ -D-glucopyranosyl 2,3-O-Isopropylidene-5-O-methyl-4-O-(tert-butylidiphenylsilyl)-D-ribonate (21a).** 3,3'-Dinitro-2,2'-dipyridyl disulfide<sup>14</sup> (30 mg) and Ph<sub>3</sub>P (26 mg) were added to a solution of 4-O-(tert-butylidiphenylsilyl)-2,3-O-isopropylidene-5-O-methyl-D-ribonic acid<sup>11</sup> (30 mg) in dry CH<sub>2</sub>Cl<sub>2</sub> (1 mL) under N<sub>2</sub> at room temperature, and the slurry was stirred for 4 h. The solids were filtered off, and the resultant thioester was purified by preparative TLC on silica (hexanes/EtOAc 3:1, R<sub>f</sub> 0.41). *n*-Butyllithium (0.17 M, 0.4 mL) was added to a solution of 2,3,4,6-tetra-O-methyl-D-glucose (19 mg), in dry THF (0.5 mL) at –78 °C under N<sub>2</sub>. After 10 min, a solution of the thioester in dry THF (0.5 mL) was added and stirring continued at –78 °C for 32 h. The mixture was quenched by adding it to a saturated aqueous NH<sub>4</sub>Cl solution (10 mL), and the ester was extracted with Et<sub>2</sub>O. Drying (Na<sub>2</sub>SO<sub>4</sub>) of the extract followed by evaporation and chromatography on silica (eluant hexanes/EtOAc 3:1) gave the ester (29 mg, 66%) as a 18:1 (21a:18a) mixture of anomers. The  $\alpha$ -anomer 21a was obtained as a colorless syrup by chromatography on silica (eluant hexanes/Et<sub>2</sub>O 3:2): [ $\alpha$ ]<sub>D</sub> +32.8° (*c* = 1.2, CHCl<sub>3</sub>); TLC R<sub>f</sub> 0.2 (silica, hexanes/Et<sub>2</sub>O 3:2); IR (neat) 1771, 1757, 1433, 1388, 1150, 1110, 1000, 708 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.78–7.68, 7.44–7.32 (2 m, 10 H), 6.29 (d, 1 H, *J* = 3.6 Hz), 4.42 (d, 1 H, *J* = 6.8 Hz), 4.38 (dd, 1 H, *J* = 4.2, 6.8 Hz), 4.22–4.16 (m, 1 H), 3.55–3.20 (m, 8 H), 3.59 (s, 3 H), 3.53 (s, 3 H), 3.37 (s, 3 H), 3.27 (s, 3 H), 3.03 (s, 3 H), 1.59 (s, 3 H), 1.32 (s, 3 H), 1.05 (s, 9 H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  168.6, 136.1, 135.9, 134.2, 133.7, 129.6, 129.5, 127.4, 110.4, 90.6, 83.0, 80.9, 79.8, 78.1, 75.5, 73.1, 73.0, 70.4, 70.1, 60.8, 60.5, 59.1, 58.9, 58.2, 27.0, 26.6, 25.3, 19.3; mass spectrum (EI), *m/e* 661 (M<sup>+</sup> – Me), 441, 385, 255, 219, 187, 175, 167, 155, 145, 127, 111, 101, 59. Anal. Calcd for C<sub>35</sub>H<sub>52</sub>O<sub>11</sub>Si: C, 62.11; H, 7.74. Found: C, 62.08; H, 7.81.

**2,3,4,6-Tetra-O-methyl- $\beta$ -D-glucopyranosyl 2,3-O-Isopropylidene-5-O-methyl-4-O-(tert-butylidiphenylsilyl)-D-ribonate (18a).** NaH (0.5 mg) was added to a stirred solution of 2,3,4,6-tetra-O-methyl-D-glucopyranose (50 mg) in dry CH<sub>2</sub>Cl<sub>2</sub> (1 mL) at –40 °C under N<sub>2</sub>. After 10 min CCl<sub>3</sub>CN (0.5 mL) was added, and the stirring was continued for 1 h, before the solvent and excess CCl<sub>3</sub>CN were removed under reduced pressure. Chromatography on silica (eluant hexanes/EtOAc 3:1) gave

the corresponding imidate (69 mg). The imidate was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (1 mL) and 4-O-(tert-butylidiphenylsilyl)-2,3-O-isopropylidene-5-O-methyl-D-ribonic acid (83 mg) was added. After stirring of the mixture at room temperature under N<sub>2</sub> for 1 h, CH<sub>2</sub>Cl<sub>2</sub> (30 mL) and ice were added, and the mixture was washed with 2 M HCl (4 × 20 mL) and saturated aqueous NaHCO<sub>3</sub> (1 × 20 mL). Drying (MgSO<sub>4</sub>), evaporation at reduced pressure, and chromatography on silica (eluant hexanes/Et<sub>2</sub>O 3:2) of the CH<sub>2</sub>Cl<sub>2</sub> extract gave 21a and 18a (87 mg, 71%) as a 1:12 mixture of anomers. Repurification using the same system yielded the pure  $\beta$ -isomer 18a as a colorless syrup: [ $\alpha$ ]<sub>D</sub> –29.5° (*c* = 2.0, CHCl<sub>3</sub>); TLC R<sub>f</sub> 0.3 (silica, hexanes/Et<sub>2</sub>O 3:2); IR (neat) 1763, 1590, 1230, 710 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.76–7.68, 7.45–7.30 (2 m, 10 H), 5.39 (d, 1 H, *J* = 8.0 Hz), 4.47 (d, 1 H, *J* = 6.8 Hz), 4.41 (dd, 1 H, *J* = 4.2, 6.8 Hz), 4.32–4.20 (m, 1 H), 3.52–3.05 (m, 8 H), 3.63 (s, 3 H), 3.52 (s, 3 H), 3.41 (s, 3 H), 3.31 (s, 3 H), 2.98 (s, 3 H), 1.57 (s, 3 H), 1.32 (s, 3 H), 1.04 (s, 9 H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  168.8, 136.2, 135.9, 134.3, 133.8, 129.5, 129.4, 127.4, 127.3, 110.5, 94.4, 86.8, 82.1, 79.7, 78.6, 75.4, 73.3, 72.9, 70.3, 70.0, 60.7, 60.4, 60.3, 59.2, 58.1, 27.0, 26.5, 25.4, 19.4; mass spectrum (EI), *m/e* 661 (M<sup>+</sup> – Me), 441, 417, 385, 311, 255, 219, 187, 155, 145, 127, 111, 101. Anal. Calcd for C<sub>35</sub>H<sub>52</sub>O<sub>11</sub>Si: C, 62.11; H, 7.74. Found: C, 62.30; H, 7.78.

**2,2-Dimethyl-5(R)-[2-methoxy-1(R)-[(tert-butylidiphenylsilyl)oxy]ethyl]-4(R)-[1-[(2,3,4,6-tetra-O-methyl- $\beta$ -D-glucopyranosyl)oxy]ethenyl]-1,3-dioxolane (18b).** Tebbe reagent 6 in PhMe (0.50 M, 1 mL) was added dropwise to 18a in dry PhMe (0.5 mL), dry THF (0.1 mL), and dry pyridine (0.1 mL) at –40 °C. The reaction mixture was stirred at that temperature for 30 min followed by 2 h at 0 °C and 1 h at room temperature. The mixture was cooled to –40 °C and quenched with a 15% NaOH solution (0.5 mL). The reaction mixture was diluted with Et<sub>2</sub>O (10 mL) and filtered through Celite. The filtrate was evaporated at reduced pressure, and the crude product was purified by chromatography on silica (eluant hexanes/Et<sub>2</sub>O 3:2). Alkene 18b (40 mg, 72%) was recovered as a cloudy syrup: [ $\alpha$ ]<sub>D</sub> +2.4° (*c* = 1.1, CHCl<sub>3</sub>); TLC R<sub>f</sub> 0.3 (silica, hexanes/Et<sub>2</sub>O 3:2); IR (neat) 1650, 1468, 1442, 1387, 1250, 1100, 1000, 705 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.80–7.70, 7.45–7.30 (2 m, 10 H), 4.56 (d, 1 H, *J* = 7.6 Hz), 4.46 (d, 1 H, *J* = 7.6 Hz), 4.43 (d, 1 H, *J* = 2.0 Hz), 4.40–4.35 (m, 2 H), 4.27–4.22 (m, 1 H), 3.62 (s, 3 H), 3.53 (s, 3 H), 3.53–3.00 (m, 8 H), 3.38 (s, 3 H), 3.30 (s, 3 H), 2.85 (s, 3 H), 1.49 (s, 3 H), 1.31 (s, 3 H), 1.01 (s, 9 H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  157.0, 135.9, 134.9, 134.8, 129.2, 127.3, 127.1, 108.5, 100.3, 91.2, 86.8, 83.1, 80.5, 79.2, 74.9, 74.0, 71.8, 71.2, 60.7, 60.4, 60.3, 59.3, 57.8, 27.0, 26.1, 24.5, 19.4; mass spectrum (EI), *m/e* 659 (M<sup>+</sup> – Me), 527, 341, 219, 187, 155, 127, 111. Anal. Calcd for C<sub>36</sub>H<sub>54</sub>O<sub>10</sub>Si: C, 64.07; H, 8.06. Found: C, 64.08; H, 8.16.

**2,2-Dimethyl-5(R)-[2-methoxy-1(R)-[(tert-butylidiphenylsilyl)oxy]ethyl]-4(R)-[1-[(2,3,4,6-tetra-O-methyl- $\alpha$ -D-glucopyranosyl)oxy]ethenyl]-1,3-dioxolane (21b).** Reaction of 21a (50 mg) with the Tebbe reagent 6, as described for 18a, gave 21b as a colorless solid (62 mg, 84%). Recrystallization from hexanes gave colorless cubes: mp 103 °C; [ $\alpha$ ]<sub>D</sub> +40.5° (*c* = 1.1, CHCl<sub>3</sub>); IR (KBr) 1650, 1480, 1440, 1390, 1304, 1230, 1170, 1115, 1102, 1050, 1000, 890, 856, 835 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.76–7.65, 7.42–7.29 (2 m, 10 H), 4.93 (d, 1 H, *J* = 4.0 Hz), 4.47 (d, 1 H, *J* = 6.8 Hz), 4.44 (d, 1 H, *J* = 2.0 Hz), 4.34 (dd, 1 H, *J* = 6.6, 6.8 Hz), 4.18 (d, 1 H, *J* = 2.0 Hz), 4.14–4.10 (m, 1 H), 3.52–3.17 (m, 8 H), 3.52 (s, 3 H), 3.48 (s, 3 H), 3.37 (s, 3 H), 3.05 (s, 3 H), 3.01 (s, 3 H), 1.50 (s, 3 H), 1.34 (s, 3 H), 1.01 (s, 9 H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  156.8, 136.2, 135.9, 134.8, 134.4, 129.4, 129.1, 127.3, 127.1, 108.6, 94.5, 92.0, 83.4, 81.2, 79.5, 78.9, 78.7, 74.1, 70.6, 70.5, 60.6, 60.2, 59.1, 58.6, 58.0, 27.2, 26.8, 25.0, 19.4; mass spectrum (EI), *m/e* 659 (M<sup>+</sup> – Me), 527, 367, 341, 219, 187, 155, 145, 127, 111, 101. Anal. Calcd for C<sub>36</sub>H<sub>54</sub>O<sub>10</sub>Si: C, 64.07; H, 8.06. Found: C, 63.96; H, 8.21.

**2,3,4,6-Tetra-O-methyl- $\beta$ -D-glucopyranosyl 1-Deoxy-3-*epi*-1-iodo-3,4-O-isopropylidene- $\beta$ -D-fructofuranoside (19) and 2,3,4,6-Tetra-O-methyl- $\beta$ -D-glucopyranosyl 1-Deoxy-3-*epi*-1-iodo-3,4-O-isopropylidene- $\alpha$ -D-fructofuranoside (20).** The vinyl ether 18b (50 mg) was dissolved in tetrabutylammonium fluoride in THF (1.0 M; 1.5 mL), and the mixture was stirred at room temperature under N<sub>2</sub> for 24 h. The solvent was removed at reduced pressure and the product alcohol was purified by chromatography on silica (eluant Et<sub>2</sub>O, R<sub>f</sub> 0.2). To a stirred solution of the alcohol in dry THF (2 mL) under N<sub>2</sub> at room temperature was added <sup>t</sup>BuOK (35 mg), and the mixture stirred for 5 min before it was cooled to –78 °C and I<sub>2</sub> (55 mg) was added. After stirring of the mixture for 4 h, the temperature was raised to –40 °C and the stirring was continued for another 18 h. The reaction mixture was quenched with saturated aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution (3 mL) and the mixture was extracted with Et<sub>2</sub>O (3 × 2 mL). After drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation, chromatography on silica (eluant hexanes/Et<sub>2</sub>O 2:3) gave 19 and 20 (26 mg, 69%) as a 4.8:1 mixture of isomers: TLC R<sub>f</sub> 0.4 (silica, hexanes/Et<sub>2</sub>O 2:3). Anal. Calcd for C<sub>20</sub>H<sub>32</sub>I<sub>2</sub>O<sub>10</sub>: C, 42.71; H, 6.27. Found: C,

(15) All new compounds were fully authenticated by spectral data and microanalyses or high-resolution mass ion measurement. The stereochemistries of the iodomethyl substituents were confirmed by <sup>1</sup>H and <sup>13</sup>C spectral assignments and by extensive NOE experiments (see supplementary material).

(16) For examples of iodoetherification reactions, see: Reitz, A. B.; Nortey, S. O.; Maryanoff, B. E.; Liotta, D.; Monahan, R. III *J. Org. Chem.* **1987**, *52*, 4191. Tamamu, Y.; Hojo, M.; Kawanura, S.; Sawada, S.; Yoshida, Z.; *Ibid.*, **1987**, *52*, 4062. Rychnovsky, S. D.; Bartlett, P. A. *J. Am. Chem. Soc.* **1981**, *103*, 3963.

(17) For a discussion of matched and mismatched stereochemical biases, see Masamune, S.; Choy, W. *Aldrichimica Acta* **1982**, *15*, 47.

(18) Vliegenghart, F. G.; Kamerling, J. P. In *Sialic Acids: Chemistry, Metabolism and Function*; Schauer, R., Ed.; Springer-Verlag: New York, 1982; p 59–76.

42.50; H, 6.20. The isomers of  $R_f$  0.4 (15 mg) and  $R_f$  0.3 (4 mg) were separated by chromatography on silica (eluant hexanes/Et<sub>2</sub>O 1:1). The major isomer **19** ( $R_f$  0.4) was obtained as a colorless syrup:  $[\alpha]_D^{25} -94^\circ$  ( $c = 1.15$ , CHCl<sub>3</sub>); IR (neat) 1460, 1413, 1390, 1380, 1255, 1216, 1160, 1100 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  4.70 (d, 1 H,  $J = 7.6$  Hz), 4.68 (d, 1 H,  $J = 5.6$  Hz), 4.57 (dd, 1 H,  $J = 2.8, 5.6$  Hz), 4.43-4.38 (m, 1 H), 3.76 (d, 1 H,  $J = 10.0$  Hz), 3.67-3.41 (m, 5 H), 3.62 (s, 3 H), 3.53 (s, 3 H), 3.52 (s, 3 H), 3.42 (s, 3 H), 3.41 (s, 3 H), 3.29-3.04 (m, 4 H), 1.53 (s, 3 H), 1.36 (s, 3 H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  113.3, 109.4, 95.4, 87.4, 86.9, 85.8, 83.5, 82.1, 79.4, 74.7, 72.7, 71.5, 60.8, 60.5, 60.4, 59.6, 58.9, 26.9, 25.6, 5.8; mass spectrum (EI),  $m/e$  547 ( $M^+ - Me$ ), 403, 371, 328, 327, 235, 175, 157, 147, 115, 101. The minor isomer **20** ( $R_f$  0.3) was obtained as a colorless syrup:  $[\alpha]_D^{25} +37^\circ$  ( $c = 0.90$ , CHCl<sub>3</sub>); IR (CCl<sub>4</sub>) 1728, 1691, 1676, 1650, 1555, 1377, 1085 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  4.85 (d, 1 H,  $J = 7.2$  Hz), 4.68-4.65 (m, 1 H), 4.62 (dd, 1 H,  $J = 4.0, 7.2$  Hz), 4.58 (d, 1 H,  $J = 8.0$  Hz), 3.63 (s, 3 H), 3.58 (s, 3 H), 3.52 (s, 3 H), 3.40 (s, 3 H), 3.37 (s, 3 H), 3.63-2.95 (m, 10 H), 1.61 (s, 3 H), 1.36 (s, 3 H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  116.5, 103.1, 96.3, 87.0, 84.7, 83.5, 82.4, 81.4, 79.6, 74.8, 72.2, 71.5, 61.2, 60.8, 60.3, 59.5, 59.4, 26.4, 26.2, 6.6; mass spectrum (EI),  $m/e$  547 ( $M^+ - Me$ ), 371, 328, 327, 285, 235, 207, 187, 175, 147.

**2,3,4,6-Tetra-O-methyl- $\alpha$ -D-glucopyranosyl 1-Deoxy-3-*epi*-1-iodo-3,4-isopropylidene- $\alpha$ -D-fructofuranoside (22) and 2,3,4,6-Tetra-O-methyl- $\alpha$ -D-glucopyranosyl 1-Deoxy-3-*epi*-1-iodo-3,4-isopropylidene- $\beta$ -D-fructofuranoside (23).** Treatment of the  $\alpha$ -vinyl ether **21b** (50 mg) with Bu<sub>4</sub>NF in THF (1.0 M; 1.5 mL) followed by <sup>t</sup>BuOK (35 mg) and I<sub>2</sub> (55 mg), as described above for **18b**, gave the disaccharides **22** and **23** (24 mg, 65%) as a 2:1 mixture, after chromatography on silica (eluant hexanes/Et<sub>2</sub>O 1:4):  $[\alpha]_D^{25} +86^\circ$  ( $c = 1.15$ , CHCl<sub>3</sub>); TLC  $R_f$  0.5 (silica, hexanes/Et<sub>2</sub>O 1:4); IR (neat) 1460, 1385, 1260, 1220, 1200, 1170, 1110 cm<sup>-1</sup>; mass spectrum (EI),  $m/e$  547 ( $M^+ - Me$ ), 371, 343, 327, 285, 235, 207, 187, 175, 157, 147, 115, 101. Anal. Calcd for C<sub>20</sub>H<sub>35</sub>O<sub>10</sub>: C,

42.71; H, 6.27. Found: C, 42.74; H, 6.17. The two anomers respectively showed the following NMR data.  $\alpha$ -Anomer **22**: <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>)  $\delta$  5.45 (d, 1 H,  $J = 3.6$  Hz), 4.69 (d, 1 H,  $J = 8.0$  Hz), 4.62-4.58 (m, 1 H), 4.49 (dd, 1 H,  $J = 4.8, 8.0$  Hz), 4.27-4.22 (m, 1 H), 3.90 (dd, 1 H,  $J = 9.4$  Hz,  $J = 9.4$  Hz), 3.70-3.44 (m, 2 H), 3.65 (s, 3 H), 3.56 (s, 3 H), 3.54 (d, 1 H,  $J = 11.2$  Hz), 3.40 (d, 1 H,  $J = 11.2$  Hz), 3.37-3.08 (m, 4 H), 3.19 (s, 3 H), 3.15 (s, 3 H), 3.04 (s, 3 H), 1.74 (s, 3 H), 1.25 (s, 3 H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  116.8, 103.7, 90.1, 85.8, 82.9, 81.9, 81.7, 81.1, 78.8, 72.3, 71.3, 70.7, 60.7, 60.4, 59.6, 59.2, 58.6, 26.7, 26.4, 8.2.  $\beta$ -Anomer **23**: <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>)  $\delta$  5.46 (d, 1 H,  $J = 3.6$  Hz), 4.98 (d, 1 H,  $J = 5.6$  Hz), 4.93 (dd, 1 H,  $J = 5.0, 5.6$  Hz), 4.36-4.31 (m, 1 H), 4.23-4.18 (m, 1 H), 3.84 (dd, 1 H,  $J = 9.2$  Hz,  $J = 9.2$  Hz), 3.70-3.08 (m, 8 H), 3.63 (s, 3 H), 3.49 (s, 3 H), 3.25 (s, 3 H), 3.16 (s, 3 H), 3.01 (s, 3 H), 1.35 (s, 3 H), 1.24 (s, 3 H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  113.6, 108.2, 91.1, 84.6, 83.6, 83.2, 81.7, 81.5, 79.2, 72.6, 71.1, 60.8, 59.2, 58.7, 27.0, 25.4, 5.9.

**Acknowledgment.** We thank the National Institutes of Health and G. D. Searle and Co. for support of our research; G. D. Searle and Co. for granting study leave to Alan F. Gasielki; FRD CSIR, Pretoria for financial support to Barend C. B. Bezuidenhout; the Midwest Center for Mass Spectrometry, an NSF Regional Instrument Facility (Grant CHE-8211164), for obtaining mass spectral data; and the National Institutes of Health (Grant RR-02314) for the purchase of a 400-MHz NMR spectrometer used in these studies.

**Supplementary Material Available:** Full experimental details describing the preparation and authentication of all new compounds (40 pages). Ordering information is given on any current masthead page.

## Asymmetric Synthesis with $\alpha,\beta$ -Bis[(methoxymethyl)oxy] Ketones. Enantioselective Total Synthesis of Natural (+)-Indolizidine 195B (Bicyclic Gephyrotoxin 195B) and (-)-Pinidine and Their Enantiomers from a Common Chiral Synthone

Naoki Yamazaki and Chihiro Kibayashi\*

Contribution from the Tokyo College of Pharmacy, Horinouchi, Hachioji, Tokyo 192-03, Japan. Received June 13, 1988

**Abstract:** The first enantioselective total synthesis of naturally occurring (+)-indolizidine 195B (bicyclic gephyrotoxin 195B) and (-)-pinidine and their enantiomers has been achieved starting from 4-*O*-benzyl-2,3-*O*-bis(methoxymethyl)-L-threitol as a single and common chiral synthon, readily available from L-tartaric acid. This synthesis establishes the absolute stereochemistry of (+)-indolizidine 195B as 3*S*,5*S*,9*S*. The strategy for the synthesis of the alkaloids in both enantiomeric forms is based on the process involving a combination of the creation of new stereogenic centers by 1,2-asymmetric induction and destruction of the original chirality inducing group, i.e., *threo*-bis(methoxymethyl) ether. Actually, these processes consist of highly diastereoselective hydride addition to  $\alpha,\beta$ -bis(methoxymethyl)oxy ketones via chelation or nonchelation control by using a variety of borohydride reagents and stereospecific transformation of *threo*-vicinal diols into both *E* and *Z* olefins.

Stereocontrolled addition of hydride from metal hydride reagents to acyclic ketones has been widely used for the preparation of optically active acyclic secondary alcohols. One of the most widely studied processes for this is the enantioselective hydride addition to prochiral ketones by metal hydride reagents modified with chiral ligands (reagent control).<sup>1</sup> Alternative efforts have been focused on the diastereoselective hydride addition of achiral metal hydride reagents (1,2- or 1,3-asymmetric induction) to ketone substrates having a stereogenic center or linked with a chiral

auxiliary (substrate control).<sup>2</sup> Often there is a need for the preparation of molecules in both enantiomeric forms for studies, for example, on their physical or biological properties. In obtaining both enantiomers of a molecule by the methods for asymmetric induction mentioned above, both enantiomers of chiral reagents, chiral building blocks, or chiral auxiliaries are required. In many cases, however, one of the enantiomers of the chiral agents is not readily accessible from commercial sources or may have a high cost (for example, D- vs. L-sugars and L- vs. D-amino acids). To

(1) *Asymmetric Synthesis*; Morrison, J. D., Ed.; Academic Press: New York, 1983; Vol. 2A.

(2) (a) Eliel, E. L., in ref 1, Chapter 5. (b) Ohishi, T.; Nakata, T. *Acc. Chem. Res.* 1984, 17, 388.