

The α -glucoside **7** is a stable compound that decomposes only very slowly in water solution with a half-life of 680 h. Incubation of yeast α -glucosidase¹⁴ with **7** resulted in a time-dependent loss of enzyme activity that followed pseudo-first-order kinetics (Figure 1). A Kitz and Wilson¹⁵ replot of the data indicated that saturation was attained. The K_I value for **7** is 0.7 ± 0.1 mM, and the k_{inact} value is 0.21 ± 0.02 min⁻¹. Phenyl α -D-glucopyranoside, a substrate of the enzyme, as well as α -D-glucose and tris(hydroxymethyl)aminomethane ("Tris"), two competitive inhibitors, protected the enzyme from inactivation. These results demonstrate that the inactivation takes place in the active site. Furthermore, addition of dithiothreitol (5 mM) in the preincubation medium had no effect on the rate of inactivation, indicating that the species responsible for inactivation was not released from the active site.¹⁶ Incubation with 5 mM **7** for 7 min at 37 °C resulted in 80% inactivation of the enzyme. Prolonged dialysis of this inactivated enzyme for 24 h at 4 °C did not regenerate any enzyme activity,

(13) **7**: ¹H NMR (CD₃OD, 360 MHz, TMS as reference) large d (6.6 ppm, CHFCl, ²J_{H-F} = 48 Hz); d (5.85 ppm, anomeric H, J_{H-H} = 3.6 Hz); m (3.6–3.95 ppm, m, 6 H), s (5.1 ppm, 4 H, OH). ¹⁹F NMR (CD₃OD, 338 MHz, CF₃CO₂H as reference) -11 ppm (m, CF₂, 2 AB parts of ABX; ²J_{FF} = 144 Hz, ³J_{FF} = 12.6 Hz); -78.5 ppm (2 dt, CHF₂). MS (CI/NH₃) 314 (MNH₄⁺), 212, 180. Elemental analysis found for (C₈H₁₂O₆F₃Cl): C: 33.00; H: 4.24. Calcd: C, 32.39; H, 4.08.

(14) α -Glucosidase, type III, from yeast (Sigma). The enzyme activity was measured at 37 °C in 0.2 M sodium phosphate buffer (pH 6.9), using *p*-nitrophenyl α -D-glucopyranoside (PNPG) as the substrate, according to Halvorson, H. *Methods Enzymol.* **1966**, *8*, 559.

(15) Kitz, R.; Wilson, I. B. *J. Biol. Chem.* **1962**, *237*, 3245.

(16) Rando, R. R. *Biochem. Pharmacol.* **1974**, *23*, 2328.

suggesting the formation of a covalent linkage of the inhibitor to the enzyme-active site (Scheme II, path A; R = CHFCl).

Contrary to what was observed with yeast α -glucosidase, **7** was found to be a substrate of the sucrase-isomaltase complex purified from rat small intestine.¹⁷ The enzyme-catalyzed hydrolysis of **7** resulted in the liberation of two molecules of HF for one molecule of glucose^{18,19} according to Scheme II, path B. One possible reason that would explain why the mammalian glucosidases, contrary to the yeast enzyme, are not inactivated by **7** might lie in a difference of nucleophilic residues involved in the respective active site of these enzymes.²⁰

In conclusion, we have shown for the first time that a 1,1-di-fluoroalkylglucoside is an enzyme-activated irreversible inhibitor of yeast α -glucosidase. All the biochemical data reported in this paper are in agreement with a novel process of enzyme-activated inhibition due to the inactivatory property of the leaving group released during the glucosidase-catalyzed hydrolysis.

Work is in progress in our laboratories to extend this approach to the inhibition of other hydrolytic enzymes of therapeutic interest.

(17) Danzin, C.; Ehrhard, A. *Arch. Biochem. Biophys.* **1987**, *257*, 472.

(18) Liberation of fluoride ions was measured with an Orion 96-09 fluoride specific electrode connected to a Mettler semiautomatic titration system (C. Gaget, unpublished results). Glucose formation was measured according to the method of Dahlqvist (ref 19).

(19) Dahlqvist, A. *Anal. Biochem.* **1964**, *7*, 18.

(20) For instance, an essential thiol group has been found in the yeast α -glucosidase by Halvorson, 1966 (see ref 14), but not in the mammalian sucrase-isomaltase complex (ref 21).

(21) Kolinska, J.; Semenza, G. *Biochim. Biophys. Acta* **1967**, *146*, 181.

Additions and Corrections

Carbon and Proton Basicity [*J. Am. Chem. Soc.* **1988**, *110*, 5611–5613]. JOHN I. BRAUMAN* and CHAU-CHUNG HAN

This paper showed that proton basicity and methyl cation affinities are better correlated than might have been expected from earlier work, particularly if one used a non-unit-slope linear correlation of heats of formation of HA and CH₃A molecules. The equations derived in the paper are mathematically correct, but the non-unit-slope correlation between heats of formation of HA and CH₃A is only a statistical one that has no physical significance.

The relationship between proton affinity and methyl cation affinity of various A⁻ depends only on the heats of formation of the HA and CH₃A species

$$\text{MCA}(\text{A}^-) - \text{PA}(\text{A}^-) =$$

$$\Delta H_f^\circ(\text{CH}_3^+) - \Delta H_f^\circ(\text{H}^+) - \Delta H_f^\circ(\text{CH}_3\text{A}) + \Delta H_f^\circ(\text{HA}) \quad (1)$$

where MCA(A⁻) is the methyl cation affinity, and PA(A⁻) is the proton affinity, of A⁻. The specific values of the heats of formation of the various species, however, have no physical content and depend on the arbitrary assignment of $\Delta H_f^\circ = 0$ for the elements in their most stable states at 298 K. The differences in heats of formation of reactants and products in any reaction are independent of the choice of the standard states of the elements. Similarly, the differences in heats of formation on the right-hand side of eq 1 are independent of the choice of standard states.

The suggestion that the correlation of MCA and PA could be improved by allowing a non-unit-slope correlation of heats of

formation of HA and CH₃A is thus true in a statistical sense only. Although we can improve the correlation of the heats of formation by allowing a non-unit slope, the slope, and the standard deviation of the correlation, will depend on the arbitrary choice of standard states of the elements.

In short, the correlation of heats of formation of CH₃A and HA, Figure 2 and eq 4 of this paper, is simply a statistical one. The correlation, and the equation derived from it for calculating MCA from PA, eq 5 of this paper, has neither predictive content nor chemical pertinence.

It remains true, however, that there is a rough unit-slope correlation between heats of formation of HA and CH₃A which is equivalent to that between PA and MCA for the various A⁻. The deviations from these unit-slope correlations, as originally pointed out by Hine and Weimer,¹ can, however, be quite large. Other relationships of proton affinities with metal cation affinities that have been reported recently² were found to have unit slope so that their significance is not compromised by the arguments presented here.

Acknowledgment. We are indebted to Professor Calvin D. Ritchie for discovering this problem and his helpful advice regarding it.

(1) Hine, J.; Weimer, R. D., Jr. *J. Am. Chem. Soc.* **1965**, *87*, 3387.

(2) Bryndza, H. E.; Fong, L. K.; Paciellow, R. A.; Tam, W.; Bercaw, J. E. *J. Am. Chem. Soc.* **1987**, *109*, 1444. Labinger, J. A.; Bercaw, J. *Organometallics* **1988**, *7*, 926.