

# Kinetics of Hydrolysis of Iodine Monochloride Measured by the Pulsed-Accelerated-Flow Method

Yi Lai Wang, Julius C. Nagy, and Dale W. Margerum\*

Contribution from the Department of Chemistry, Purdue University, West Lafayette, Indiana 47907. Received March 27, 1989



**Abstract:** First-order rate constants from 75 000 to 150 000 s<sup>-1</sup> are measured by the pulsed-accelerated-flow technique for the rate of loss of ICl<sub>2</sub><sup>-</sup> in neutral and basic solutions. The rate varies inversely with the Cl<sup>-</sup> concentration because ICl(aq) is the reactive form. The hydrolysis of ICl(aq) to give HOI and Cl<sup>-</sup> is extremely rapid ( $k_{\text{H}_2\text{O}} = 2.4 \times 10^6 \text{ s}^{-1}$  at 25.0 °C,  $\mu = 0.50$ ). Hydroxide ion accelerates the rate of hydrolysis. The second-order rate constant for the reaction of OH<sup>-</sup> with ICl(aq) is very large ( $k_{\text{OH}} = 4.5 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ). Reactions of ICl(aq) with H<sub>2</sub>PO<sub>4</sub><sup>-</sup>/HPO<sub>4</sub><sup>2-</sup> and with HCO<sub>3</sub><sup>-</sup>/CO<sub>3</sub><sup>2-</sup> buffers are also very rapid and are first order in the total buffer concentration, with resulting second-order rate constants,  $(1.3 \pm 0.2) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ , that do not depend on the basicity of the buffer or the ratio of buffer species. Two ICl(aq) species are proposed to be in rapid equilibria: H<sub>2</sub>OICl (82%) and ICl (18%). Kinetic data indicate that ICl (the less hydrated species) is so reactive that it is unselective and reacts with H<sub>2</sub>PO<sub>4</sub><sup>-</sup>, HPO<sub>4</sub><sup>2-</sup>, HCO<sub>3</sub><sup>-</sup>, CO<sub>3</sub><sup>2-</sup>, and OH<sup>-</sup> at the diffusion limit ( $7 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ). The more hydrated species, H<sub>2</sub>OICl, also reacts rapidly with OH<sup>-</sup> ( $3.9 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ) by proton transfer to form H<sub>2</sub>O and HOICl<sup>-</sup>, which rapidly loses Cl<sup>-</sup> to give HOI. Spectrophotometric measurements of the equilibrium constants at 25.0 °C give  $([\text{ICl}(\text{aq})][\text{Cl}^-])/[\text{ICl}_2^-] = 0.013 \text{ M}$  ( $\mu = 1.0$  and 0.5) and  $([\text{HOI}][\text{Cl}^-]^2[\text{H}^+])/[\text{ICl}_2^-] = 1.06 \times 10^{-6} \text{ M}^3$  ( $\mu = 0.5$ ).

Kinetics of the hydrolysis of Cl<sub>2</sub>, Br<sub>2</sub>, and I<sub>2</sub> have been determined by relaxation techniques<sup>1</sup> and for Cl<sub>2</sub> by stopped-flow<sup>2</sup> and continuous-flow<sup>3</sup> methods. However, the kinetics of hydrolysis of interhalogens, such as ICl, have not been examined. This had been due to the lack of a method suitable to the study of reactions that are too fast to measure by stopped-flow methods and cannot be maintained under equilibrium conditions as necessary to apply relaxation methods. The development of the pulsed-accelerated-flow (PAF) method<sup>4-6</sup> permits very fast reactions ( $t_{1/2} \geq 2 \mu\text{s}$ ) to be studied. This makes it possible for the first time to study the kinetics and mechanism of iodine monochloride hydrolysis. The PAF method uses integrating observation of the reaction mixture under continuous-flow conditions for short time periods (pulse) during which the flow is accelerated.<sup>4,5</sup> The reactants are observed as they mix in order to eliminate the dead time (time between mixing and observation) that is typically 3-5 ms for stopped-flow instruments.<sup>7</sup> Variation of the flow velocity in the PAF instrument enables the physical mixing process and the chemical reaction process that occur in the observation tube to be resolved from each other so that the rate constant of the actual chemical reaction can be calculated.<sup>4,5</sup> The PAF method has been applied in studies of a number of very fast non-metal redox reactions.<sup>8-10</sup>


Iodine monochloride is proposed as an intermediate in oxidation reactions of iodide by chloramines,<sup>11</sup> nitrogen trichloride,<sup>9</sup> and hypochlorous acid.<sup>9,11</sup> All these reactions involve the transfer of a Cl<sup>+</sup> ion to I<sup>-</sup>. Iodine monochloride also is proposed as a key

intermediate<sup>12</sup> in the chlorite-iodide oscillating reaction system.<sup>13,14</sup> It has long been known that iodine monochloride is labile and that it hydrolyzes extensively in aqueous solutions.<sup>15</sup> In a study of the autocatalytic reaction between I<sub>2</sub> and ClO<sub>2</sub><sup>-</sup>, an estimated value of 100 s<sup>-1</sup> is given as the rate constant for ICl hydrolysis.<sup>14</sup> We shall see that this estimate is too small by a factor of  $2.4 \times 10^4$ . Nevertheless, the hydrolysis path was not observed in the study of the ICl<sub>2</sub><sup>-</sup> reaction with I<sup>-</sup> in 0.1-1.0 M Cl<sup>-</sup> concentration for either 0.10 or 1.0 M H<sup>+</sup> concentrations.<sup>8</sup> It is therefore of interest to measure how rapidly iodine monochloride hydrolyzes and to determine the reaction mechanism for this process.


The final products of the hydrolysis of ICl in acid are iodate and iodine (eq 1a) and in base are iodate and iodide (eq 1b).




Iodine monochloride in water is written as ICl(aq) to indicate that it includes both ICl and H<sub>2</sub>OICl. The latter species (I) will have two electron-pair bonds and three nonbonding electron pairs around iodine in a structure similar to ICl<sub>2</sub><sup>-</sup> (II). It is apparent



that the overall reactions in eq 1a,b involve multiple steps because the products are derived from disproportionation as well as hydrolysis reactions. The initial product of ICl hydrolysis is HOI (eq 2). The disproportionation of HOI in acid gives IO<sub>3</sub><sup>-</sup> and I<sub>2</sub>.<sup>16-18</sup> Kinetic studies have shown that the disproportionation of HOI is both slow and complex.<sup>17</sup>




Iodine monochloride readily forms dichloroiodate(I) (eq 3), ICl<sub>2</sub><sup>-</sup>, which is very stable in solutions where the HCl concentration is 0.5 M or greater.<sup>19</sup> We will show that the value of the




- (1) Eigen, M.; Kustin, K. *J. Am. Chem. Soc.* **1962**, *84*, 1355-1361.  
 (2) Margerum, D. W.; Gray, E. T., Jr.; Huffman, R. P. In *Organometals and Organometalloids, Occurrence and Fate in the Environment*; Brinckman, F. E., Bellama, J. M., Eds.; ACS Symposium Series 82; American Chemical Society: Washington, DC, 1978; pp 278-291.  
 (3) Lifshitz, A.; Perlmutter-Haymann, B. *J. Phys. Chem.* **1960**, *64*, 1663-1665.  
 (4) Jacobs, S. A.; Nemeth, M. T.; Kramer, G. W.; Ridley, T. Y.; Margerum, D. W. *Anal. Chem.* **1984**, *56*, 1058-1065.  
 (5) Nemeth, M. T.; Fogelman, K. D.; Ridley, T. Y.; Margerum, D. W. *Anal. Chem.* **1987**, *59*, 283-291.  
 (6) Bowers, C. P.; Fogelman, K. D.; Nagy, J. C.; Ridley, T. Y.; Wang, Y. L.; Margerum, D. W. To be submitted for publication.  
 (7) Dickson, P. N.; Margerum, D. W. *Anal. Chem.* **1986**, *58*, 3153-3158.  
 (8) Margerum, D. W.; Dickson, P. N.; Nagy, J. C.; Kumar, K.; Bowers, C. P.; Fogelman, K. D. *Inorg. Chem.* **1986**, *25*, 4900-4904.  
 (9) Nagy, J. C.; Kumar, K.; Margerum, D. W. *Inorg. Chem.* **1988**, *27*, 2773-2780.  
 (10) Fogelman, K. D.; Walker, D. M.; Margerum, D. W. *Inorg. Chem.* **1989**, *28*, 986-993.  
 (11) Kumar, K.; Day, R. A.; Margerum, D. W. *Inorg. Chem.* **1986**, *25*, 4344-4350.

- (12) Beck, M. T.; Rabai, G. *J. Phys. Chem.* **1986**, *90*, 2204-2205.  
 (13) Citri, O.; Epstein, I. R. *J. Phys. Chem.* **1987**, *91*, 6034-6040.  
 (14) Rabai, G.; Beck, M. T. *Inorg. Chem.* **1987**, *26*, 1195-1199.  
 (15) Faull, J. H. *J. Am. Chem. Soc.* **1934**, *56*, 522-526.  
 (16) Li, C. H.; White, C. F. *J. Am. Chem. Soc.* **1943**, *65*, 335-339.  
 (17) Thomas, T. R.; Pence, D. T.; Hasty, R. A. *J. Inorg. Nucl. Chem.* **1980**, *42*, 183-186.  
 (18) Wren, J. C.; Paquette, J.; Sander, S.; Ford, B. L. *Can. J. Chem.* **1986**, *64*, 2284-2296.



**Figure 1.** (a) Absorbance change in the spectrophotometric titration of  $\text{ICl}_2^-$  with 1.00 M HCl ( $V_{\text{HCl}}$ ). Conditions: 2.50 mL of  $1.90 \times 10^{-3}$  M  $[\text{ICl}_2^-]_{\text{init}}$  in 1.00 M  $\text{HClO}_4$  at 25.0 °C, 1.00-cm cell, 343 nm. (b) Absorbance versus the iteratively calculated chloride concentration.

equilibrium constant  $K_3$  is 0.013 M, which differs significantly from the long-established value of 0.006 M.<sup>15</sup> The UV spectrum of  $\text{ICl}_2^-$  has characteristic absorption bands at 224 nm ( $\epsilon = 47\,000 \text{ M}^{-1} \text{ cm}^{-1}$ )<sup>19</sup> and 343 nm, where  $\epsilon$  ( $\text{M}^{-1} \text{ cm}^{-1}$ ) values of 243<sup>19</sup> and 275<sup>20</sup> have been reported. Our results indicate that the latter value is correct. Equations 2 and 3 can be combined to give the equilibrium in terms of  $\text{ICl}_2^-$  and  $\text{HOI}$  (eq 4). We have measured




this equilibrium constant and find a  $K_4$  value of  $1.06 \times 10^{-6} \text{ M}^3$ . A  $K_2$  value of  $8.2 \times 10^{-5} \text{ M}^2$  can be calculated from our values of  $K_3$  and  $K_4$ . These equilibrium constants are obtained by more direct measurements than those previously reported.<sup>15</sup>

The hydrolysis of  $\text{ICl}$  can be suppressed by  $\text{Cl}^-$  due to the formation of  $\text{ICl}_2^-$ , but the hydrolysis is nevertheless far too rapid to be measured by stopped-flow spectrophotometry. The PAF method extends accessible first-order rate constants by a factor of  $10^3$  beyond the limit of stopped-flow methods. This makes it possible to measure the rate of reaction of iodine monochloride with water, with hydroxide ion, and with phosphate and carbonate buffers.

### Experimental Section

A 0.060 M stock solution of  $\text{ICl}_2^-$  was prepared<sup>8,21</sup> by addition of standardized<sup>22</sup> NaI solution to a solution of primary standard-grade  $\text{NaIO}_3$  in HCl. Two moles of NaI was added for each mole of  $\text{NaIO}_3$  (eq 5). A brown precipitate of  $\text{I}_2$  formed initially but redissolved with



stirring after 30 min. The final concentrations of  $\text{H}^+$  and  $\text{Cl}^-$  are 0.20 and 1.0 M, respectively, and under these conditions the  $\text{ICl}_2^-$  stock solution is very stable. The  $\text{ICl}_2^-$  solutions used in the hydrolysis experiments were made by dilution of the stock solution with the reaction medium immediately prior to the experiments.

Solutions of  $\text{HClO}_4$  and HCl were standardized against NaOH, which had been standardized with primary standard potassium hydrogen phthalate. Solutions used in the PAF kinetic experiments were filtered through 0.22- $\mu\text{m}$  Triton-free Millipore type HA filter and were degassed by aspiration for at least 10 min. An Orion 601A pH meter with a combination electrode was used to measure the pH of solutions after the reactions were complete. These readings were corrected to  $-\log [\text{H}^+]$  values on the basis of electrode calibration by titration of standardized  $\text{HClO}_4$  with standardized NaOH at  $\mu = 0.50$  ( $\text{NaClO}_4$ ) and 25.0 °C.

A Perkin-Elmer Model 320 spectrophotometer was used to acquire UV-vis spectra and to obtain spectrophotometric titration data. The titration was carried out by adding HCl to a dilute acidic  $\text{ICl}_2^-$  solution.

(19) Cason, D. L.; Neumann, H. M. *J. Am. Chem. Soc.* **1961**, *83*, 1822–1828.

(20) Gillam, A. E.; Morton, R. A. *Proc. R. Soc. London, A* **1931**, *132*, 152–167.

(21) Philbrick, F. A. *J. Chem. Soc.* **1930**, 2254–2260.

(22) Kolthoff, I. M.; Sandell, E. B.; Meehan, E. J.; Bruckenstein, S. *Quantitative Chemical Analysis*, 4th ed.; Macmillan: London, 1969; p 852.

**Table I.** Stopped-Flow Determination of  $K_4$  Based on the Immediate Absorbance Decrease on Mixing  $\text{ICl}_2^-$  with Buffers<sup>a</sup>

$[\text{Cl}^-]$ , M	$\text{p}[\text{H}^+]$	$A_0(343 \text{ nm})$	$[\text{Cl}^-]$ , M	$\text{p}[\text{H}^+]$	$A_0(343 \text{ nm})$
0.3	3.56 <sup>b</sup>	0.218	0.4	5.51 <sup>c</sup>	0.109
0.3	4.12 <sup>b</sup>	0.199	0.4	6.05 <sup>c</sup>	0.074
0.3	4.64 <sup>b</sup>	0.167	0.4	6.54 <sup>c</sup>	0.061
0.3	5.06 <sup>b</sup>	0.128	0.5	3.99 <sup>b</sup>	0.215
0.3	5.61 <sup>c</sup>	0.087	0.5	4.97 <sup>b</sup>	0.172
0.3	6.08 <sup>c</sup>	0.067	0.5	5.29 <sup>b</sup>	0.146
0.4	3.59 <sup>b</sup>	0.221	0.5	5.98 <sup>c</sup>	0.084
0.4	4.17 <sup>b</sup>	0.204	0.5	6.44 <sup>c</sup>	0.068
0.4	4.66 <sup>b</sup>	0.179	0.5	6.85 <sup>c</sup>	0.060
0.4	5.08 <sup>b</sup>	0.146			

<sup>a</sup> 25.0 °C,  $\mu = 0.50$ ,  $[\text{ICl}_2^-]_0 = 4.3 \times 10^{-4} \text{ M}$ ,  $A_0 = 0.223$ , 1.89-cm cell. <sup>b</sup>  $[\text{OAc}^-]_{\text{T}} = 0.05 \text{ M}$ . <sup>c</sup>  $[\text{PO}_4]_{\text{T}} = 0.05 \text{ M}$ .

A stoppered 1-cm spectral cell (4-mL capacity) was loaded with 2.50 mL of a  $1.9 \times 10^{-3} \text{ M}$   $\text{ICl}_2^-$  solution in 1.0 M  $\text{HClO}_4$ . The absorbance at 343 nm was measured as small increments of 1.0 M HCl were added to give a final volume of 3.89 mL. The absorbance during the titration increased initially as more  $\text{ICl}_2^-$  formed and then decreased due to dilution (Figure 1a).

A Durrum stopped-flow spectrometer interfaced<sup>23</sup> to a Hewlett-Packard 2100S minicomputer was used to determine the total change of absorbance of  $\text{ICl}_2^-$  after it was mixed with phosphate buffer. The rate of loss of  $\text{ICl}_2^-$  is too fast to be measured by this method, but the equilibrium in eq 4 can be established within 5–20 ms without significant disproportionation of  $\text{HOI}$ .


A pulsed-accelerated-flow spectrometer was used to obtain kinetic data for the reaction of  $\text{ICl}_2^-$  with dilute NaOH solutions, with phosphate buffer, and with carbonate buffer. This instrument employs integrating observation with continuous flow<sup>4</sup> over a short time period to conserve reagents (only 5–6 mL is needed per push). Key features of the PAF spectrometer include a twin-path mixing/observation cell<sup>5</sup> and a programmed-flow velocity. The two reactant solutions enter the observation cell via 10 jets, which intersect in the middle of the observation tube, and the reacting mixture then flows parallel to the direction of observation.<sup>5</sup> The velocity of the solution in the observation tube is programmed to change from 3.0  $\text{m s}^{-1}$  at the beginning of the push to 12.5  $\text{m s}^{-1}$  at the end of the push in the acceleration mode or from 12.5 to 3.0  $\text{m s}^{-1}$  in the deceleration mode. Turbulent flow that is necessary for mixing the solutions exists at these velocities, and the mixing rate constant is proportional to the velocity. The change of the velocity permits resolution of the physical mixing process and the chemical reaction process. Calibration reactions have been used to verify the resolved rate constants.<sup>4,5</sup> The Model IV PAF instrument used in this work has a double-grating monochromator with a wavelength range of 200–850 nm and minimal stray light. The instrumental features have been summarized in a previous application.<sup>10</sup> Details of the construction, computer interfacing, and measurement of rate constants as large as 350 000  $\text{s}^{-1}$  will be provided elsewhere.<sup>6</sup> In the present work first-order rate constants are measured in the range 75 000–150 000  $\text{s}^{-1}$  by the PAF method.

All PAF experiments were carried out under pseudo-first-order conditions at 25.0 °C and  $\mu = 0.50$  ( $\text{NaClO}_4$ ). Reactions were monitored at 230 nm in order to take advantage of the high molar absorptivity of  $\text{ICl}_2^-$  ( $\epsilon_{230} = 42\,000 \text{ M}^{-1} \text{ cm}^{-1}$ ) compared to the much lower absorptivities of  $\text{HOI}$  and  $\text{OI}^-$  at this wavelength. A typical initial absorbance ( $A_0$ ) was 1.22 ( $[\text{ICl}_2^-] = 1.4 \times 10^{-3} \text{ M}$ , 2.05-cm cell), and typical  $A_\infty$  values were 0.11. The observed first-order rate constant,  $k_r$ , is the result of a linear regression of the  $M_{\text{exptl}}^{5,10}$  values versus the flow velocity in the observation tube.  $M_{\text{exptl}} = (A_v - A_\infty)/(A_0 - A_\infty)$ , where  $A_v$  is the instantaneous absorbance for each of 250 points taken over a velocity range of 12–7  $\text{m s}^{-1}$ . These high velocities are needed to give linear plots of  $M_{\text{exptl}}$  against  $v$  because of the magnitude of the rate constants in this study. The rate constants are corrected in accordance with the calibration of the PAF instrument for large  $k_r$  values ( $>100\,000 \text{ s}^{-1}$ ).<sup>6</sup> The reported  $k_r$  values and standard deviations are obtained from at least seven trials for each set of conditions. The average relative standard deviation is  $\pm 9\%$ .

### Results and Discussion

**Equilibrium Constant for the Hydrolysis of  $\text{ICl}_2^-$ .** A stopped-flow instrument was used to study the extent of hydrolysis of  $\text{ICl}_2^-$  to give  $\text{HOI}$  (eq 4) in the  $\text{p}[\text{H}^+]$  range from 3.56 to 6.91, with  $[\text{Cl}^-] = 0.30\text{--}0.50 \text{ M}$ . The loss of  $\text{ICl}_2^-$  was monitored at 343 nm. The

(23) Wills, B. G.; Bittikofer, J. A.; Pardue, H. L.; Margerum, D. W. *Anal. Chem.* **1970**, *42*, 1340–1349.



**Figure 2.** Plot of eq 7 with slope =  $K_4 = (1.06 \pm 0.03) \times 10^{-6} \text{ M}^3$  at 25.0 °C,  $\mu = 0.50$ , based on the molar absorptivities given in Table 11. The inset shows the validity of the fit at lower HOI concentrations.

initial absorbance change takes place within the dead time of the instrument (<4 ms). The stopped-flow spectrometer was used so that the subsequent disproportionation of HOI did not interfere. The data (Table I) show that at a given  $[\text{Cl}^-]$  the loss of  $\text{ICl}_2^-$  increases as  $p[\text{H}^+]$  increases as expected from eq 4. The equilibrium constant  $K_4$  is defined by eq 6. The ratio of [HOI] to


$$K_4 = \frac{[\text{Cl}^-]^2[\text{H}^+][\text{HOI}]}{[\text{ICl}_2^-]} \quad (6)$$

$[\text{ICl}_2^-]$  can be obtained from the original absorbance of the  $\text{ICl}_2^-$  solution ( $A_0 = b\epsilon_{\text{ICl}_2^-}[\text{ICl}_2^-]_0$ ) and the equilibrium absorbance ( $A_e = b(\epsilon_{\text{ICl}_2^-}[\text{ICl}_2^-]_e + \epsilon_{\text{HOI}}[\text{HOI}]_e)$ ). A plot of the ratio against  $1/([\text{H}^+][\text{Cl}^-]^2)$  should give a straight line with slope of  $K_4$ . In eq 7, the molar absorptivity value for  $\text{ICl}_2^-$  is  $275 \text{ M}^{-1} \text{ cm}^{-1}$  but

$$\frac{[\text{HOI}]}{[\text{ICl}_2^-]} = \frac{A_0 - A_e}{A_e - A_0(\epsilon_{\text{HOI}}/\epsilon_{\text{ICl}_2^-})} = \frac{K_4}{[\text{H}^+][\text{Cl}^-]^2} \quad (7)$$

the  $\epsilon$  value for HOI at 343 nm has not been reported. However,  $\epsilon$  values for HOI and  $\text{OI}^-$  at other UV wavelengths are available. For HOI, the known value is  $\epsilon_{278}^{\text{max}} = 200$ ,<sup>24</sup> and for  $\text{OI}^-$ ,  $\epsilon_{365} = 5824$  and  $\epsilon_{363} = 60$ .<sup>18</sup> We tested a range of the  $\epsilon$  values from 60 to  $70 \text{ M}^{-1} \text{ cm}^{-1}$  for HOI at 343 nm in order to give a linear fit for eq 7. The results of the weighted linear regression show that the best fit (Figure 2) is achieved when the  $\epsilon$  value for HOI at 343 nm is  $67 \text{ M}^{-1} \text{ cm}^{-1}$ . The slope from the regression treatment gives a new value of  $K_4$  equal to  $(1.06 \pm 0.03) \times 10^{-6} \text{ M}^3$  (25.0 °C,  $\mu = 0.50$ ). This is the first direct measurement of this equilibrium constant, as previous measurements had interference from the disproportionation of HOI. Our experimental value for  $K_4$  is larger than the previously estimated value ( $\sim 6 \times 10^{-7} \text{ M}^3$ ).<sup>15</sup>

**Equilibrium Constant for  $\text{ICl}_2^-/\text{ICl}$ .** The equilibrium constant for the hydrolysis of  $\text{ICl}_2^-$  indicates that the concentration of HOI is only 1% of the initial  $\text{ICl}_2^-$  concentration in 1.0 M  $\text{H}^+$  and 0.010 M  $\text{Cl}^-$ . However,  $\text{ICl}_2^-$  will partially dissociate into  $\text{ICl}(\text{aq})$  and  $\text{Cl}^-$  under these conditions, and another species,  $\text{H}_2\text{OI}^+$ , also must be considered. Bell and Gelles<sup>25</sup> found that the hydrated iodine cation is in equilibrium with HOI in highly acidic solutions (eq 8). The equilibrium constant,  $K_5$ , is  $35 \text{ M}^{-1}$ . Since the formation



of  $\text{H}_2\text{OI}^+$  is favorable in high acid, it needs to be considered in the calculation of the equilibrium constant for  $\text{ICl}_2^-/\text{ICl}$ . Therefore, three equilibria (eq 3, 4, and 8) must be considered in the calculation of the concentration of each species. The concentration of HOI can be neglected since it is only 2.8% of the  $\text{H}_2\text{OI}^+$  concentration in 1.0 M acid. The concentrations of

$\text{ICl}_2^-$ ,  $\text{ICl}(\text{aq})$ , and  $\text{H}_2\text{OI}^+$  can be derived as shown by eq 9–11, where  $[\text{ICl}_2^-]_{\text{init}} = [\text{ICl}_2^-] + [\text{ICl}(\text{aq})] + [\text{H}_2\text{OI}^+]$ ,  $K_2K_3 = K_4 = 1.06 \times 10^{-6} \text{ M}$ , and  $K_5 = 35 \text{ M}^{-1}$ .

$$[\text{ICl}_2^-] = \frac{[\text{ICl}_2^-]_{\text{init}}[\text{Cl}^-]^2}{K_4K_5 + K_3[\text{Cl}^-] + [\text{Cl}^-]^2} \quad (9)$$

$$[\text{ICl}(\text{aq})] = \frac{[\text{ICl}_2^-]_{\text{init}}K_3[\text{Cl}^-]}{K_4K_5 + K_3[\text{Cl}^-] + [\text{Cl}^-]^2} \quad (10)$$

$$[\text{H}_2\text{OI}^+] = \frac{[\text{ICl}_2^-]_{\text{init}}K_4K_5}{K_4K_5 + K_3[\text{Cl}^-] + [\text{Cl}^-]^2} \quad (11)$$

A spectrophotometric titration with HCl was used to measure the  $K_3$  equilibrium constant. A  $1.9 \times 10^{-3} \text{ M}$   $\text{ICl}_2^-$  solution ( $[\text{Cl}^-] = 0.0020 \text{ M}$ ) in 1.0 M  $\text{HClO}_4$  was titrated with 1.0 M HCl so that the  $[\text{H}^+]$  concentration remained constant while the  $[\text{Cl}^-]$  concentration increased. The titration data are shown in Figure 1a. The increase of absorbance at the first half of the titration indicates the progressive formation of  $\text{ICl}_2^-$  with added  $\text{Cl}^-$ . The subsequent decrease of the absorbance is due to the dilution caused by adding more HCl. The increase of absorbance due to  $\text{ICl}_2^-$  formation as the  $[\text{Cl}^-]$  increases has been observed previously,<sup>19</sup> although another report<sup>20</sup> claimed that no absorbance change was observed under the same conditions.

Assuming that all three species ( $\text{ICl}_2^-$ ,  $\text{ICl}(\text{aq})$ , and  $\text{H}_2\text{OI}^+$ ) absorb at 343 nm, a relationship between the absorbance ( $A_{\text{obsd}}$ ) and  $[\text{Cl}^-]$  can be derived (eq 12 and 13), where the cell path is 1.0 cm. In eq 13,  $K_3$ ,  $\epsilon_{\text{ICl}_2^-}$ ,  $\epsilon_{\text{ICl}}$ , and  $\epsilon_{\text{H}_2\text{OI}^+}$  are the parameters to

$$A_{\text{obsd}} = [\text{ICl}_2^-]\epsilon_{\text{ICl}_2^-} + [\text{ICl}(\text{aq})]\epsilon_{\text{ICl}} + [\text{H}_2\text{OI}^+]\epsilon_{\text{H}_2\text{OI}^+} \quad (12)$$

$$A_{\text{obsd}} = \left[ \frac{[\text{ICl}_2^-]_{\text{init}}}{K_4K_5 + K_3[\text{Cl}^-] + [\text{Cl}^-]^2} \right] \times \left( [\text{Cl}^-]^2\epsilon_{\text{ICl}_2^-} + K_3[\text{Cl}^-]\epsilon_{\text{ICl}} + K_4K_5\epsilon_{\text{H}_2\text{OI}^+} \right) \quad (13)$$

be calculated,  $[\text{Cl}^-]$  and  $A_{\text{obsd}}$  are the independent and dependent variables, and  $K_4$  ( $1.06 \times 10^{-6} \text{ M}^3$ ) and  $K_5$  ( $35 \text{ M}^{-1}$ ) are known values. The parameters can be resolved when eq 13 is fitted to the titration data by use of a nonlinear regression curve-fitting procedure, which employs a Marquardt algorithm<sup>26</sup> to change the initially estimated values of the parameters iteratively until the minimum on the  $\chi^2$  surface is found. The fitting, however, needs a value for the total chloride concentration,  $[\text{Cl}^-]$ , in the solution, which comes from three sources: One source is the original  $\text{Cl}^-$  in the solution to be titrated, the second source is from the HCl added during the titration, and the third source is from the dissociation of  $\text{ICl}_2^-$ . To calculate the  $\text{Cl}^-$  contribution,  $[\text{Cl}^-]_d$ , from the dissociation of  $\text{ICl}_2^-$  a quadratic equation (eq 14) must be

$$[\text{Cl}^-]_d = \left[ -([\text{Cl}^-] + K_3) + \left\{ ([\text{Cl}^-] + K_3)^2 + 4K_3[\text{ICl}_2^-]_{\text{init}} \right\}^{1/2} \right] / 2 \quad (14)$$

solved. This requires knowledge of  $K_3$ , which in turn, is not known until the  $[\text{Cl}^-]$  can be calculated. This circular logic situation is resolved iteratively. The fitting function (eq 13) is expanded to include a nested iterative process that calculates  $[\text{Cl}^-]_d$  from eq 14 by using an assumed  $K_3$  value. The  $[\text{Cl}^-]_d$  value is then added to the other sources of chloride ion to give  $[\text{Cl}^-]$ , and a new  $[\text{Cl}^-]_d$  value is calculated. This process continues until  $[\text{Cl}^-]$  does not change. The total  $\text{Cl}^-$  concentration,  $[\text{Cl}^-]$ , is then substituted into the fitting function (eq 13) to produce a new set of parameters including  $K_3$ . In the second iteration the new  $K_3$  value is used to calculate  $[\text{Cl}^-]$  again in the process just described. The curve-fitting procedure eventually converges upon a set of parameters that best fit the data. The line through the data points in Figure 1a is the result of this fit in terms of the absorbance versus the volume of HCl added. Figure 1b is the plot of absorbance against the calculated  $[\text{Cl}^-]$ . The fitted parameters are  $K_3 = 0.013 \pm 0.001 \text{ M}$  (25.0 °C,  $\mu = 1.0$ ) with molar absorptivity

(24) Palmer, D. A.; Van Eldik, R. *Inorg. Chem.* **1986**, *25*, 928–931.

(25) Bell, R. P.; Gelles, E. *J. Chem. Soc.* **1951**, 2734–2740.

(26) Bevington, P. R. *Data Reduction and Error Analysis for the Physical Sciences*; McGraw-Hill: New York, 1969; p 235.

**Table II.** Equilibrium Constants and Molar Absorptivities of Iodine(I) Species in Aqueous Solutions<sup>a</sup>

Equilibration Constants		
$K_3 = ([\text{ICl}(\text{aq})][\text{Cl}^-])/[\text{ICl}_2^-]$	$= (1.3 \pm 0.1) \times 10^{-2} \text{ M}$	
$K_4 = ([\text{HOI}][\text{Cl}^-]^2[\text{H}^+])/[\text{ICl}_2^-]$	$= (1.06 \pm 0.03) \times 10^{-6} \text{ M}^3$	
$K_2 = ([\text{HOI}][\text{Cl}^-][\text{H}^+])/[\text{ICl}(\text{aq})]$	$= 8.2 \times 10^{-5} \text{ M}^2$ <sup>b</sup>	
$K_5 = [\text{H}_2\text{OI}^+]/([\text{HOI}][\text{H}^+])$	$= 35 \text{ M}^{-1}$ <sup>c</sup>	
$[\text{H}_2\text{OICl}]/[\text{ICl}]$	$= 4.5$ <sup>d</sup>	
Molar Absorptivities ( $\text{M}^{-1} \text{ cm}^{-1}$ )		
species	343 nm	230 nm
$\text{ICl}_2^-$	$274 \pm 3$	$(4.2 \pm 0.5) \times 10^4$
$\text{ICl}(\text{aq})$	$176 \pm 8$	
$\text{HOI}$	$67 \pm 2$ <sup>e</sup>	$(4 \pm 1) \times 10^3$
$\text{H}_2\text{OI}^+$	$149 \pm 7$	



<sup>a</sup> Conditions: 25.0 °C,  $\mu = 0.50$ ,  $[\text{ICl}(\text{aq})] = [\text{ICl}] + [\text{H}_2\text{OICl}]$ .  
<sup>b</sup> Calculated from  $K_3$  and  $K_4$ . <sup>c</sup> Reference 25. <sup>d</sup> Ratio is estimated from kinetic behavior of  $\text{ICl}(\text{aq})$  with bases, where  $[\text{ICl}(\text{aq})] = [\text{H}_2\text{OICl}] + [\text{ICl}]$ . <sup>e</sup> The uncertainty is estimated from the best fit of the data in Figure 2.

values at 343 nm for  $\epsilon_{\text{ICl}_2^-} = 274 \pm 3$ ,  $\epsilon_{\text{ICl}} = 176 \pm 8$ , and  $\epsilon_{\text{H}_2\text{OI}^+} = 149 \pm 7$ . All values are averages from four experiments. The value of  $K_3$  is the same, within experimental error, at  $\mu = 0.50$  as it is at  $\mu = 1.0$ . The  $\epsilon$  value found for  $\text{ICl}_2^-$  agrees within experimental error with the earlier report<sup>20</sup> of  $275 \text{ M}^{-1} \text{ cm}^{-1}$ .

The results show that  $\text{ICl}(\text{aq})$  absorbs significantly at 343 nm with  $\epsilon = 176 \text{ M}^{-1} \text{ cm}^{-1}$ . This value is comparable to the  $\epsilon$  values given<sup>27</sup> for  $\text{ICl}$  in polar organic solvents: 181 at 343 nm in  $\text{C}_2\text{H}_5\text{OH}$ , 189 at 344 nm in  $\text{CH}_3\text{OH}$ , and 188 at 346 nm in  $(\text{CH}_3)_3\text{COH}$ . The absorption band is due to a  $\pi^* \rightarrow \sigma^*$  transition.<sup>28</sup> The molar absorptivity value for  $\text{H}_2\text{OI}^+$  ( $149 \text{ M}^{-1} \text{ cm}^{-1}$  at 343 nm) is larger than that of  $\text{HOI}$  ( $67 \text{ M}^{-1} \text{ cm}^{-1}$  at 343 nm).


The equilibrium constant,  $K_3$ , differs by slightly more than a factor of 2 from the value of 0.0060 M that was reported previously.<sup>15</sup> The previous method was less accurate because it involved more indirect measurements of different constants or concentrations, such as the distribution coefficients for  $\text{ICl}$  between aqueous and  $\text{CCl}_4$  solutions, and because the  $\text{H}_2\text{OI}^+$  contribution was neglected. Our determination of the overall equilibrium constant  $K_4$  for the hydrolysis of  $\text{ICl}_2^-$  to  $\text{HOI}$  also disagrees with the value estimated by Faull.<sup>15</sup> Thermodynamic data for  $\text{ICl}_2^-$ ,  $\text{ICl}$ , and  $\text{IOH}$  and values of electrode potentials<sup>29</sup> all appear to be based on Faull's values.<sup>15</sup> We recommend use of the revised values for the equilibrium constants that are given in Table II.

**Chloride Ion Suppression of the Rate of Hydrolysis of Iodine Monochloride.** The reaction in eq 4 is too fast for the stopped-flow time scale, but it can be measured by the pulsed-accelerated-flow method. A wavelength of 230 nm was selected to observe the reaction in order to take advantage of the large absorbance decrease as  $\text{ICl}_2^-$  is converted to  $\text{HOI}$  (Table II). Pseudo-first-order rate constants ( $k_r$ ) decrease from 150 000 to 77 000  $\text{s}^{-1}$  as the chloride ion concentration increases from 0.25 to 0.50 M (Figure 3). The  $[\text{ICl}_2^-]_{\text{init}}$  is  $3.0 \times 10^{-5} \text{ M}$ , and the hydrolysis of  $\text{ICl}_2^-$  occurs in dilute phosphate buffer at  $\text{p}[\text{H}^+] = 6.50$ . The results are consistent with a rapid preequilibrium between  $\text{ICl}(\text{aq})$  and  $\text{ICl}_2^-$  followed by a rate-determining hydrolysis reaction (eq 15). The curved line in Figure 3 is a fit to eq 16 where  $k'$  is  $3.1 \times 10^6 \text{ s}^{-1}$  and  $K_3$  is 0.013 M.



$$k_r = k'K_3/(K_3 + [\text{Cl}^-]) \quad (16)$$

**Hydroxide Ion Effect.** The rate of hydrolysis of  $\text{ICl}_2^-$  increases greatly with increase in hydroxide ion concentration. The reaction

**Figure 3.** Rate constants from PAF measurements of the hydrolysis of  $\text{ICl}_2^-$  as a function of chloride ion concentration. Conditions: 25.0 °C, 0.50  $\mu\text{M}$ ,  $[\text{ICl}_2^-]_{\text{init}} = 3.0 \times 10^{-5} \text{ M}$ ,  $[\text{PO}_4]_{\text{T}} = 0.6 \times 10^{-3} \text{ M}$ ,  $\text{p}[\text{H}^+] = 6.50$ , 230 nm.**Table III.** Effect of Hydroxide Ion and Buffers on the First-Order Rate Constant for the Hydrolysis of  $\text{ICl}_2^-$  Measured by the PAF Method<sup>a</sup>


$\text{p}[\text{H}^+]$	$10^3[\text{B}], \text{M}$	$10^5 k_r, \text{s}^{-1}$
<b>B = OH<sup>-</sup></b>		
	0.27	$0.93 \pm 0.07$
	0.41	$1.1 \pm 0.1$
	0.51	$1.2 \pm 0.1$
<b>B = H<sub>2</sub>PO<sub>4</sub><sup>-</sup> + HPO<sub>4</sub><sup>2-</sup></b>		
6.11	1.00	$0.91 \pm 0.08$
6.20	1.25	$1.0 \pm 0.1$
6.14	1.50	$1.0 \pm 0.1$
6.14	2.00	$1.2 \pm 0.1$
6.48	0.60	$0.77 \pm 0.07$
6.49	1.20	$1.0 \pm 0.1$
6.50	1.80	$1.2 \pm 0.1$
6.50	2.40	$1.3 \pm 0.1$
6.50	3.00	$1.5 \pm 0.2$
7.11	0.50	$0.75 \pm 0.06$
7.11	0.75	$0.82 \pm 0.06$
7.10	1.00	$0.89 \pm 0.07$
7.11	1.50	$1.1 \pm 0.1$
7.12	2.00	$1.2 \pm 0.1$
7.09	3.00	$1.5 \pm 0.2$
<b>B = HCO<sub>3</sub><sup>-</sup> + CO<sub>3</sub><sup>2-</sup></b>		
9.19	0.75	$0.94 \pm 0.08$
9.19	1.00	$1.0 \pm 0.1$
9.18	1.25	$1.1 \pm 0.1$
9.18	1.50	$1.2 \pm 0.1$
9.17	2.00	$1.4 \pm 0.2$
9.68	0.90	$1.04 \pm 0.09$
9.69	1.20	$1.1 \pm 0.1$
9.68	1.50	$1.3 \pm 0.1$

<sup>a</sup>  $[\text{Cl}^-] = 0.50 \text{ M}$ , 25.0 °C,  $\mu = 0.5$ ,  $[\text{ICl}_2^-]_{\text{init}} = 1.1 \times 10^{-5} \text{ M}$  for reactions with  $\text{OH}^-$  and  $3.0 \times 10^{-5} \text{ M}$  for reactions in buffers.  $\text{p}K_a(\text{H}_2\text{PO}_4^-) = 6.46$  (Mesmer, R. E.; Baes, C. F. *J. Solution Chem.* **1974**, *3*, 307–321) and  $\text{p}K_a(\text{HCO}_3^-) = 9.70$  (Odier, M.; Plichon, V. *Anal. Chim. Acta* **1971**, *55*, 209–220; ref 10).

is so rapid that  $[\text{Cl}^-]$  is adjusted to 0.50 M, and even with  $[\text{OH}^-]$  concentrations of only 0.27–0.51 mM the  $k_r$  values increase from 93 000 to 120 000  $\text{s}^{-1}$  (Table III). A linear regression of the  $k_r$  values as a function of hydroxide ion concentration is shown in Figure 4a, where the intercept is  $(6.2 \pm 0.2) \times 10^4 \text{ s}^{-1}$  and the slope is  $(1.1 \pm 0.1) \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ . A chloride-independent rate constant for the hydroxide path can be calculated from the product of the slope  $\times (K_3 + [\text{Cl}^-])/K_3$ . This gives a value of  $(4.5 \pm 0.4) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  for the reaction of  $\text{ICl}(\text{aq})$  with  $\text{OH}^-$ .


Because  $\text{ICl}$  is a strong Lewis acid,<sup>30</sup> it is very easy for it to accept lone-pair (or  $\pi$ ) electrons from electron-donating solvents. Charge-transfer molecular complexes have been reported<sup>31</sup> to form

(27) Buckles, R. E.; Mills, J. F. *J. Am. Chem. Soc.* **1954**, *76*, 4845–4849.(28) Mason, J. F. *Q. Rev. Chem. Soc.* **1961**, *15*, 287–371.(29) Desideri, P. G.; Lepri, L.; Heimler, D. In *Standard Potentials in Aqueous Solution*; Bard, A. J., Parsons, R., Jordan, J., Eds.; Marcel Dekker: New York, 1985; pp 84–90.(30) Drago, R. S.; Wenz, A. *J. Am. Chem. Soc.* **1962**, *84*, 526–528.



**Figure 4.** Rate constants from PAF measurements of the hydrolysis of  $\text{ICl}_2^-$  as a function of base concentrations.  $[\text{Cl}^-] = 0.50 \text{ M}$ , 230 nm. Key: (a)  $[\text{B}] = [\text{OH}^-]$ ,  $[\text{ICl}_2^-]_{\text{init}} = 1.1 \times 10^{-5} \text{ M}$ ; (b)  $[\text{B}] = [\text{CO}_3]_{\text{T}}$ ,  $\text{p}[\text{H}^+] = 9.2\text{--}9.7$ ; (c)  $[\text{B}] = [\text{PO}_4]_{\text{T}}$ ,  $\text{p}[\text{H}^+] = 6.1\text{--}7.1$ .  $[\text{ICl}_2^-]_{\text{init}} = 3.0 \times 10^{-5} \text{ M}$  for (b) and (c).

**Scheme I.** Pathways for the Hydrolysis of  $\text{ICl}$  with  $\text{H}_2\text{O}$ ,  $\text{OH}^-$ , and Buffers (B)



between electron-donating solvents and  $\text{ICl}$ . It is inevitable that some  $\text{ICl}$  would be specifically hydrated by water to form  $\text{H}_2\text{OICl}$ , as the coordination sphere of the iodine in  $\text{ICl}$  is unsaturated and the oxygen atom of the water molecule can donate its lone-pair electrons to form a complex. The  $\text{H}_2\text{OICl}$  complex will be in equilibrium with  $\text{ICl}$  molecules that maintain an octet of valence electrons around iodine. We will refer to the latter species as unhydrated  $\text{ICl}$  or more accurately as a less hydrated form because it will have dipole-dipole interactions with water. The possible presence of both  $\text{ICl}$  and  $\text{H}_2\text{OICl}$  in water indicates that  $\text{OH}^-$  could react with  $\text{ICl}(\text{aq})$  by a direct nucleophilic attack of  $\text{OH}^-$  on  $\text{ICl}$  or by deprotonation of  $\text{H}_2\text{OICl}$ . Both reactions give the same product,  $\text{HOICl}^-$ , which can dissociate to form  $\text{HOI}$  as shown in Scheme I.

The rate constant for the hydroxide path,  $k_{\text{OH}}$ , is the sum of the two paths defined in eq 17 where  $[\text{ICl}(\text{aq})]$  denotes the sum of the  $[\text{ICl}]$  and  $[\text{H}_2\text{OICl}]$  concentrations, and  $k_{\text{dOH}}$  and  $k_{\text{bOH}}$  are the rate constants for the path of direct attack and base catalysis, respectively.

$$k_{\text{OH}} = k_{\text{dOH}} \frac{[\text{ICl}]}{[\text{ICl}(\text{aq})]} + k_{\text{bOH}} \frac{[\text{H}_2\text{OICl}]}{[\text{ICl}(\text{aq})]} \quad (17)$$

The intercept in the linear plot (Figure 4a) indicates the existence of a base-independent or water path in the hydrolysis of  $\text{ICl}(\text{aq})$ . The rate constant for the water path is the product of the intercept and  $(K_3 + [\text{Cl}^-])/K_3$ . This gives a value of  $k_{\text{H}_2\text{O}} = 2.4 \times 10^6 \text{ s}^{-1}$  for the hydrolysis of  $\text{ICl}(\text{aq})$ . The water path of the  $\text{ICl}$  hydrolysis would proceed via the  $\text{H}_2\text{OICl}$  species, which

could either deprotonate to form  $\text{HOICl}^-$  or ionize to form  $\text{H}_2\text{OI}^+$  as shown in Scheme I. Although  $\text{H}_2\text{OI}^+$  is a stable species only in high acid,<sup>32,33</sup> it cannot be ruled out as an intermediate. Both  $\text{HOICl}^-$  and  $\text{H}_2\text{OI}^+$  would proceed rapidly to form  $\text{HOI}$ , by loss of  $\text{Cl}^-$  and  $\text{H}^+$ , respectively. We cannot distinguish between these two paths, but an  $\text{HOICl}^-$  intermediate seems more likely.

**$\text{ICl}(\text{aq})$  Hydrolysis with Buffers.** The kinetics of the hydrolysis of  $\text{ICl}(\text{aq})$  were studied with variation of buffer concentrations. Pseudo-first-order conditions, where  $[\text{buffer}] \gg [\text{ICl}_2^-]_{\text{init}}$ , were maintained for all reactions. The reactions are so fast that low concentrations of buffer ( $< 3 \times 10^{-3} \text{ M}$ ) must be used because buffers accelerate the rate.

In phosphate buffer, the observed rate constants for the hydrolysis of  $\text{ICl}_2^-$  increase from 75 000 to 150 000  $\text{s}^{-1}$  as the buffer concentration increases from 0.50 to 3.00 mM (Table III). The increase is proportional to the total concentration of phosphate,  $[\text{PO}_4]_{\text{T}}$ , rather than to the concentration of the more basic form of the buffer,  $[\text{HPO}_4^{2-}]$ . Figure 4c is a plot of the observed rate constants in 0.50 M  $[\text{Cl}^-]$  measured at different  $\text{p}[\text{H}^+]$  values versus the total concentration of phosphate. A linear regression of the data for the phosphate study gives an intercept of  $(6.1 \pm 0.2) \times 10^4 \text{ s}^{-1}$  and a slope of  $(3.0 \pm 0.2) \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ . A chloride-independent rate constant for the phosphate path of the hydrolysis of  $\text{ICl}(\text{aq})$  can be calculated by use of eq 18. This

$$k_{\text{PO}_4} = \text{slope} \times (K_3 + [\text{Cl}^-])/K_3 \quad (18)$$

gives a  $k_{\text{PO}_4}$  value of  $(1.2 \pm 0.1) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ . The intercept again means that a water path exists, and its value agrees very well with that obtained from the study of the  $\text{OH}^-$  dependence.

The observed rate constants for the hydrolysis of  $\text{ICl}_2^-$  in carbonate buffer also increase from 94 000 to 140 000  $\text{s}^{-1}$  with an increase of total carbonate concentration from 0.75 to 2.00 mM (Table III). Again, the rate constant is proportional to the total concentration of buffer,  $[\text{CO}_3]_{\text{T}}$ , rather than to the concentration of the more basic form of carbonate,  $[\text{CO}_3^{2-}]$ . Calculations show that in carbonate buffer the contribution of  $\text{OH}^-$  to the observed rate constant cannot be ignored and therefore must be taken into account. At  $\text{p}[\text{H}^+] = 9.20$  and  $9.70$  with  $\text{p}K_{\text{w}} = 13.74$  ( $\mu = 0.5$ ),<sup>34</sup> the contributions of  $\text{OH}^-$  to the observed rate constants are  $3 \times 10^3$  and  $1.1 \times 10^4 \text{ s}^{-1}$ , respectively. These values are subtracted from each of the corresponding observed rate constants for the carbonate path to give the rate constants plotted in Figure 4b. The slope of the plot for  $[\text{CO}_3]_{\text{T}}$  is  $(3.5 \pm 0.2) \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ , and the intercept is  $(6.3 \pm 0.2) \times 10^4 \text{ s}^{-1}$ . This gives a chloride-independent rate constant for the carbonate path equal to  $(1.4 \pm 0.1) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ . The intercept of the plot (Figure 4b) indicates that the water path also exists in the hydrolysis of  $\text{ICl}(\text{aq})$  in carbonate buffer, and the value is very close to the intercepts for the phosphate and hydroxide studies. These results confirm that the base-independent  $\text{ICl}$  hydrolysis (i.e., the water path) has an average rate constant of  $(6.2 \pm 0.2) \times 10^4 \text{ s}^{-1}$  in 0.50 M  $[\text{Cl}^-]$ . A chloride-independent rate constant for the water path,  $k_{\text{H}_2\text{O}}$ , is  $2.4 \times 10^6 \text{ s}^{-1}$ .

**Mechanism.** As is the case for the  $\text{OH}^-$  path, there are two possible reaction pathways for the reaction of  $\text{ICl}(\text{aq})$  with buffers. The buffers (B) could react as bases to deprotonate the hydrated  $\text{H}_2\text{OICl}$  and give  $\text{HOICl}^-$ , or they could react directly with the unhydrated  $\text{ICl}$  to form molecular complexes,  $\text{B-ICl}$ . The first pathway, in which the buffers act as proton-transfer agents, can be eliminated because  $\text{H}_2\text{PO}_4^-$ ,  $\text{HPO}_4^{2-}$ ,  $\text{HCO}_3^-$ , and  $\text{CO}_3^{2-}$  all have the same rate constant ( $\text{av} = 1.3 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ) with  $\text{ICl}(\text{aq})$ . Proton-transfer rate constants would depend on the basicity of the buffer species, and this is not the case. The indiscriminate reactivity of these species is explained by the second mechanism, where the reaction of the unhydrated form of  $\text{ICl}$  with all the buffer species occurs at diffusion-controlled rate constants ( $7 \times$

(32) Berliner, E. *J. Am. Chem. Soc.* **1956**, *78*, 3632–3637.

(33) Rao, M. D. P.; Padmanabha, J. *Indian J. Chem.* **1981**, *19A*, 1179–1182.

(34) Martell, A. E.; Smith, R. M. *Critical Stability Constants*; Plenum Press: New York, 1976; Vol. 4, p 1.


(31) Andrews, L. J.; Keefer, R. M. *Adv. Inorg. Chem. Radiochem.* **1961**, *3*, 91–131.

Table IV. Rate Constants for the Reactions of ICl(aq)<sup>a</sup>

reaction	rate const	value <sup>b</sup>
ICl(aq) + H <sub>2</sub> O	k <sub>H<sub>2</sub>O</sub>	(2.4 ± 0.2) × 10 <sup>6</sup> s <sup>-1</sup>
ICl(aq) + OH <sup>-</sup>	k <sub>OH</sub>	(4.5 ± 0.4) × 10 <sup>9</sup> M <sup>-1</sup> s <sup>-1</sup>
ICl(aq) + [PO <sub>4</sub> ] <sub>T</sub>	k <sub>PO<sub>4</sub></sub>	(1.2 ± 0.1) × 10 <sup>9</sup> M <sup>-1</sup> s <sup>-1</sup>
ICl(aq) + [CO <sub>3</sub> ] <sub>T</sub>	k <sub>CO<sub>3</sub></sub>	(1.4 ± 0.1) × 10 <sup>9</sup> M <sup>-1</sup> s <sup>-1</sup>
ICl(aq) + I <sup>-</sup>	k <sub>2</sub> (in ref 8)	(5.1 ± 0.4) × 10 <sup>8</sup> M <sup>-1</sup> s <sup>-1</sup>
H <sub>2</sub> OICl + OH <sup>-</sup>	k <sub>b OH</sub>	~4 × 10 <sup>9</sup> M <sup>-1</sup> s <sup>-1</sup>
ICl + OH <sup>-</sup>	k <sub>b OH</sub>	~7 × 10 <sup>9</sup> M <sup>-1</sup> s <sup>-1</sup>
ICl + B <sup>c</sup>	k <sub>d B</sub>	~7 × 10 <sup>9</sup> M <sup>-1</sup> s <sup>-1</sup>

<sup>a</sup> [ICl(aq)] = [ICl] + [H<sub>2</sub>OICl]. <sup>b</sup> 25.0 °C, μ = 0.50. <sup>c</sup> B = CO<sub>3</sub><sup>2-</sup>, HCO<sub>3</sub><sup>-</sup>, HPO<sub>4</sub><sup>2-</sup>, H<sub>2</sub>PO<sub>4</sub><sup>-</sup>.

10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup>) to form B-ICl (III and IV). The breakup of the B-ICl species to form HOI could involve loss of Cl<sup>-</sup> and subsequent P-O or C-O bond cleavage. Our studies only monitor the loss of ICl<sub>2</sub><sup>-</sup>, so the subsequent reactions need not be fast.



It is well-known that iodine monochloride, as a strong electron acceptor, forms 1:1 charge-transfer molecular complexes with electron-pair donors.<sup>31</sup> These electron donors include benzene and its derivatives,<sup>35</sup> amine,<sup>36</sup> pyridine,<sup>37,38</sup> and oxygen-containing organic solvents.<sup>27</sup> Numerous kinetic studies have been carried out for the reactions of ICl with alcohols,<sup>39,40</sup> aniline and substituted anilines,<sup>32,33,41,42</sup> alkenes,<sup>43-45</sup> and phenol and substituted phenols.<sup>46,47</sup> All these studies show that a 1:1 molecular complex is formed in the first step of the reaction. Some of the molecular complexes proceed to form iodination products, such as ROI in alcohols where the O-I bond is confirmed by infrared spectroscopy.<sup>39</sup> Similar reactions could occur in the hydrolysis of ICl with phosphate and carbonate buffers, where molecular complexes between ICl and phosphate or carbonate are formed.

We propose that ICl(aq) exists as two forms in rapid equilibrium, ICl and H<sub>2</sub>OICl, and that the direct attack of B on ICl is diffusion controlled. Because the average k<sub>B</sub> value is 1.3 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup> and the diffusion-controlled rate constant is 7 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup>, the concentration of unhydrated ICl in solution is estimated to be 18% of the total ICl(aq) concentration. This value gives the observed second-order rate constant of 1.3 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup>. The mechanism explains the nonselective attack by HPO<sub>4</sub><sup>2-</sup>/H<sub>2</sub>PO<sub>4</sub><sup>-</sup> and by CO<sub>3</sub><sup>2-</sup>/HCO<sub>3</sub><sup>-</sup>, because all species react with the less hydrated ICl at diffusion-controlled rates.



The free energy of hydration for the reaction of ICl(g) to give ICl(aq) is -9.26 kJ mol<sup>-1</sup>, while the corresponding ΔG<sup>o</sup> change for I<sub>2</sub> is -2.94 kJ mol<sup>-1</sup>.<sup>29</sup> If the difference between these hydration energies (-6.34 kJ mol<sup>-1</sup>) was entirely due to the specific formation of H<sub>2</sub>OICl, the equilibrium constant for [H<sub>2</sub>OICl]/[ICl] would be 13. Our experimental value of 4.6 for this equilibrium constant is reasonable because dipole-dipole interactions between ICl and

H<sub>2</sub>O will also make some contribution to its free energy of hydration.

In the case where hydroxide is the only base, the reaction must proceed by both direct attack on unhydrated ICl and deprotonation of H<sub>2</sub>OICl because the rate constant is larger than the k<sub>B</sub> values. So OH<sup>-</sup> must accelerate the reaction by a proton-transfer reaction with H<sub>2</sub>OICl to form HOICl<sup>-</sup>. The rate constants k<sub>d OH</sub> for direct attack, and k<sub>b OH</sub> for base catalysis, are defined in eq 17 and shown in Scheme I. The k<sub>d OH</sub> value should be 7 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup> for the diffusion-controlled direct reaction of OH<sup>-</sup> with ICl. The rate constant for the base-catalyzed path, k<sub>b OH</sub> = 3.9 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup>, can be calculated by use of eq 17 with k<sub>OH</sub> = 4.5 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup>, [ICl]/[ICl(aq)] = 0.18, and [H<sub>2</sub>OICl]/[ICl(aq)] = 0.82. The resolved rate constants are summarized in Table IV. Actually, k<sub>b OH</sub> is not very sensitive to changes in the ratio of [ICl]/[ICl(aq)]. The corresponding proton-transfer reactions between H<sub>2</sub>OICl and CO<sub>3</sub><sup>2-</sup> or HPO<sub>4</sub><sup>2-</sup> should be much slower because they are much weaker bases than OH<sup>-</sup>. Hence, the buffers react preferentially with ICl.


The hydrolysis of ICl(aq) (k<sub>H<sub>2</sub>O</sub> = 2.4 × 10<sup>6</sup> s<sup>-1</sup>) is many orders of magnitude faster than that of Cl<sub>2</sub> (29 s<sup>-1</sup>) and I<sub>2</sub> (3 s<sup>-1</sup>).<sup>1</sup> The huge difference is related to the polarity of the ICl molecule. Unlike Cl<sub>2</sub> and I<sub>2</sub>, ICl has a large dipole moment (1.3 D).<sup>30</sup> It is, therefore, more susceptible to attack by water and other donors. It has been noted<sup>1</sup> that in the hydrolysis of Cl<sub>2</sub>, Br<sub>2</sub>, and I<sub>2</sub> the tendency to form a ternary complex such as X<sub>2</sub>OH<sup>-</sup> increases in the order Cl<sub>2</sub> < Br<sub>2</sub> < I<sub>2</sub>. The degree of polarizability of the halogen atoms and the relative ease of valence shell expansion play important roles in the acceptance of an electron-donating group to saturate the coordination sphere. Since I<sub>2</sub> and Cl<sub>2</sub> are nonpolar species, while ICl is a highly polarized species, the formation of H<sub>2</sub>OICl as a major species in aqueous solution is reasonable and also reflects the tendency of iodine to expand its valence shell. The reactivity of H<sub>2</sub>OICl is not surprising, because it can readily lose a proton to form HOICl<sup>-</sup> or might lose Cl<sup>-</sup> to form H<sub>2</sub>OI<sup>+</sup>, and both these species will rapidly give HOI.

**Kinetics of ICl(aq) and I<sup>-</sup>.** A previous PAF study<sup>8</sup> showed that ICl<sub>2</sub><sup>-</sup> and I<sup>-</sup> react via ICl(aq) (eq 3 and 19-21) where k<sub>2</sub> is the rate-determining step and the formation of I<sub>3</sub><sup>-</sup> was observed. Our



new value for K<sub>3</sub> changes the k<sub>2</sub> value from 1.1 × 10<sup>9</sup> to 5.1 × 10<sup>8</sup> M<sup>-1</sup> s<sup>-1</sup>, as reported in Table IV. In terms of our ratio of ICl/H<sub>2</sub>OICl, we would have expected a value of 1.3 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup> for k<sub>2</sub>, based on a diffusion-controlled rate constant between ICl and I<sup>-</sup>. The reason for this discrepancy is not clear, but the ICl<sub>2</sub><sup>-</sup>/I<sup>-</sup> study was carried out at higher ionic strength, μ = 1.0. This ionic strength difference does not affect the K<sub>3</sub> value but might influence the k<sub>2</sub> rate constant or change the ratio of ICl/H<sub>2</sub>OICl.

Since we now know that the hydrolysis rate of ICl(aq) is extremely fast, we need to consider why this path was not seen in the study with I<sup>-</sup>.<sup>8</sup> There are several reasons: First, the equilibrium in eq 4 in 1 M [H<sup>+</sup>] concentrations gives only 1% HOI at the lowest Cl<sup>-</sup> concentration used (0.1 M) and in 0.1 M [H<sup>+</sup>] only 10% HOI when [Cl<sup>-</sup>] is 0.1 M. Second, the rate constant for the reaction between HOI and I<sup>-</sup> (eq 22) is only 3.2 × 10<sup>5</sup>



M<sup>-1</sup> s<sup>-1</sup> and the pathway between H<sub>2</sub>OI<sup>+</sup> and I<sup>-</sup> does not contribute.<sup>48</sup> (This suggests that H<sub>2</sub>OI<sup>+</sup> should be written as H<sub>2</sub>OIOH<sub>2</sub><sup>+</sup> in order to account for its lower reactivity.) Thus, even if 10% of the ICl<sub>2</sub><sup>-</sup> formed HOI, the rate of production of I<sub>3</sub><sup>-</sup> via HOI would be negligible compared to the formation via

(35) Keefer, R. M.; Andrews, L. J. *J. Am. Chem. Soc.* **1950**, *72*, 5170-5173.

(36) Hassel, O.; Hope, H. *Acta Chem. Scand.* **1960**, *2*, 391-397.

(37) Gupta, A. D.; Basu, R. *J. Chim. Phys. Phys.-Chim. Biol.* **1975**, *72*, 271.

(38) Schuster, I. I.; Roberts, J. D. *J. Org. Chem.* **1979**, *44*, 2658-2662.

(39) Das, C. R.; Bose, A. N. *Bull. Chem. Soc. Jpn.* **1973**, *46*, 818-821.

(40) Terentyev, A. P.; Belenkey, L. I.; Yanovkaya, L. A. *J. Gen. Chem. USSR Engl. Transl.* **1954**, *24*, 1251-1255.

(41) Vainstein, F. M.; Tomilenko, E. I.; Shilov, E. A. *Kinet. Catal. Engl. Transl.* **1966**, *7*, 26-34.

(42) Berliner, E. *J. Am. Chem. Soc.* **1950**, *72*, 4003-4009.

(43) Schmid, G. H.; Gordon, J. W. *Can. J. Chem.* **1986**, *64*, 2171-2174.

(44) Schmid, G. H.; Gordon, J. W. *Can. J. Chem.* **1984**, *62*, 2526-2534.

(45) Amirtha, N.; Viswanathan, S.; Ganesan, R. *Bull. Chem. Soc. Jpn.* **1983**, *56*, 314-317.

(46) Shashidhar, G. V. S.; Satyanatayana, N.; Sundaram, E. *Indian J. Chem.* **1986**, *25A*, 289-291.

(47) Rao, M. D. P.; Padmanabha, J. *Indian J. Chem.* **1981**, *20A*, 133-135.

(48) Palmer, D. A.; Eldik, van R. *Inorg. Chem.* **1986**, *25*, 928-931.

ICl(aq) and I<sup>-</sup>. Hence, the absence of a hydrolysis path in the previous study<sup>8</sup> is verified.

### Conclusion

The hydrolysis of iodine monochloride in water is 5-6 orders of magnitude faster than that of Cl<sub>2</sub> and I<sub>2</sub>. The difference is attributed to the dipole moment and polarizability of ICl and to the ease of valence shell expansion of iodine. This leads to the formation of ICl molecular complexes with H<sub>2</sub>O, OH<sup>-</sup>, and buffers.

We propose that both ICl and H<sub>2</sub>OICl exist in aqueous solution and that the reaction with ICl to form molecular complexes is diffusion controlled. Hydroxide ion has a dual role in the hydrolysis of ICl(aq). It can directly attack ICl to give HOICl<sup>-</sup>, and it can deprotonate H<sub>2</sub>OICl to give HOICl<sup>-</sup>. The HOICl<sup>-</sup> intermediate rapidly loses Cl<sup>-</sup> to give HOI.

**Acknowledgment.** This work was supported by National Science Foundation Grants CHE-8616666 and CHE-8720318.

## Organo-f-Element Thermochemistry. Absolute Metal-Ligand Bond Disruption Enthalpies in Bis(pentamethylcyclopentadienyl)samarium Hydrocarbyl, Hydride, Dialkylamide, Alkoxide, Halide, Thiolate, and Phosphide Complexes. Implications for Organolanthanide Bonding and Reactivity

Steven P. Nolan, David Stern, and Tobin J. Marks\*


Contribution from the Department of Chemistry, Northwestern University, Evanston, Illinois 60201-3113. Received February 8, 1989

**Abstract:** Absolute samarium-ligand bond disruption enthalpies in the series Cp'<sub>2</sub>Sm-R/X (Cp' = η<sup>5</sup>-(CH<sub>3</sub>)<sub>5</sub>C<sub>5</sub>) have been measured by iodolytic and alcoholic isoperibol titration calorimetry of Cp'<sub>2</sub>Sm/(Cp'<sub>2</sub>Sm-I)<sub>n</sub>, (Cp'<sub>2</sub>Sm-O<sup>t</sup>Bu)<sub>2</sub>/Cp'<sub>2</sub>Sm-R/X ensembles in toluene. Derived D(Cp'<sub>2</sub>Sm-R/X) values in toluene solution are as follows (kcal/mol, R/X): 47.0 (1.5), CH(SiMe<sub>3</sub>)<sub>2</sub>; 45.0 (1.5), η<sup>3</sup>-C<sub>3</sub>H<sub>5</sub>; 93.2, CCPh; 54.2 (3.0), H; 48.2 (1.8), NMe<sub>2</sub>; 82.4 (3.5), O<sup>t</sup>Bu; 81.3 (1.0), OCH<sup>t</sup>Bu; 97.1 (3.0), Cl; 83.6 (1.5), Br; 69.4 (2.4), I; 73.4 (2.4), S<sup>n</sup>Pr; 32.6 (2.0), PEt<sub>2</sub>. D(Cp'<sub>2</sub>Sm-THF) and D(Cp'<sub>2</sub>Sm(THF)-THF) values were also determined in toluene and are 7.3 (0.4) and 4.9 (1.0) kcal/mol, respectively, while D(Cp'<sub>2</sub>Sm(THF)-I) was found to be 72.7 (2.9) kcal/mol. The observed D(Sm-halogen) parameters are close to the D<sub>1</sub> values of the corresponding samarium trihalides. Important trends in D(Cp'<sub>2</sub>Sm-R/X) include a relatively small value of D(Sm-H) - D(Sm-alkyl), a large value of D(Sm-I) - D(Sm-alkyl), and generally strong bonds to group 15 and group 16 ligands. A variety of Sm-centered ligand transposition and oxidative addition/reductive elimination processes are analyzed in light of the present data. The formation of strong Sm-heteroelement bonds makes an important contribution to the driving force. Hydrocarbon functionalization via dinuclear Sm(II) → Sm(III) oxidative addition processes is only expected to be exothermic in special cases.

Recent developments in organometallic thermochemistry have demonstrated the considerable insight into bonding and reaction chemistry that metal-ligand bond disruption enthalpy (eq 1 for

$$D(L_nM-R) = \Delta H_f^\circ(L_nM) + \Delta H_f^\circ(R^\bullet) - \Delta H_f^\circ(L_nM-R) \quad (1)$$

$L_n$  = ancillary ligands



the adiabatic, homolytic process of eq 2) information can provide.<sup>1,2</sup> In the solution phase, relative metal-ligand bond enthalpies have been obtained by equilibration or various calorimetric techniques<sup>1,2</sup> and have shed considerable light on bonding patterns and the

driving forces for important reactions. Absolute metal-ligand bond enthalpies provide, in addition, an absolute quantitation of the energetics of eq 2 and a means of comparing bond enthalpies for different M and L<sub>n</sub>M centers. Relative metal-ligand enthalpies can be placed on an approximately absolute scale by judicious choice of a reference ("anchor") point, such as the corresponding D<sub>1</sub>(MX<sub>n</sub>) datum for an L<sub>n</sub>MR/L<sub>n</sub>MX series.<sup>3,4</sup> However, this approximation assumes aspects of transferability that may not always be general and certainly warrants further scrutiny. Direct measurements of absolute metal-ligand bond enthalpies have included kinetic,<sup>2c,h,5</sup> photoacoustic,<sup>2b</sup> laser pyrolytic,<sup>7</sup> two-electron

(1) Marks, T. J., Ed. Metal-Ligand Bonding Energetics in Organotransition Metal Compounds. *Polyhedron Symposium-in-Print* 1988, 7.

(2) For some leading references, see also: (a) Pilcher, G.; Skinner, H. A. In *The Chemistry of the Metal-Carbon Bond*; Harley, F. R., Patal, S., Eds.; Wiley: New York, 1982; pp 43-90. (b) Connor, J. A. *Top. Curr. Chem.* 1977, 71, 71-110. (c) Halpern, J. *Acc. Chem. Res.* 1982, 15, 238-244. (d) Mondal, J. U.; Blake, D. M. *Coord. Chem. Rev.* 1983, 47, 204-238. (e) Mansson, M. *Pure Appl. Chem.* 1983, 55, 417-426. (f) Skinner, H. A.; Connor, J. A. *Pure Appl. Chem.* 1985, 57, 79-88. (g) Pearson, R. G. *Chem. Rev.* 1985, 85, 41-59. (h) Halpern, J. *Inorg. Chim. Acta* 1985, 100, 41-48. (i) Bryndza, H. E.; Fong, L. K.; Pacjello, R. A.; Tam, W.; Bercaw, J. E. *J. Am. Chem. Soc.* 1987, 109, 1444-1456. (j) Nolan, S. P.; Hoff, C. D.; Stoutland, P. O.; Newman, L. J.; Buchanan, J. M.; Bergman, R. G.; Yang, G. K.; Peters, K. G. *J. Am. Chem. Soc.* 1987, 109, 3143-3145, and references therein. (k) Dias, A. R.; Salema, M. S.; Martinho-Simoes, J. A. *J. Organomet. Chem.* 1981, 222, 69-78.

(3) (a) Schock, L. E.; Marks, T. J. *J. Am. Chem. Soc.* 1988, 110, 7701-7715. (b) Dias, A. R.; Simoes, J. A. M. In ref 1, pp 1531-1544. (c) Lappert, M. F.; Patil, D.; Pedley, J. B. *J. Chem. Soc., Chem. Commun.* 1975, 830-831.

(4) (a) Bruno, J. W.; Marks, T. J.; Morss, L. R. *J. Am. Chem. Soc.* 1983, 105, 6824-6832. (b) Sonnenberger, D. C.; Morss, L. R.; Marks, T. J. *Organometallics* 1985, 4, 352-355. (c) Bruno, J. W.; Stecher, H. A.; Morss, L. R.; Sonnenberger, D. C.; Marks, T. J. *J. Am. Chem. Soc.* 1986, 108, 7275-7280, and ref 4. (d) Lin, Z. Ph.D. Thesis, Northwestern University, Evanston, IL, December 1988, Appendix A (D(Th-S<sup>n</sup>Pr) = 105 (5) kcal/mol).

(5) (a) Halpern, J. In ref 1, pp 1483-1490. (b) Hay, B. P.; Finke, R. G. In ref 1, pp 1469-1481. (c) Koenig, T. W.; Hay, B. P.; Finke, R. G. In ref 1, pp 1499-1516. (d) Wayland, B. B. In ref 1, pp 1545-1555. (e) Geno, M. K.; Halpern, J. *J. Chem. Soc., Chem. Commun.* 1987, 1052-1053.

(6) (a) Yang, G. K.; Valda, V.; Peters, K. S. In ref 1, pp 1619-1622. (b) Rothberg, L. J.; Simon, J. D.; Bernstein, M.; Peters, K. S. *J. Am. Chem. Soc.* 1983, 105, 3464-3470.