

Inhibition of Chorismate Synthase by (6*R*)- and (6*S*)-6-Fluoro-5-enolpyruvylshikimate 3-Phosphate

Shankar Balasubramanian,[†] Gareth M. Davies,[‡]
John R. Coggins,[§] and Chris Abell^{*†}

University Chemical Laboratory, University of Cambridge
Lensfield Road, Cambridge CB2 1EW, England
Department of Biochemistry, University of Glasgow
Glasgow G12 8QQ, Scotland
ICI Pharmaceuticals, Alderley Park
Macclesfield, Cheshire SK10 4TG, England


Received May 28, 1991

The seventh step of the shikimate pathway,¹ mediated by chorismate synthase (EC 4.6.1.4), is the conversion of 5-enolpyruvylshikimate 3-phosphate (1, EPSP) to chorismate (2) (Scheme I). This unusual enzymatic transformation proceeds by an overall trans-1,4-elimination of phosphate with abstraction of the C-6 *pro-R* hydrogen.^{2,3} The mechanism of this reaction is unknown. Experiments with model systems⁴ and arguments based on molecular orbital considerations⁵ have been used to discount a concerted E2' elimination. Some of the other postulated mechanisms are summarized in Scheme I.^{1-3,6} The rearrangement mechanism¹ proceeding via 3 is unlikely as this compound is a competitive inhibitor but not a substrate for the enzyme.⁷ It has also been shown that phosphate loss is not a fast step prior to the rate-determining step of the reaction.⁶ More recently we have shown that the reaction proceeds with an associated primary kinetic isotope effect at C-6 on *V* and *V*/*K*.⁸


Modification of the reactivity of EPSP by stereospecific substitution of hydrogen by fluorine at C-6 gives compounds which could potentially discriminate between the mechanisms in Scheme I. These 6-fluoro-EPSPs (7b and 7c) could act as substrates, suicide inhibitors, or competitive inhibitors of chorismate synthase. We report the synthesis of (6*R*)-6-fluoro-EPSP (7b) and (6*S*)-6-fluoro-EPSP (7c) and their interaction with chorismate synthase.

6-Fluoro-EPSPs were synthesized from the corresponding 6-fluoroshikimates (5b and 5c)^{9,10} following our protocol for the enzymatic transformation of shikimate (5a) to EPSP (7a) via shikimate 3-phosphate (6a).^{10,11} The transformations were performed sequentially in deuteriated buffer using shikimate kinase and EPSP synthase, each isolated from overexpressing strains of *Escherichia coli* (Scheme II).^{12,13} The reaction was monitored

Scheme I. Postulated Mechanisms for the Chorismate Synthase Catalyzed Reaction



Scheme II^a



^a (i) Shikimate kinase, 1.8 units, 24 h. (ii) EPSP synthase, 0.22 units, 24 h. Transformations i and ii were carried out sequentially on 50 mM (6*R*)- and (6*S*)-6-fluoroshikimate in 1.0 mL of D₂O, pD 7.1 (Tris-DCI, 300 mM), at 25 °C, containing 50 mM MgCl₂, 50 mM ATP, and 50 mM phosphoenol pyruvate, and were followed by ¹H NMR spectroscopy.

by ¹H NMR spectroscopy in which the C-2 vinyl proton resonance of each compound was clearly visible. Both 6-fluoroshikimates 5b and 5c are good substrates for shikimate kinase and were transformed at rates comparable to that of shikimate under the same conditions. (6*R*)-6-Fluoroshikimate 3-phosphate (6b) was produced quantitatively, and (6*S*)-6-fluoroshikimate 3-phosphate (6c) was produced in greater than 85% yield (by ¹H NMR spectroscopy). On addition of EPSP synthase, 6b and 6c were each transformed at a rate which was about an order of magnitude slower than that of shikimate 3-phosphate under the same conditions. (6*R*)-6-Fluoro-EPSP (7b) was produced quantitatively, and (6*S*)-6-fluoro-EPSP (7c) was produced in approximately 85% yield. The final reaction mixtures were treated with apyrase (Sigma, grade VII) to degrade ATP and ADP, which facilitated purification of 7b and 7c by ion-exchange chromatography on Dowex 1X8.^{11,14} Both purified analogues were isolated as the dibarium salts in 30-40% overall yield.

To explore the possibility that either 7c or, more likely, 7b is a substrate for chorismate synthase, each was incubated with the purified *Neurospora crassa* enzyme under *V*_{max} conditions.¹⁵ UV spectroscopy was used to detect the appearance of a diene chromophore by monitoring changes in absorbance in the region 240-300 nm.¹⁶ No diene formation was detected for either

* Address correspondence to Dr. C. Abell, University Chemical Laboratory, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, U.K. Telephone +44-223-336405; FAX +44-223-336362.

[†] University of Cambridge.

[‡] ICI Pharmaceuticals.

[§] University of Glasgow.

(1) Ganem, B. *Tetrahedron* 1978, 34, 3353-3383.

(2) Hill, R. K.; Newkome, G. R. *J. Am. Chem. Soc.* 1969, 91, 5893-5894.

(3) Onderka, D. K.; Floss, H. G. *J. Am. Chem. Soc.* 1969, 91, 5894-5896.

Onderka, D. K.; Floss, H. G.; Carroll, M. *J. Biol. Chem.* 1972, 247, 736-744.

(4) Hill, R. K.; Bock, M. G. *J. Am. Chem. Soc.* 1978, 100, 637-639.

Toromanoff, E. C. *R. Sceances Acad. Sci., Ser. C* 1980, 290, 81-84.

(5) Fukui, K. *Tetrahedron Lett.* 1965, 2427-2432. Anh, N. T. *J. Chem. Soc., Chem. Commun.* 1968, 1089-1090.

(6) Hawkes, T. R.; Lewis, T.; Coggins, J. R.; Mousdale, D. M.; Lowe, D. L.; Thorneley, R. N. F. *Biochem. J.* 1990, 265, 899-902.

(7) Bartlett, P. A.; Maitra, U.; Chouinard, P. M. *J. Am. Chem. Soc.* 1986, 108, 8068-8071.

(8) Balasubramanian, S.; Abell, C.; Coggins, J. R. *J. Am. Chem. Soc.* 1990, 112, 8581-8583.

(9) Sutherland, J. K.; Watkins, W. J.; Bailey, J. P.; Chapman, A. K.; Davies, G. M. *J. Chem. Soc., Chem. Commun.* 1989, 1386-1387. The full experimental details have been published in a thesis¹¹ (Watkins, W. J. Ph.D. Thesis, University of Manchester, 1987) and a patent (European Patent Application Number 393923, published October 1990).

(10) The experimental procedure for the conversion of fluoroshikimates to fluoro-EPSPs and the spectroscopic characterization of the fluoroshikimates and fluoro-EPSPs are in the supplementary material accompanying this paper.

(11) Balasubramanian, S.; Abell, C. *Tetrahedron Lett.* 1991, 32, 963-966.

(12) Millar, G.; Lewendon, A.; Hunter, M. G.; Coggins, J. R. *Biochem. J.* 1986, 237, 427-437.

(13) Duncan, K.; Lewendon, A.; Coggins, J. R. *FEBS Lett.* 1984, 170, 59-63.

(14) Knowles, P. F.; Levin, J. G.; Sprinson, D. B. *Methods Enzymol.* 1970, 17, 360-362.

(15) White, P. J.; Millar, G.; Coggins, J. R. *Biochem. J.* 1988, 251, 313-322.




Figure 1. Cornish-Bowden plot showing competitive inhibition of chorismate synthase by (6*S*)-6-fluoro-EPSP (**7c**). UV assays (275 nm) were carried out at 25 °C, pH 7.0 (triethanolamine hydrochloride, 50 mM), and included 1.2 millimoles of chorismate synthase, 20 μM NADPH, 10 μM FMN, 50 mM KCl, and 2.5 mM MgCl₂ in addition to substrate EPSP (**7a**) and inhibitor (6*S*)-6-fluoro-EPSP (**7c**) in a final volume of 1 mL. Inhibitor concentrations were 0, 0.5, 1, 2, 3.5, and 5 μM, and substrate concentrations were (■) 20 μM EPSP, (▲) 35 μM EPSP, and (○) 50 μM EPSP.

compound under conditions which would have easily detected a turnover rate 0.2% that of EPSP itself.¹⁷

Competition experiments were performed in which chorismate synthase was assayed at various fixed concentrations of EPSP in the presence of a range of concentrations of **7b** or **7c**. Figure 1 shows a Cornish-Bowden plot¹⁸ of the data obtained for (6*S*)-6-fluoro-EPSP (**7c**). The parallel plots clearly signify a competitive mode of inhibition. The inhibition constant K_i was determined from a Dixon plot.¹⁹ It is found that both fluoro-EPSPs show clean competitive inhibition with **7c** having an affinity an order of magnitude greater than **7b**: K_i ((6*S*)-6-fluoro-EPSP) = 0.2 ± 0.1 μM, K_i ((6*R*)-6-fluoro-EPSP) = 3.0 ± 0.3 μM. These values compare with K_i (iso-EPSP **3**) = 8.7 μM,⁷ and K_m (EPSP) = 2.2 μM.⁸ The lack of irreversible inhibition by either compound was confirmed by incubation of *N. crassa* chorismate synthase with 50 μM of each inhibitor at 25 °C. Over a period 1 h, no loss of enzyme activity was observed relative to a control which lacked inhibitor.

The lack of irreversible inhibition is inconsistent with a mechanism involving a covalent enzyme-intermediate adduct such as **4**. While the observation that both 6-fluoro-EPSPs are potent competitive inhibitors does not itself support or preclude any of the other mechanisms in Scheme I, it does provide a useful tool for future mechanistic studies of the enzyme.

Acknowledgment. We thank B. Wright and D. Jude for assistance in preliminary studies, Dr. S. Crossland for assistance with mass spectrometry, and ICI Agrochemicals and the SERC for a case studentship for S.B.

Supplementary Material Available: Preparation of **7b,c** from **5b,c**, spectroscopic characterization of **5b,c** and **7b,c**, and Dixon plot showing inhibition of chorismate synthase by **7c** (4 pages). Ordering information is given on any current masthead page.

(16) Chorismate formation is normally monitored by the appearance of the diene chromophore which has its λ_{max} at 275 nm. In these experiments, absorbance was monitored in the range 240–300 nm in order to accommodate a possible shift in the absorbance maximum due to a fluorine substituent.

(17) This experiment does not rigorously preclude either compound being a substrate but puts an upper limit on their turnover rate.

(18) Cornish-Bowden, A. *Biochem. J.* 1974, 137, 143–144.

(19) Dixon, M. *Biochem. J.* 1953, 55, 170–171.

Modulation of Physical and Chemical Properties of η -H₂ Complexes of Osmium Amines by Facile Substitution

Zai-Wei Li and Henry Taube*

Department of Chemistry, Stanford University
Stanford, California 94305

Received May 28, 1991

Since the discovery of the first dihydrogen complex by Kubas et al.,¹ many dihydrogen complexes have been synthesized. In 1971,² the preparation in our laboratory of [Os(en)₂H₂]²⁺ as the chloride salt was reported. It was described as a dihydride and was assigned a *cis* configuration on the strength of ¹H NMR results which revealed two sets of amine protons in equal number. Our investigation of the analogous species [Os(NH₃)₄H₂]²⁺, not heretofore reported, throws new light on that structural assignment and, as well, provides ready access to a series of complexes arrived at by the simple addition of a variety of ligands to these 16e⁻ moieties.

When [Os(NH₃)₄H₂](B(C₆H₅)₄)₂³ (**1**) is dissolved in (C-D₃)₂CO, the ¹H NMR spectrum reveals only two kinds of protons ascribable to the cation, in the abundance ratio 6:1 at δ = 3.82 ppm and -11.37 ppm, respectively. For the purposes of species differentiation, the value of T_1 for the coordinated hydrogen was also measured (T_1 = 572 ms, 20 °C), as it was for the other species to be dealt with. When a trace of acid, for example, HO₃SCF₃, is present, slow H/D exchange between the solvent and coordinated hydrogen ensues, and, in a partially exchanged sample, J_{HD} was measured as 4.0 Hz. When any of a large number of solutes is added in excess, among them acetonitrile (AN), pyridine (Py), imidazole (Im), I⁻, Cl⁻, D₂O, and Br⁻, δ (ppm) J_{HD} (Hz), and T_1 (ms, 400 MHz) change and new characteristic values are registered. (See Table I.) In every case except with D₂O and (C-D₃)₂CO as addend, the corresponding solid salt was also prepared,⁵ and dissolved, with no discernible differences in the ¹H NMR signals. Because the solute level is low (0.010 M), we can conclude, at least in the case of the labile systems I⁻, Br⁻, or Cl⁻ as addend, that in acetone the affinity of the osmium center for the ligand is very high. As expected, it is much reduced in D₂O as solvent.

Of potential anionic ligands, the only one among those we have introduced which does not change the values of δ , J_{HD} , and T_1 is PF₆⁻ (even CF₃SO₃⁻ produces a set of characteristic values). This indicates that neither B(C₆H₅)₄⁻ nor PF₆⁻ enters the coordination sphere of the osmium complex, a supposition which, in the case of the former at least, is reasonable and, in view of the bulk and almost spherical shape of PF₆⁻, is reasonable for it also. However, it leaves open the question of whether (CD₃)₂CO also is a ligand when the B(C₆H₅)₄⁻ and PF₆⁻ salts are dissolved. That this is in fact the case is indicated by observations made for [Os(en)₂H₂]²⁺, and it is therefore likely also the case for [Os(NH₃)₄H₂]²⁺.

In preparing [Os(en)₂H₂]²⁺, we followed the literature procedure,² but with the difference that, instead of Cl⁻, B(C₆H₅)₄⁻ was

(1) Kubas, G. J.; Ryan, R. R.; Swanson, B. I.; Vergamini, P. J.; Wasserman, J. J. *J. Am. Chem. Soc.* 1984, 106, 451–452.

(2) Malin, J.; Taube, H. *Inorg. Chem.* 1971, 10, 2403.

(3) [Os(NH₃)₄H₂](B(C₆H₅)₄)₂ was made by the following procedure: Os(NH₃)₄(O₃SCF₃)₃ (100 mg) in 15 mL of H₂O was reduced by Zn/Hg (3 g) for 3 h, and then 15 mL of 0.2 M NaB(C₆H₅)₄ solution was added. The resulting precipitate was dried under vacuum. Microanal. Calcd for [Os(NH₃)₄H₂](B(C₆H₅)₄)₂·2H₂O: C, 61.67; H, 6.25; N, 5.99. Found: C, 61.50; H, 6.20; N, 5.80. Yield: >70%.

(4) Li, Z.-W.; Harman, W. D.; Lay, P. A.; Taube, H. *Inorg. Chem.*, submitted.

(5) The preparation of the pyridine adduct is typical of the others. The compound [Os(NH₃)₄(H₂)·Py][B(C₆H₅)₄]₂ (**2**) was prepared by dissolving **1** (100 mg) in pyridine (5 mL); after 1 h, ether was added to cause precipitation. The precipitate was collected, washed with ether, and dried. Yield: 90%. Microanal. Calcd for [Os(NH₃)₄(H₂)·Py][B(C₆H₅)₄]₂·2H₂O: C, 62.81; H, 5.82; N, 6.91. Found: C, 62.76; H, 6.03; N, 6.64. ¹H NMR in (CD₃)₂CO (ppm): 8.83 (d, 2 H, Py), 8.14 (t, 1 H, Py), 7.75 (t, 2 H, Py), 7.40–6.70 (m, 40 H, C₆H₅), 3.74 (s, br, 12 H, 4 NH₃), -7.44 (s, 2 H, OsH₂).