

Hyperconjugative Stabilization of Silicenium Ions: Kinetics of Hydride Abstractions from β -Element Substituted Silanes¹

Nils Basso,[†] Solvig Görs,[‡] Eckhard Popowski,[‡] and Herbert Mayr^{*,§}

Contribution from the Institut für Chemie, Medizinische Universität zu Lübeck, D-23562 Lübeck, Germany, and the Fachbereich Chemie, Universität Rostock, D 18055 Rostock, Germany

Received November 30, 1992. Revised Manuscript Received March 1, 1993

Abstract: Rates of hydride abstractions from β -silyl-, β -germyl-, and β -stannyl-substituted silanes [$\text{HSiMe}_2(\text{CH}_2\text{-MMe}_3)$, M = Si, Ge, Sn] and related compounds by diarylcarbenium ions have been measured. The intermediate silicenium ions, produced in the rate-determining step, are stabilized by hyperconjugation, but the effects are much smaller than in comparable carbenium ions.

Introduction

The stabilization of carbenium ions by β -silyl, β -germyl, and β -stannyl groups has been derived from solvolysis rates of β -element-substituted alkyl derivatives,¹ rates of electrophilic addition to allyl-² and vinylenes³ compounds, and rates of hydride abstraction from element-substituted alkanes.⁴⁻⁶

Traylor investigated hydride abstractions from tetraethylelement compounds by the trityl cation.^{4b} The rate was found to strongly depend on the nature of the metal, indicating increasing hyperconjugative stabilization of carbenium ions in the order Si-C < Ge-C < Sn-C < Pb-C (Table I). Similar effects have been discussed for the interaction between $\sigma(\text{Si-CH}_2)$ and the singly-occupied p-orbital at germanium in germly radicals $\text{Me}_3\text{-SiCH}_2(\text{Me}_2)\text{Ge}^{\cdot}$.⁷ The comparison with the rate of hydride abstraction in cycloheptatriene shows that the stabilization of

Table I. Rate Constants for the Hydride Transfer from the Organometallic Compounds Et_4M (M = Si, Ge, Sn, Pb) to $\text{Ph}_3\text{C}^+\text{BF}_4^-$ (H_3CCN , +29.8 °C)^{4b}

compd	$k_2, \text{L}\cdot\text{mol}^{-1}\cdot\text{s}^{-1}$	compd	$k_2, \text{L}\cdot\text{mol}^{-1}\cdot\text{s}^{-1}$
Et_4Si	5.7×10^{-8}	Et_4Pb	4.9×10^{-1}
Et_4Ge	4.0×10^{-6}	cycloheptatriene	2.1×10^{-1}
Et_4Sn	8.3×10^{-4}		

^a k_2 per hydrogen.

Table II. Rate Constants for the Reaction of $\text{Ph}_3\text{C}^+\text{BF}_4^-$ with 1,3-Dimetallopropanes $\text{Me}_3\text{M-CH}_2\text{CH}_2\text{CH}_2\text{-M}'\text{Me}_3$ (M, M' = C, Si, Ge, Sn, Pb) in H_3CCN at +30 °C^{4a}

compd	$k_2, \text{L}\cdot\text{mol}^{-1}\cdot\text{s}^{-1}$
$\text{Me}_3\text{Si-CH}_2\text{CH}_2\text{-SiMe}_3$	$6.2 \times 10^{-6} \text{ }^b$
$\text{Me}_3\text{Si-CH}_2\text{CH}_2\text{-GeMe}_3$	1.9×10^{-5}
$\text{Me}_3\text{Si-CH}_2\text{CH}_2\text{-SnMe}_3$	5.9×10^{-4}
$\text{Me}_3\text{Si-CH}_2\text{CH}_2\text{-PbMe}_3$	1.3×10^{-1}
$\text{Me}_3\text{Ge-CH}_2\text{CH}_2\text{-GeMe}_3$	1.8×10^{-4}
$\text{Me}_3\text{Ge-CH}_2\text{CH}_2\text{-SnMe}_3$	7.1×10^{-3}
$\text{Me}_3\text{Ge-CH}_2\text{CH}_2\text{-PbMe}_3$	2×10^{-1}
$\text{Me}_3\text{Sn-CH}_2\text{CH}_2\text{-CMe}_3$	$2 \times 10^{-4} \text{ }^b$
$\text{Me}_3\text{Sn-CH}_2\text{CH}_2\text{-SnMe}_3$	4×10^{-2}

^a k_2 per hydrogen. ^b Solvent: 50% CD_3CN in CD_2Cl_2 .

the β -lead-substituted carbenium ion $\text{Et}_3\text{Pb-CH}_2\text{CH}_2^+$ is comparable to that of the tropylium ion ($\text{p}K_{\text{R}^+} = +4.76^8$).

Disubstituted alkanes $\text{Me}_3\text{M-CH}_2\text{CH}_2\text{CH}_2\text{-M}'\text{Me}_3$ (1,3-dimetallopropanes, M or M' = C, Si, Ge, Sn, Pb) react considerably faster with Ph_3C^+ (Table II). From the magnitude of the substituent effects it was concluded that bridging of the metal is not important in the transition state.

Recently, we reported rates of hydride abstractions from silanes and derived the effect of various substituents for stabilizing silicenium ions.^{9a} In agreement with related experimental investigations^{9b,c} and theoretical predictions,⁶ π -conjugation has been found to be much less effective for silicenium ions than for carbenium ions.^{9a} Comparison of the hydride abstractions from silanes, germanes, and stannanes showed the decreasing importance of π -conjugation with increasing size of the central atom.^{9d}

(8) Ritchie, C. D. *Can. J. Chem.* 1986, 64, 2239.

(9) (a) Mayr, H.; Basso, N.; Hagen, G. *J. Am. Chem. Soc.* 1992, 114, 3060. (b) Chojnowski, J.; Fortuniak, W.; Stanczyk, W. *J. Am. Chem. Soc.* 1987, 109, 7776. (c) Chojnowski, J.; Wilczek, L.; Fortuniak, W. *J. Organomet. Chem.* 1977, 135, 13. (d) Mayr, H.; Basso, N. *Angew. Chem.* 1992, 104, 1103; *Angew. Chem., Int. Ed. Engl.* 1992, 31, 1046. (e) Mayr, H.; Schneider, R.; Schade, C.; Bartl, J.; Bederke, R. *J. Am. Chem. Soc.* 1990, 112, 4446.

[†] Dedicated to Professor R. P. Kreher on the occasion of his 60th birthday.
[†] Medizinische Universität zu Lübeck.

[‡] Universität Rostock.

[§] New address: Institut für Organische Chemie, Technische Hochschule Darmstadt, D-64287 Darmstadt, Germany.

(1) (a) Sommer, L. H.; Baughman, G. A. *J. Am. Chem. Soc.* 1961, 83, 3346. (b) Cook, M. A.; Eaborn, C.; Walton, D. R. M. *J. Organomet. Chem.* 1970, 24, 301. (c) Lambert, J. B.; Wang, G.-t.; Finzel, R. B.; Teramura, D. H. *J. Am. Chem. Soc.* 1987, 109, 7838. (d) Lambert, J. B.; Wang, G.-t. *Tetrahedron Lett.* 1988, 29, 2551. (e) Lambert, J. B.; Wang, G.-t.; Teramura, D. H. *J. Org. Chem.* 1988, 53, 5422. (f) Lambert, J. B.; Wang, G.-t. *J. Phys. Org. Chem.* 1988, 1, 169. (g) Lambert, J. B.; Chelius, E. C. *J. Am. Chem. Soc.* 1990, 112, 8120.

(2) (a) Hagen, G.; Mayr, H. *J. Am. Chem. Soc.* 1991, 113, 4954. (b) Mayr, H.; Hagen, G. *J. Chem. Soc., Chem. Commun.* 1989, 91. (c) Mayr, H.; Pock, R. *Tetrahedron* 1986, 42, 4211. (d) Bartl, J.; Steenken, S.; Mayr, H. *J. Am. Chem. Soc.* 1991, 113, 7710.

(3) (a) Soderquist, J. A.; Hassner, A. *Tetrahedron Lett.* 1988, 29, 1899. (b) Cochran, J. C.; Williams, L. E.; Bronk, B. S.; Calhoun, J. A.; Fassberg, J.; Clark, K. G. *Organometallics* 1989, 8, 804. (c) Hagen, G. Ph.D. Thesis, Medizinische Universität Lübeck, 1990.


(4) Solution studies: (a) Traylor, T. G.; Koerner, G. S. *J. Org. Chem.* 1981, 46, 3651. (b) Jerkunica, J. M.; Traylor, T. G. *J. Am. Chem. Soc.* 1971, 93, 6278. See also: (c) Davis, D. D.; Jacocks, H. M., III. *J. Organomet. Chem.* 1981, 206, 33.

(5) Gas-phase studies of R_3Si^+ : (a) Hajdasz, D.; Squires, R. *J. Chem. Soc., Chem. Commun.* 1988, 1212. (b) Li, X.; Stone, J. A. *J. Am. Chem. Soc.* 1989, 111, 5586. (c) Ciommer, B.; Schwarz, H. *J. Organomet. Chem.* 1983, 244, 319. (d) Drewello, T.; Burgers, P. C.; Zummack, W.; Apeloig, Y.; Schwarz, H. *Organometallics* 1990, 9, 1161.



(6) Theoretical investigations of R_3Si^+ : (a) Wierschke, S. G.; Chandrasekhar, J.; Jorgensen, W. L. *J. Am. Chem. Soc.* 1985, 107, 1496. (b) White, J. C.; Cave, R. J.; Davidson, E. R. *J. Am. Chem. Soc.* 1988, 110, 6308. (c) Ibrahim, M. R.; Jorgensen, W. L. *J. Am. Chem. Soc.* 1989, 111, 819. (d) Eaborn, C. *J. Chem. Soc., Chem. Commun.* 1972, 1255. (e) Lambert, J. B. *Tetrahedron* 1990, 46, 2677. (f) Apeloig, Y. In *The Chemistry of Organic Silicon Compounds*; Patai, S., Rappoport, Z., Eds.; John Wiley & Sons: Chichester, 1989; Part 1, p 196. (g) Apeloig, Y. In *Heteroatom Chemistry*; Block, E., Ed.; VCH-Verlagsgesellschaft: Weinheim 1990; p 27.

(7) Mochida, K. *Bull. Chem. Soc. Jpn.* 1984, 57, 796.

Scheme I



Analogously, hyperconjugation has been calculated to be more important in carbenium ions than in silicenium ions. Thus, at MP2/6-31G*//3-21G(*) eq 1 is exothermic by 33.1 kcal·mol⁻¹ compared with only 10.3 kcal·mol⁻¹ for eq 2 (MP3/6-31G*//6-31G*).⁶⁸



We have now studied the kinetics of hydride abstractions from β -silyl-, β -germyl-, and β -stannyl-substituted silanes in order to derive the magnitude of hyperconjugative stabilization in silicenium ions.

Methods

Colored solutions of mono- and di(*p*-methoxy)-substituted diphenylcarbenium ions were obtained by addition of the corresponding diarylchloromethanes to solutions of TiCl₄ or Me₃SiOSO₂CF₃ (TMSOTf) in CH₂Cl₂ as described previously.^{9a} The decay of absorbance upon addition of the silanes was followed photometrically using fiber optics and the work station described earlier.^{9a}

Results

As reported for other silanes,^{9a} the reactions were found to follow a second-order rate law, first order with respect to silane and first order with respect to carbenium ion, in accord with the reaction mechanism described in Scheme I. Previous work^{9a} has shown that the reaction rates are independent of the negative counterions, indicating that these are not involved in the rate-determining step. In this way, a transition state involving pentacoordinate silicon was excluded, and SET processes were eliminated by studying kinetic isotope effects. Rate-determining formation of silicenium ions has thus been derived, though silicenium ions have not yet been observed as persistent entities in the condensed phase.¹⁰

In this work most hydride abstractions were carried out with the (*p*-anisyl)(phenyl)carbenium ion (*p*-H₃COC₆H₄)PhCH⁺ in dichloromethane at -70 °C (Table III). In some cases, also the less electrophilic bis(*p*-anisyl)carbenium ion (*p*-H₃COC₆H₄)₂CH⁺ has been employed, which proved to be 60 to 117 times less reactive than (*p*-H₃COC₆H₄)PhCH⁺.

(10) (a) Prakash, G. K. S.; Keyaniyan, S.; Aniszfeld, R.; Heiliger, L.; Olah, G. A.; Stevens, R. C.; Choi, H.-K.; Bau, R. *J. Am. Chem. Soc.* 1987, 109, 5123. (b) Olah, G. A.; Heiliger, L.; Li, X.-Y.; Prakash, G. K. S. *J. Am. Chem. Soc.* 1990, 112, 5991. (c) Wang, N.; Hwu, J. R.; White, E. H. *J. Org. Chem.* 1991, 56, 471. (d) Eaborn, C. *J. Organomet. Chem.* 1991, 405, 173. (e) Lickiss, P. D. *J. Chem. Soc., Dalton Trans.* 1992, 1333. (f) Kira, M.; Hino, T.; Sakurai, H. *J. Am. Chem. Soc.* 1992, 114, 6697. (g) Olah, G. A.; Rasul, G.; Heiliger, L.; Bausch, J.; Prakash, G. K. S. *J. Am. Chem. Soc.* 1992, 114, 7737.

(11) (a) Taft, R. W., Jr.; Lewis, I. C. *J. Am. Chem. Soc.* 1958, 80, 2436. (b) Taft, R. W., Jr.; Lewis, I. C. *Tetrahedron* 1959, 5, 210. (c) Levitt, L. S.; Levitt, B. W. *Chem. Ind. (London)* 1970, 990. (d) Levitt, L. S.; Levitt, B. W. *Tetrahedron* 1973, 29, 941. (e) Levitt, L. S.; Widing, H. F. *Progr. Phys. Org. Chem.* 1976, 12, 119. (f) Reynolds, W. F.; Hamer, G. K.; Bassindale, A. R. *J. Chem. Soc., Perkin Trans. II* 1977, 971.

(12) Calculated according to eq 3 in: Davis, D. D. *J. Organomet. Chem.* 1981, 206, 21. See also ref 2a.

Table III. Second-Order Rate Constants for the Reactions of Diarylcarbenium Ions with β -Element-IV-Substituted Silanes in CH₂Cl₂ at -70 °C

silane	X, Y from (<i>p</i> -XC ₆ H ₄)(<i>p</i> -YC ₆ H ₄)CH ⁺	<i>k</i> ₂ (-70 °C), L·mol ⁻¹ ·s ⁻¹
HSiMe ₂ (CH ₂ CMe ₃)	OMe, H	1.65 × 10 ²
HSiMe ₂ (CH ₂ SiMe ₃)	OMe, H	4.99 × 10 ²
HSiMe ₂ (CH ₂ GeMe ₃)	OMe, H	8.69 × 10 ²
HSiMe ₂ (CH ₂ SnMe ₃)	OMe, OMe	5.82 × 10 ¹
	OMe, H	3.53 × 10 ³
HSiMe(CH ₂ SiMe ₃) ₂	OMe, OMe	5.89
	OMe, H	5.12 × 10 ²
HSi(CH ₂ SiMe ₃) ₃	OMe, H	5.84 × 10 ¹
HSi(CH ₂ SiMe ₂ ⁿ Bu) ₃	OMe, OMe	8.34 × 10 ⁻¹
	OMe, H	9.79 × 10 ¹
HSi(CH ₂ SiMe ₂ Ph) ₃	OMe, OMe	3.29 × 10 ⁻¹
	OMe, H	3.44 × 10 ¹

Scheme II. Influence of the Number of β -(Trimethylsilyl) Substituents on the Hydride Donation Abilities of the Silanes HSiMe_{3-n}(CH₂SiMe₃)_n (*n* = 0–3)

Silane	Me	CH ₂ SiMe ₃	CH ₂ SiMe ₃	CH ₂ SiMe ₃
	H-Si-Me	H-Si-Me	H-Si-CH ₂ SiMe ₃	H-Si-CH ₂ SiMe ₃
	Me	Me	Me	CH ₂ SiMe ₃
<i>k</i> _{rel}	= 1.0 ^a	7.8	8.0	0.9

^a Reference value: *k*₂ = 63.7 L·mol⁻¹ s⁻¹.^{9a}

Table IV. Comparison of the Rates of Hydride Transfer from β -Substituted Silanes HSi(CH₂SiMe₂R)₃ toward (*p*-H₃COC₆H₄)PhCH⁺ with the Substituent Constants $\sigma_1(\text{R})$ ¹¹

HSi(CH ₂ SiMe ₂ R) ₃ R =	<i>k</i> _{rel}	$\sigma_1(\text{R})$
Ph	= 1.00	+0.100
Me	1.70	-0.046
ⁿ Bu	2.85	-0.054

In order to corroborate that the observed decay of (*p*-H₃COC₆H₄)PhCH⁺ in the presence of HSiMe₂(CH₂SnMe₃) indeed corresponds to the hydride abstraction from silicon and not to a reaction at another position, we have combined this carbenium ion with the analogous tetrasubstituted silane Me₃-Si(CH₂SnMe₃). Now, the carbenium ion concentration remained constant within 15 min, while in the corresponding experiment with HSiMe₂(CH₂SnMe₃) the reaction was complete after approximately 1 min.

Discussion

The introduction of a β -trimethylsilyl group in trimethylsilane increases the rate of hydride transfer by a factor of 7.8 (Scheme II). While the second β -(trimethylsilyl) group has no effect on reactivity, the third depresses the nucleophilicity to give a rate constant almost identical to that of trimethylsilane. These observations can be interpreted by a balance of steric and electronic effects. Whereas electronic effects are dominant when the first substituent is introduced, electronic and steric effects cancel for the second substituent, and steric effects dominate when the third trimethylsilyl group is added.

Table IV shows that a slight reactivity increase occurs as the inductive donation of the substituents at the β -silicon is growing.

A plot of log *k*₂ versus the σ_1 substituent constants (Figure 1) of the β -substituents shows a linear correlation for X = Cl to CMe₃ (σ_1 = -0.070 to 0.470). The β -metal-substituted silanes, however, react faster than expected on the basis of σ_1 (Figure 1) in accord with hyperconjugative stabilization.

Figure 2 shows a linear correlation between the reactivities of HSiMe₂(CH₂MMe₃) and σ^* of CH₂MMe₃, but points for other substituents (CH₂-H, CH₂-Ph, and CH₂-Cl) deviate from this line. The limited range of this correlation is not surprising, as σ^*




Figure 1. Correlation between the reactivities of the silanes $\text{HSiMe}_2(\text{CH}_2\text{X})$ (toward $(\rho\text{-H}_3\text{COC}_6\text{H}_4)\text{PhCH}^+$, CH_2Cl_2 , -70°C) with the σ_1 values of X^{I} ($\rho = -5.24$, $r = 0.998$). Regression analysis was made without $\text{X} = \text{SiMe}_3$, GeMe_3 , and SnMe_3 ; the k_2 values for $\text{X} = \text{Cl}$, Ph , H , and Me were taken from ref 9a.





Figure 2. Correlation of the reactivities of the β -element-substituted silanes $\text{HSiMe}_2(\text{CH}_2\text{MMe}_3)$ ($\text{M} = \text{C}, \text{Si}, \text{Ge}, \text{Sn}$) toward $(\rho\text{-H}_3\text{COC}_6\text{H}_4)\text{PhCH}^+$ (CH_2Cl_2 , -70°C) with Davis' σ^+ substituent constants¹² ($\rho = -2.76$, $r = 0.994$).

values also are not generally suitable for describing the effects of substituents directly attached to carbenium centers.¹³

The absolute value of the slope in Figure 2 shall, therefore, not be overinterpreted, but it can be used for comparison with the effects of the substituents CH_2MMe_3 on other reactions. As the slope of this correlation ($\rho = -2.76$) is considerably smaller than in the analogous plot of $\log k$ values versus $\sigma^+(\text{CH}_2\text{MMe}_3)$ for hydride abstractions from the tetraethylelement compounds listed in Table I ($\rho = 13.8$),^{4b,c} it is concluded that σ - π interaction is considerably weaker in the transition state of hydride abstractions yielding silicium ions than in the transition state yielding carbenium ions (Scheme III). This is probably a result of poorer orbital overlap for silicon than for carbon and the reluctance of silicon to participate in multiple bonding.^{6f,g} An analogous dependence of the rate constants on σ^+ of CH_2MR_3 has been observed for the attack of benzhydryl cations at allylelement compounds, which also produces β -element-stabilized carben-

Scheme III. Reaction Constants ρ for Reactions with Rate-Limiting Formation of a β -Stabilized Cationic Center



$\text{M}, \text{M}' = \text{C}, \text{Si}, \text{Ge}, \text{Sn}; \text{R} = \text{Cl}, \text{Ph}, \text{Me}, ^n\text{Bu}$.

ium ions ($\rho = -13.6$, Scheme III). A smaller ρ value is calculated, however, when $\log k$ for the hydride abstractions from 1,3-dimetallopropanes is plotted against $\sum\sigma^+(\text{CH}_2\text{MMe}_3)$ ($\rho = -7.32$).^{4a}

Conclusion

All observations concur that hyperconjugation is effective for stabilizing silicium ions, but the magnitude of this effect is considerably smaller than in comparable carbenium ions.

Experimental Section

Details of the kinetic procedure and preparations for the carbenium precursors have been published elsewhere.^{9a} The organoelement compounds have been synthesized according to Popowski.¹⁴ For $\text{HSiMe}_2(\text{CH}_2\text{GeMe}_3)$ the published procedure has been modified by using THF as the solvent for Me_3GeCl . In this way, the yield of side products [$\text{HSiMe}_2(\text{CH}_2\text{SiMe}_3)$] could be reduced.

Dimethyl((trimethylgermyl)methyl)silane. A solution of (bromomethyl)dimethylsilane (16.0 g, 0.104 mol) in ether (50 mL) was added dropwise to a suspension of magnesium turnings (2.80 g, 0.115 mol) in ether (50 mL) in an argon atmosphere. The solution of the Grignard reagent was decanted from the magnesium, and the yield was determined by acidimetric titration (99%). Next, 70 mL of the ethereal solution of (dimethylsilyl)methylmagnesium bromide (0.070 mol) thus obtained was added dropwise to a solution of Me_3GeCl (10.0 g, 0.065 mol) in tetrahydrofuran (70 mL). The mixture was stirred for 4 h at 35°C , kept at ambient temperature for 15 h, and then hydrolyzed with an aqueous NH_4Cl solution (20%). The organic layer was separated, and the aqueous layer was extracted with ether. After the combined organic layers were washed with NaHCO_3 solution and water and dried over CaCl_2 , and the solvents evaporated, the mixture was distilled over a 10-cm Vigreux column to give $\text{HSiMe}_2(\text{CH}_2\text{GeMe}_3)$ (4.74 g, 38%) with bp 130°C (lit.¹⁴ bp $134\text{--}135^\circ\text{C}$). GC analysis indicated a purity of >99%. $^1\text{H-NMR}$ (CCl_4): δ -0.18 (CH_2), 0.09 (SiCH_3), 0.19 (GeCH_3), 4.05 (SiH). $^{29}\text{Si-NMR}$ (CDCl_3): δ -16.03, $J(^{29}\text{Si-H}) = 181.0$ Hz, SiMe_4 reference. Mass spectrum (70 eV): m/z (rel intensity) 191 (2.6%, $\text{M}^+ - \text{H}$), 177 (92.9%, $\text{M}^+ - \text{CH}_3$), 119 (28.5%, Me_3Ge^+), 73 (100%, Me_3Si^+). For IR data see ref 14.

Dimethyl(neopentyl)silane. Neopentyl bromide and lithium were combined according to published procedures¹⁵ to give neopentyl lithium in 28 and 29% yield (by Gilman titration¹⁶), respectively. A solution of neopentyl lithium (50 mmol) in *n*-hexane (100 mL) was added dropwise to a solution of chlorodimethylsilane (4.26 g, 45 mmol) in ether (50 mL, purified by distillation over Na and Ph_2CO), which was cooled in an ice bath (15 min, Ar atmosphere). The mixture was then heated under reflux for 5 h and worked up as described above to give 40–50% of $\text{HSiMe}_2(\text{CH}_2\text{CMe}_3)$ (2.35–2.93 g) with bp $110\text{--}112^\circ\text{C}$ (lit.¹⁷ bp 110.2°C). A contamination (probably 2,2,5,5-tetramethylhexane formed during the preparation of neopentyl lithium) could not be removed by distillation. The product has, therefore, been purified by preparative GC (column length 3 m, o.d. 1 cm; 20% GE SE 30 on chromosorb WAW DMCS, oven temperature 80°C). IR (*n*-hexane): $\bar{\nu}$ 2120.3 cm^{-1} (SiH). $^1\text{H-NMR}$ (CCl_4): δ 0.07 (SiCH_3), 0.70 (CH_2), 0.99 (CCH_3), 3.99 (SiH). $^{29}\text{Si-NMR}$ (CDCl_3): δ -16.03, $J(^{29}\text{Si-H}) = 181.0$ Hz, SiMe_4 reference.

(14) Popowski, E.; Marekova, U.; Reiske, T.; Schulz, H.; Kelling, H.; Jancke, H. *Z. Anorg. Allg. Chem.* **1990**, *583*, 195.

(15) (a) Beachley, O. T., Jr.; Victoriano, L. *Organometallics* **1988**, *7*, 63.

(b) Schrock, R. R.; Fellmann, J. D. *J. Chem. Soc.* **1978**, *100*, 3359.

(16) Gilman, H.; Cartledge, F. K. *J. Organomet. Chem.* **1964**, *2*, 447.

(17) Heflfejs, J.; Mares, F.; Chvalovsky, V. *Collect. Czech. Chem. Commun.* **1966**, *31*, 586.

(13) (a) Koshy, K. M.; Roy, D.; Tidwell, T. T. *J. Am. Chem. Soc.* **1979**, *101*, 357. (b) Chwang, W. K.; Knittel, P.; Koshy, K. M.; Tidwell, T. T. *J. Am. Chem. Soc.* **1977**, *99*, 3395. (c) Oyama, K.; Tidwell, T. T. *J. Am. Chem. Soc.* **1976**, *98*, 947. (d) Nowlan, V. J.; Tidwell, T. T. *Acc. Chem. Res.* **1977**, *10*, 252.

Mass spectrum (70 eV): m/z (rel intensity) 130 (0.25%, M^+), 129 (1.2%, $M^+ - H$), 115 (100%, $M^+ - CH_3$), 73 (78.7%, $M^+ - CMe_3$).

(Trimethylsilyl)(trimethylstannyl)methane. The literature procedure¹⁸ was improved as follows: trimethylstannyl chloride (15.0 g, 75.3 mmol) was dissolved in 50 mL of ether. A solution of Me_3SiCH_2Li (91.2 mmol, prepared analogous to the corresponding Grignard reagent in ref 14 in 95 mL of *n*-hexane) was added dropwise within 15 min with water-ice cooling (Ar atmosphere). The mixture was stirred for 4 h at 35 °C, kept at ambient temperature for 15 h, and then worked up as described above. Distillation over a packed column yielded 11.7 g (62%, bp 164–166 °C, lit.¹⁸ bp 165–166 °C). ¹H-NMR (CCl_4): δ -0.25 (CH_2), 0.02 ($SiCH_3$),

0.09 ($SnCH_3$). Mass spectrum (70 eV): m/z (rel intensity) 237 (100, $M^+ - CH_3$), 165 (59.2%, Me_3Sn^+), 135 (11.4%, $MeSn^+$), 73 (46.1%, Me_3Si^+).

Acknowledgment. Financial support by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie is gratefully acknowledged. We thank Dr. B. Dogan for critical discussions.

Supplementary Material Available: One table with concentrations and rate constants of the kinetic experiments described on Table III (3 pages). Ordering information is given on any current masthead page.

(18) (a) Papetti, S.; Post, H. W. *J. Org. Chem.* 1957, 22, 526. (b) Schmidbaur, H. *Chem. Ber.* 1964, 97, 270.