

Models for Surface Catalysis: A Remarkable $\text{Pt}_3(\text{ReO}_3)$ Cluster Cation

Jianliang Xiao and Richard J. Puddephatt*

Department of Chemistry
University of Western Ontario
London, Canada N6A 5B7

Ljubica Manojlović-Muir, Kenneth W. Muir,* and
Ali Ashgar Torabi

Department of Chemistry
University of Glasgow
Glasgow, Scotland G12 8QQ

Received November 9, 1993

Metal–metal bonded clusters may occur with the metal atoms in either high oxidation states, where the ligands are typically π -donors such as halide, alkoxide, oxide, or sulfide, or in low oxidation states, where the ligands are typically π -acceptors such as CO, isocyanides, or phosphines.¹ These two classes of metal clusters have little in common and are usually treated separately.^{1,2} In particular, “hybrid” metal clusters containing both types of metal–ligand fragments are essentially unknown, although there are now several binuclear complexes with two metals in different oxidation states.^{1–4} This is unfortunate since clusters containing both metal oxide and metal π -acceptor ligand fragments could not only display unique chemistry but also provide insights into the interaction of metal cluster surfaces with oxide materials in areas such as heterogeneous catalysis. This article reports the synthesis and structure of the cluster cation $[\text{Pt}_3(\text{ReO}_3)(\mu\text{-dppm})_3]^+$ (**1**, dppm = $\text{Ph}_2\text{PCH}_2\text{PPh}_2$), which contains both a high oxidation state ReO_3 fragment and a low oxidation state $\text{Pt}_3(\mu\text{-dppm})_3$ unit. The stability of this new cluster suggests that a much wider spectrum of “hybrid” clusters might exist. The cluster is also interesting as a model for possible Pt–Re bonding and metal–support interactions in the important heterogeneous bimetallic Pt–Re/ Al_2O_3 catalysts.⁵

The cluster cation $[\text{Pt}_3\{\text{Re}(\text{CO})_3(\mu\text{-dppm})_3\}]^+$ (**2**) is known to react with Me_3NO or O_2 at room temperature to give the oxidized clusters $[\text{Pt}_3\{\text{Re}(\text{CO})_3(\mu_3\text{-O})(\mu\text{-dppm})_3\}]^+$ (**3**) and $[\text{Pt}_3\{\text{Re}(\text{CO})_3(\mu_3\text{-O})_2(\mu\text{-dppm})_3\}]^+$ (**4**), respectively.⁴ It has now been shown that the reaction of **2** with O_2 in refluxing *o*-xylene gives **4** [PF_6], which reacts further with O_2 to give the remarkable cluster **1** [PF_6]. More extended reflux resulted in partial decomposition and formation of **1** [ReO_4].⁴

(1) (a) Mingos, D. M. P.; Wales, D. J. *Introduction to Cluster Chemistry*; Prentice Hall: Englewood Cliffs, NJ, 1990. (b) Shriver, D. F.; Kaesz, H. D.; Adams, R. D. *The Chemistry of Metal Cluster Complexes*; VCH: New York, 1990.

(2) (a) Cotton, F. A. *ACS Symp. Ser.* **1983**, 211, 209. (b) Ara, I.; Fanwick, P. E.; Walton, R. A. *Inorg. Chem.* **1992**, 31, 3211. (c) Chisholm, M. H.; Huffman, J. C.; Van Der Sluis, W. G. *J. Am. Chem. Soc.* **1987**, 109, 2514.

(3) Several carbonyl clusters containing one or two oxide ligands are known, but in all cases the oxide is present as a μ_3 ligand, not as an identifiable metal oxide fragment. Ge, Y.-W.; Sharp, P. R. *Inorg. Chem.* **1992**, 31, 379.

(4) Xiao, J.; Vittal, J. J.; Puddephatt, R. J.; Manojlović-Muir, K. W. *J. Am. Chem. Soc.* **1993**, 115, 7882. In one experiment using refluxing xylene, the yield of **1** [PF_6] was ca. 90% after 7 h and the yield of **1** [ReO_4] was ca. 30% after 15 h.

(5) Klabunde, K. J.; Li, Y.-X. *Clusters, Alloys, and Poisoning, An Overview. In Selectivity in Catalysis*; Davis, M. E., Suib, S. L., Eds.; ACS Symposium Series 517, American Chemical Society: Washington, DC, 1993.

(6) Spectroscopic and X-ray data for $[\text{Pt}_3(\text{ReO}_3)(\mu\text{-dppm})_3][\text{ReO}_4]$ (**1** [ReO_4]; triclinic, $P\bar{1}$; $a = 14.154(4)$, $b = 14.213(3)$, and $c = 20.522(4)$ Å; $\alpha = 78.938(14)^\circ$, $\beta = 75.812(19)^\circ$, and $\gamma = 74.957(22)^\circ$; $V = 3828.9(15)$ Å³, $Z = 2$; Enraf-Nonius CAD4 diffractometer with Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å; $R_w = 0.069$, $R = 0.077$ for 284 parameters and 5414 reflections with $I \geq 3\sigma(I)$. IR (Nujol mull): $\nu(\text{Re}=\text{O})$ (ReO_4^-) = 915 (st), 905 (m); $\nu(\text{Re}=\text{O})$ (ReO_3) = 940 (w), 890 (st). ³¹P NMR (CD_2Cl_2): $\delta = -2.8$ [s, ¹(PtP) = 3134, ³J(PP) = 133 Hz].

Table 1. Comparison of Metal–Metal Distances (Å) in **1**, **2**, and **4**

	1	2	4
Pt(1)–Pt(2)	2.598(2)	2.611(1)	2.826(1)
Pt(1)–Pt(3)	2.609(3)	2.593(1)	3.094(1)
Pt(2)–Pt(3)	2.600(3)	2.608(1)	3.081(1)
Pt(1)–Re	2.720(3)	2.684(1)	2.843(1)
Pt(2)–Re	2.748(3)	2.649(1)	2.854(1)
Pt(3)–Re	2.711(3)	2.685(1)	3.228(1)





Figure 1. View of the $[\text{Pt}_3(\text{ReO}_3)(\mu\text{-dppm})_3]^+$ cation and its associated $[\text{ReO}_4]^-$ anion. 50% probability ellipsoids are displayed. For clarity, hydrogen atoms are omitted, as are all but the *ipso* carbon atom of each of the 12 phenyl rings. The range of Pt–P distances is 2.274(9)–2.306(9) Å. Metal–oxygen distances are Re(1)–O(1) 1.74(3), Re(1)–O(2) 1.70(3), Re(1)–O(3) 1.77(3), Re(2)–O(4) 1.71(4), Re(2)–O(5) 1.80(4), Re(2)–O(6) 1.67(3), and Re(2)–O(7) 1.68(4) Å.



The structure of **1** [ReO_4] has been determined by X-ray diffraction (see Table 1 and Figure 1).⁶ The crystals contain $[\text{Pt}_3(\text{ReO}_3)(\mu\text{-dppm})_3]^+$ cluster cations based on a tetrahedral Pt_3Re core. Each Pt–Pt edge is bridged by a dppm ligand, forming the well-known latitudinal $\text{Pt}_3(\mu\text{-dppm})_3$ moiety. A face of the Pt_3 triangle is μ_3 -bridged by the $\text{Re}(1)\text{O}_3$ unit. Re(1) thereby adopts a highly distorted octahedral coordination which retains local C_3 symmetry, with Pt–Re(1)–Pt angles of 56.7(1)–57.4(1) $^\circ$ and O–Re(1)–O angles of 104(1)–110(1) $^\circ$. Thus **1** is closely related structurally to **2**, with oxo ligands replacing carbonyl. The Pt–Pt distances in both **1** and **2** are nearly constant, and their means [2.602(4) and 2.604(6) Å] are virtually identical. The Pt–Re distances in each cluster are rather less regular, and the mean Pt–Re distance in **1** [2.726(11) Å] is somewhat greater than the corresponding mean [2.673(12) Å] for **2**. The Re(1)–O distances in **1** [1.70(3)–1.77(3) Å] are unexceptional.⁷

(7) The mean length of 54 terminal Re–O (oxo) bonds is 1.709 Å. *International Tables for Crystallography*, Volume C, International Union of Crystallography; Kluwer Academic Publishers: Dordrecht, 1992; p 745.




Figure 2. Qualitative MO correlation diagram for formation of the cation **1** from the fragments $\text{Pt}_3(\mu\text{-H}_2\text{PCH}_2\text{PH}_2)_3$ and ReO_3^+ . The three highest energy occupied MOs are the M–M bonding orbitals of a_1 and e symmetry, while the LUMO is the M–M antibonding level of e symmetry.

Each cation in **1** [ReO_4] has closely associated with it a [ReO_4][−] anion which blocks the free face of the Pt_3 triangle: the Re(2) atom is nearly equidistant from Pt(2) and Pt(3) (4.97 and 4.96 Å) and is slightly further (5.29 Å) from Pt(1); the anion is oriented so that the Re(2)–O(7) vector points toward the midpoint of Pt(2)–Pt(3). The resulting short O(7)···Pt contacts, respectively 3.98, 3.41, and 3.37 Å, are, however, clearly nonbonding. We have previously noted the ability of the free face of a latitudinal $\text{Pt}_3(\mu\text{-dppm})_3$ ($M = \text{Pt}, \text{Pd}$) moiety to encapsulate a loosely bound anion: for example, above the free Pd_3 face of $[\text{Pd}_3(\mu_3\text{-CO})(\mu\text{-dppm})_3]^{2+}$ there is an associated CF_3CO_2^- anion which participates in $\text{Pd}\cdots\text{O}$ contacts of 2.77(1)–2.92(2) Å, rather shorter than the $\text{Pt}\cdots\text{O}(7)$ contacts described here.⁸ In the IR spectrum (Nujol mull) of **1** [ReO_4], two bands at 915 and 905 cm^{-1} could be assigned to $\nu(\text{Re}=\text{O})$ of coordinated ReO_4^- (free ReO_4 gives only one band).⁹ In addition, two bands due to $\nu(\text{Re}=\text{O})$ in the $\text{Re}(\text{=O})_3$ fragment were present at 940 and 890 cm^{-1} in both **1** [PF_6] and **1** [ReO_4]. For comparison, the complex $[(\eta^5\text{-C}_5\text{Me}_5)\text{Re}(\text{O})_3]$ gives $\nu(\text{Re}=\text{O})$ at 909 and 878 cm^{-1} .¹⁰ The ¹H and ³¹P NMR spectra of **1** [PF_6] and **1** [ReO_4] in CD_2Cl_2 solution were identical, indicating that the ReO_4^- ion is probably not associated with **1** in solution.^{6,11}

The overall transformation of **2** to **1** involves the replacement of the three carbonyl ligands in **2** by the three terminal oxo ligands in **1**. Since both CO and the terminal oxo ligand are formally two-electron donors, the overall cluster counts in **1** and **2** are the same. Both **1** and **2** are coordinatively unsaturated 54-electron

clusters, and it is therefore expected that the cluster cores should be similar. In contrast, cluster **4** is formed by the addition of two $\mu_3\text{-O}$ ligands to **2** without carbonyl dissociation and is a 62-electron cluster with much weaker metal–metal bonding.⁴

The bonding in **1** and **2** can be understood in terms of the donation of electron density from the three filled Pt–Pt bonding orbitals, having $a_1 + e$ symmetry, of the $\text{Pt}_3(\text{dppm})_3$ fragment to the three vacant acceptor orbitals, also having $a_1 + e$ symmetry, of a $\text{Re}(\text{=O})_3^+$ fragment as shown in Figure 2. In this formalism, the platinum and rhenium atoms in **1** may be considered as Pt(O) and Re(VII). Even though this is an extreme interpretation, the oxidation states of platinum and rhenium in **1** are clearly very different to an extent which is unprecedented in transition-metal clusters.

The ReO_3 fragment is present in several other unusual compounds, such as MeReO_3 , $[(\eta^5\text{-C}_5\text{Me}_5)\text{ReO}_3]$, and $[(\text{Me}_2\text{-PCH}_2\text{PMe}_2)_2\text{Cl}_2\text{Re-ReO}_3]$, and the first two are considered as Re(VII) complexes.^{2b,10} There is a useful analogy between complexes **1** and $[(\eta^5\text{-C}_5\text{Me}_5)\text{ReO}_3]$, since the donor orbitals of both the $\text{Pt}_3(\text{dppm})_3$ fragment and the C_5Me_5^- ligand have $a_1 + e$ symmetry and so may be considered to be isolobal.¹² The analogy, together with the proven compatibility between the ReO_3^+ and $\text{Pt}_3(\mu\text{-dppm})_3$ fragments, suggests several other possible target “hybrid” clusters, by substitution of $\text{Pt}_3(\text{dppm})_3$ units for C_5Me_5^- ligands in known $(\text{C}_5\text{Me}_5)\text{MX}_n$ complexes and hence that the study of such “hybrid” clusters could become a major field of research. Specifically, since the vast majority of organometallic oxo complexes contain cyclopentadienyl ligands,¹³ the preparation of other Pt_3 fragment-based oxide clusters should be feasible.

The oxidation sequence of **2** to **3** to **4** to **1** and then [ReO_4][−] shows how a low oxidation state rhenium atom can be oxidized to perrhenate in a stepwise manner while bound to a Pt_3 triangle. It is interesting that the reverse transformation has been proposed for the platinum-catalyzed reduction of rhenium oxides in the formation of the important reforming catalyst Pt–Re/ Al_2O_3 .¹⁴ It has been suggested that rhenium oxides such as [ReO_4][−] migrate over the support surface and are reduced in contact with a platinum particle, whereupon bimetallic Pt–Re clusters are formed.¹⁴ The mechanisms of the cluster and surface reactions may well be related, and complex **1** [ReO_4] contains two types of Pt_3^- rhenium oxide interactions which may be considered as models for proposed surface intermediates.

Acknowledgment. We thank the NSERC (Canada) and SERC (U.K.) for financial support and the Iranian Ministry of Culture and Higher Education for a studentship to A.A.T.

Supplementary Material Available: Crystallographic data for $[\text{Pt}_3(\text{ReO}_3)(\mu\text{-Ph}_2\text{PCH}_2\text{PPh}_2)_3][\text{ReO}_4]$ and tables of atomic parameters, anisotropic displacement parameters, bond lengths, and bond angles (13 pages); observed and calculated structure factors (17 pages). This material is contained in many libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS; see any current masthead page for ordering information.

(8) Ferguson, G.; Lloyd, B. R.; Manojlović-Muir, L.; Muir, K. W.; Puddephatt, R. *J. Inorg. Chem.* **1986**, *25*, 4190.

(9) Ohlhausen, L.; Cockrum, D.; Register, J.; Roberts, K.; Long, G. J.; Powell, G. L.; Hutchinson, B. B. *Inorg. Chem.* **1990**, *29*, 4886.

(10) Herrmann, W. A.; Serrano, R.; Bock, H. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 383.

(11) Jennings, M. C.; Schoettel, G.; Roy, S.; Puddephatt, R. J.; Douglas, G.; Manojlović-Muir, L.; Muir, K. W. *Organometallics* **1991**, *10*, 580.

(12) Albright, T. A.; Burdett, J. K.; Whangbo, M.-H. *Orbital Interactions in Chemistry*; Wiley: New York, 1985; Chapter 20.

(13) Bottomley, F.; Sutin, L. *Adv. Organomet. Chem.* **1988**, *28*, 339.

(14) (a) Godbey, D. J.; Somorjai, G. A. *Surf. Sci.* **1988**, *202*, 204. (b) Augustine, S. M.; Sachtler, W. M. H. *J. Catal.* **1989**, *116*, 184 and reference therein.