




Additions and Corrections

Study of Sulfinato and Sulfenato Complexes Derived from the Oxygenation of Thiolate Sulfur in [1,5-Bis(2-mercapto-2-methylpropyl)-1,5-diazacyclooctanato(2-)]nickel(II) [*J. Am. Chem. Soc.* **1995**, *117*, 963–973]. RIZALIA M. BUONOMO, IVAN FONT, MICHAEL J. MAGUIRE, JOSEPH H. REIBENSPIES, THAWATCHAI TUNTULANI, AND MARCETTA Y. DARENSBOURG


Page 970, Table 4: Ni–N, Ni–S, and S–O bond distances of compounds **1***, **2***, and **4*** should read as follows:

Compound	Ni–N (Å)	Ni–S (Å)		S–O (Å)
1*	1.995(3)	2.152(1)		
	1.995(3)	2.152(1)		
2*	1.99(1)	2.109(4)		1.47(1)
	2.00(1)	2.132(4)		1.46(1)
4*	2.016(9)	2.162(3)		1.548(8)
	1.998(7)	2.148(3)		

JA9550081

A [10]Annulene Isomer May Be Aromatic, After All! [*J. Am. Chem. Soc.* **1995**, *117*, 1369–1373]. HORST M. SULZBACH, PAUL VON RAGUE SCHLEYER,* HAIJUN JIAO, YAOMING XIE, AND HENRY F. SCHAEFER III*

Page 1369: Structure **III** should have five double bonds as shown below.



III

JA9550079

Macrocyclic Peptide Inhibitors of Serine Proteases. Convergent Total Synthesis of Cyclotheonamides A and B via a Late-Stage Primary Amine Intermediate. Study of Thrombin Inhibition under Diverse Conditions [*J. Am. Chem. Soc.* **1995**, *117*, 1225–1239]. BRUCE E. MARYANOFF,* MICHAEL N. GRECO, HAN-CHENG ZHANG, PATRICIA ANDRADE-GORDON, JACK A. KAUFFMAN, K. C. NICOLAOU, AIJUN LIU, AND PETER H. BRUNGS

Page 1233, column 1: The standard rate equation for slow-binding kinetics given in the second paragraph should read as follows: $P = v_s t + (v_0 - v_s)(1 - e^{-kt})/k$.

JA955009T