


Sidearm Effect: Improvement of the Enantiomeric Excess in the Asymmetric Michael Addition[†] of Indoles to Alkylidene Malonates

Jian Zhou and Yong Tang*


State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry,
Chinese Academy of Sciences, 200032 Shanghai, China

Received May 17, 2002

C₂-symmetric chiral bisoxazoline-metal complexes enjoy high prestige in asymmetric catalysis.¹ For example, bisoxazoline **1** derived Cu(II) complexes² have been well established as versatile catalysts for asymmetric C–C bond formations such as aldol,³ cycloaddition,⁴ ene,⁵ Michael,⁶ amination,⁷ Friedel–Crafts,⁸ Henry,⁹ and Mannich¹⁰ reactions. In most cases, excellent enantioselectivity could be achieved only when *tert*-leucinol-derived bisoxazoline **1b** was used. In sharp contrast to the great success of bisoxazolines, the development and application of trisoxazolines are rather limited.¹¹ In our efforts to develop superior catalysts which are cheap, easy to access, air-stable, and water-tolerant, we considered a sidearm approach, focusing on the improvement of bisoxazoline into pseudo-C₃-symmetric trisoxazoline **2**, for the following reasons: (1) compared with C₂-symmetric bisoxazolines **1**, coordination of three nitrogen atoms of trisoxazoline **2** to the metallic center induces a more tunable pseudo-C₃-symmetric chiral space, to shield distant reactive sites; (2) tridentate ligand **2** is expected to increase the stability of active intermediates with respect to ordinary bidentate bisoxazolines.¹² Thus, the metal complex may be tolerable to moisture, and it is possible to improve the catalytic activity. In this communication, we report our preliminary results on this subject.



The trisoxazoline **2** can be easily synthesized from **3** and L-valinol by two steps. Compound **3** reacted with L-valinol without solvent¹³ at 70 °C to afford white hygroscopic solid **4** in 61% yield. Treatment of compound **4** with PPh₃/CCl₄¹⁴ afforded desired trisoxazoline **2** as colorless oil in 75% yield.



Very recently, Jørgensen et al. pioneered asymmetric Michael addition of indoles to alkylidene malonates.^{8c} They found that chiral Lewis acid **1b**/Cu(OTf)₂ could promote this reaction in moderate-to-good ee value (up to 69% ee). To examine if our design is operative, we tried the above reaction using the newly designed trisoxazoline **2**/Cu(ClO₄)₂·6H₂O^{12b,15} complex as the chiral Lewis

Table 1. Effect of Temperature and Additive on the Reaction between Indole **5a** and Benzylidene Malonate **6a**

entry	catalyst loading (%)	additive	temp (°C)	yield ^a (%)	ee ^{b,c} (%)
1	10	None	15	88	82
2	10	None	0	50	85
3	10	HFIP	0	99	85
4 ^d	1	HFIP	0	90	78
5	10	HFIP	−20	84	89
6	10	HFIP	−25	56	93

^a Isolated yield and performed on 0.25 mmol scale unless otherwise noted. ^b Determined by chiral HPLC. ^c Absolute configuration was assigned by chemical transformation. ^d Performed on 4 mmol scale. ^e All reactions were run in acetone–ether (1:3, v/v).

acid. We are pleased to find that the catalytic activity was comparable to that of **1b**/Cu(OTf)₂ and the enantioselectivity of the reaction of indole with benzylidene malonate was enhanced greatly (entry 1, Table 1).

The alkylation of indole **5a** with malonate **6a** proceeded quite slowly at 0 °C, and the yield was only 50%; even the reaction time was prolonged to 72 h (entry 2 in Table 1) in the absence of hexafluoro-*i*-PrOH (HFIP). To our delight, we found that the addition of 2 equiv of HFIP¹⁶ can greatly improve the reactivity without loss of ee (entry 3). The reaction was proved to be temperature-dependent. When the reaction temperature decreased from 0 to −25 °C, the ee value of compound **7a** increased from 85 to 93% (entries 3, 5, and 6 in Table 1).

The high yield and excellent enantioselectivity encouraged us to study the generality of this reaction by investigating a variety of structurally different indole derivatives and alkylidene malonates (Table 2). Arylidene malonates with indole worked well to afford the desired products in excellent yields with high ee. Unlike the ee that were different between dimethyl and diethyl malonates in the case of **1b**/Cu(OTf)₂, both methyl ester and ethyl ester maintained excellent face selectivity when trisoxazoline was used (entries 1–4). Enantiomeric excesses ranging from 90 to 92% were obtained in high yields for various substituted arylidene malonates (entry 4–8). Diethyl ethylidene malonate could also react with indole smoothly, but it was less enantioselective (entry 9). The reaction of benzylidene malonate with 4-methoxyindole, 5-methoxyindole, and 5-methylindole resulted in the alkylation products in high-to-excellent yields with excellent ee value (entries 10, 11, 12), indicating that the substituents on indole ring had almost no effect on the enantioselectivity.

It is noteworthy that trisoxazoline **2**/Cu(ClO₄)₂·6H₂O complex is an air- and water-stable compound. *It maintained the same*

* To whom correspondence should be addressed. E-mail: tangy@pub.sioc.ac.cn.

[†] This reaction is also regarded as Friedel–Crafts reaction in ref 8c.

Table 2. Asymmetric alkylation of indoles **5** with various alkylidene malonates **6**

entry	malonate 6	indole 5	yield % ^b	ee % ^c
1			84	89
2			98	88 (50) ^d
3			99	91 (56) ^d
4			99	91
5			99	91
6			95	90 (50) ^d
7			99	92 (69) ^d
8			84	90
9 ^c			84	60
10			73	91 (60) ^d
11			92	93
12			97	91
13			95	91 ^f


^a Reactions were run in acetone–ether (1:3 v/v) with 10 mol % chiral catalyst at -20°C . ^b Isolated yields. ^c Determined by chiral HPLC. ^d Number in parentheses refers to the best result in the literature. ^e Run at -35°C . ^f Performed under air atmosphere.

catalytic reactivity almost without loss of enantioselectivity even when the reaction was carried out under air atmosphere (entry 13 in Table 2).

Optical purity of the products could be enhanced by recrystallization. For example, 99.6% ee was obtained in 71% yield after compound **7a** (86% ee) was recrystallized from a mixed solvent of petroleum ether, CH_2Cl_2 , and acetone. These indole derivatives prepared by the current method are potentially synthetically useful because they can undergo many chemical transformations, in particular for the preparation of β -substituted tryptophans¹⁷ and some bioactive indole derivatives.¹⁸

The high ee values obtained with L-valinol-derived trisoxazoline **2** are rather intriguing. The proposed transition-state model shown below supports all the experimental observation and is consistent with the absolute configuration of some selected products. The indoles could only approach the *re* face of the alkylidene malonates due to steric hindrance.

In conclusion, we have demonstrated that pseudo- C_3 -symmetric trisoxazoline **2** has the encouraging ability to achieve high face selectivity in asymmetric Michael reaction of indoles to alkylidene malonates. The cheap and easy synthesis of trisoxazoline–Cu (II) complexes, the high selectivity, and the mild reaction conditions including water- and air-tolerance make our method potentially



useful. In particular, our ligand design strategy may also provide a basis for further optimization of the sidearmed bisoxazoline to enhance both enantioselectivity and catalytic efficiency.

Acknowledgment. We are grateful for the financial support from the National Natural Sciences Foundation of China.

Supporting Information Available: Characterization data for all compounds, absolute configuration, and experimental procedures (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Ghosh, A. K.; Mathivanan, P.; Cappiello, J. *Tetrahedron: Asymmetry* **1998**, *9*, 1.
- (a) Evans, D. A.; Rovis, T.; Johnson, J. S. *Pure Appl. Chem.* **1999**, *71*, 1407. (b) Jørgensen, K. A.; Johannsen, M.; Yao, S.; Audrain, H.; Thorhaug, J. *Acc. Chem. Res.* **1999**, *32*, 605. (c) Johnson, J. S.; Evans, D. A. *Acc. Chem. Res.* **2000**, *33*, 325.
- (a) Evans, D. A.; Murry, J. A.; Kozłowski, M. C. *J. Am. Chem. Soc.* **1996**, *118*, 5814. (b) Evans, D. A.; Kozłowski, M. C.; Burgey, C. S.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **1997**, *119*, 7893 and references therein.
- For DA reactions, see: (a) Evans, D. A.; Miller, S. J.; Lectka, T. *J. Am. Chem. Soc.* **1993**, *115*, 6460. (b) Evans, D. A.; Miller, S. J.; Lectka, T.; von Matt, P. *J. Am. Chem. Soc.* **1999**, *121*, 7559 and references therein. For HAD reaction, see: (c) Johannsen, M.; Jørgensen, K. A. *J. Org. Chem.* **1995**, *60*, 5757. (d) Yao, S.; Johannsen, M.; Audrain, H.; Hazell, R. G.; Jørgensen, K. A. *J. Am. Chem. Soc.* **1998**, *120*, 8599. (e) Evans, D. A.; Johnson, J. S. *J. Am. Chem. Soc.* **1998**, *120*, 4895 and references therein. For cycloadditions of silyl ketenes, see: (f) Evans, D. A.; Janey, J. M. *Org. Lett.* **2001**, *3*, 2125.
- Evans, D. A.; Burgey, C. S.; Paras, N. A.; Vojkovsky, T.; Tregay, S. W. *J. Am. Chem. Soc.* **1998**, *120*, 5824 and references therein.
- (a) Evans, D. A.; Willis, M. C.; Johnston, J. N. *Org. Lett.* **1999**, *1*, 865. (b) Evans, D. A.; Rovis, T.; Kozłowski, M. C.; Tedrow, J. S. *J. Am. Chem. Soc.* **1999**, *121*, 1994 and references therein.
- (a) Evans, D. A.; Johnson, D. S. *Org. Lett.* **1999**, *1*, 595. (b) Juhl, K.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2002**, *124*, 2420.
- (a) Gathergood, N.; Zhuang, W.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2000**, *122*, 12517. (b) Jensen, K. B.; Thorhaug, J.; Hazell, R. G.; Jørgensen, K. A. *Angew. Chem., Int. Ed.* **2001**, *40*, 160. (c) Zhuang, W.; Hansen, T.; Jørgensen, K. A. *Chem. Commun.* **2001**, 347.
- (a) Christensen, C.; Juhl, K.; Jørgensen, K. A. *Chem. Commun.* **2001**, 2222. For Aza-Henry reaction: (b) Knudsen, K. R.; Risgaard, T.; Nishiwaki, N.; Gothelf, K. V.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2001**, *123*, 5843.
- Juhl, K.; Gathergood, N.; Jørgensen, K. A. *Angew. Chem., Int. Ed.* **2001**, *40*, 2995.
- (a) Kawasaki, K.; Tsumura, S.; Katsuki, T. *Synlett.* **1995**, 1245. (b) Chuang, T.-H.; Fang, J.-M.; Bolm, C. *Synth. Commun.* **2000**, *30*, 1627. (c) Kohmura, Y.; Katsuki, T. *Tetrahedron Lett.* **2000**, *41*, 3941.
- For introducing additional donating atoms into the chiral ligand to stabilize the active intermediates, see: (a) Fache, F.; Schulz, E.; Tommasino, M. L.; Lemaire, M. *Chem. Rev.* **2000**, *100*, 2159. (b) Kanemasa, S.; Oderaotoshi, Y.; Sakaguchi, S.; Yamamoto, H.; Tanaka, J.; Wada, E.; Curran, D. P. *J. Am. Chem. Soc.* **1998**, *120*, 3074.
- Kawasaki, K.; Katsuki, T. *Tetrahedron* **1997**, *53*, 6337.
- Vorbrüggen, H.; Krolikiewicz, K. *Tetrahedron* **1993**, *49*, 9353.
- For using $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ as Lewis acid, see: Ghosh, A. K.; Cho, H.; Cappiello, J. *Tetrahedron: Asymmetry* **1998**, *9*, 3687.
- Evans, D. A.; Rovis, T.; Kozłowski, M. C.; Downey, C. W.; Tedrow, J. S. *J. Am. Chem. Soc.* **2000**, *122*, 9134.
- Jeannin, L.; Nagy, T.; Vassileva, E.; Sapi, J.; Laronze, J. Y. *Tetrahedron Lett.* **1995**, *36*, 2057 and references therein.
- (a) Leditschke, H.; Ehrhart, G. U.S. Patent 2,752,358. (b) Black, W. C.; Bayly, C.; Belley, M.; Chan, C.-C.; Charleson, S.; Denis, D.; Gauthier, J. Y.; Gordon, R.; Guay, D.; Kargman, S.; Lau, C. K.; Leblanc, Y.; Mancini, J.; Ouellet, M.; Percival, D.; Roy, P.; Skorey, K.; Tagari, P.; Vickers, P.; Wong, E.; Xu, L.; Prasit, P. *Bioorg. Med. Chem. Lett.* **1996**, *6*, 725. (c) Takeda, T.; Mukaiyama, T. *Chem. Lett.* **1980**, 163. (d) Nagy, T.; Jeannin, L.; Sapi, J.; Laronze, J. Y.; Renard, P.; Pfeiffer, B.; Bizot-Espiard, J. G. *Eur. J. Med. Chem.* **1995**, *30*, 575. (e) Austin, J. F.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2002**, *124*, 1172.

JA026936K