


## Oxygen–Oxygen Bond Homolysis in a Novel Titanium(IV) Alkylperoxide Complex, Cp<sub>2</sub>Ti(OO<sup>t</sup>Bu)Cl

Antonio G. DiPasquale, Werner Kaminsky,<sup>†</sup> and James M. Mayer\*

Department of Chemistry, Campus Box 351700, University of Washington, Seattle, Washington 98195-1700


Received September 10, 2002

Metal-catalyzed oxidations by peroxides are of importance from industrial to biological chemistry.<sup>1</sup> Metal ions can bind peroxides, can activate them toward oxidation of substrates, and can catalytically decompose them. Examples include the large-scale epoxidation of propylene by alkyl hydroperoxides, the metal-mediated autoxidation of cyclohexane, <sup>1a–c,2</sup> and the Sharpless titanium-tartrate chiral epoxidation.<sup>3</sup> Metal-peroxide species are key intermediates in the reactions of a variety of oxidizing metalloenzymes,<sup>1d–f</sup> and a variety of biomimetic metal catalysts have been developed.<sup>1,4</sup> In many cases, the metal-peroxide complex is thought to undergo O–O bond cleavage to give a metal-oxo species that is the reactive oxidant.<sup>5</sup> This O–O bond cleavage, whether homolytic or heterolytic, is thought to be facilitated by oxidation of the metal center (eq 1).<sup>6</sup>




We report here a new d<sup>0</sup> titanocene *tert*-butylperoxide complex and mechanistic studies of its decomposition that implicate O–O bond homolysis *without* metal oxidation.

Cp<sub>2</sub>TiCl<sub>2</sub> (Cp = η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>) reacts with NaOO<sup>t</sup>Bu in THF at –20 °C to give Cp<sub>2</sub>Ti(OO<sup>t</sup>Bu)Cl (**1**) which is extracted into hexane and isolated in 84% yield (eq 2).<sup>7,8</sup> This synthesis follows the briefly



reported in situ generation of Cp<sub>2</sub>Zr(OO<sup>t</sup>Bu)Cl.<sup>9</sup> Related Cp<sup>\*</sup><sub>2</sub>Hf(OO<sup>t</sup>Bu)R complexes have been prepared by protolytic reactions.<sup>10</sup> Complex **1** has been characterized by <sup>1</sup>H and <sup>13</sup>C NMR, IR, high-resolution mass spectrometry, elemental analysis, and X-ray diffraction (Figure 1).<sup>7,8,11</sup> The complex has a typical bent-metallocene geometry and is similar to the structures of Cp<sub>2</sub>Ti(OEt)Cl<sup>12</sup> and Cp<sup>\*</sup><sub>2</sub>Hf(OO<sup>t</sup>Bu)Et.<sup>10</sup> The Ti–O distance in **1** of 1.9090(14) Å is close to the Ti–ethoxide distance of 1.855(2) Å.<sup>12</sup> The peroxide distances of 1.4668(19) Å in **1** and 1.489(12) Å in the hafnium derivative<sup>10</sup> are typical of peroxides.<sup>13</sup> The *tert*-butylperoxo ligand in **1** is bound through only one oxygen, as indicated by the long Ti···O(2) distance (2.952(2) Å) and the open Ti(1)–O(1)–O(2) angle (121.5(1)°). This contrasts with the one other structurally characterized titanium alkylperoxide, which has an η<sup>2</sup>-<sup>t</sup>BuOO ligand.<sup>14</sup>

Complex **1** decomposes in CD<sub>2</sub>Cl<sub>2</sub> at 300 K to give *tert*-butyl alcohol and a number of Cp-containing products in small yield, including Cp<sub>2</sub>TiCl<sub>2</sub> (Table 1). Decay of **1** follows first-order kinetics (by <sup>1</sup>H NMR), proceeding more slowly in cyclohexane (t<sub>1/2</sub> = 2.1 h) and benzene (1.0 h) than in CD<sub>2</sub>Cl<sub>2</sub> (0.5 h). An Eyring plot of rate constants for decomposition from 273 to 313 K in CD<sub>2</sub>Cl<sub>2</sub> gives ΔH<sup>‡</sup> = 27 ± 2 kcal mol<sup>–1</sup> and ΔS<sup>‡</sup> = 15 ± 5 eu. Preliminary



**Figure 1.** ORTEP diagram of **1**, with thermal ellipsoids drawn at 30% probability. Selected data not included in the text: Ti(1)–Cl(1), 2.3956(6) Å; O(1)–Ti(1)–Cl(1), 97.31(4)°; C(11)–O(2)–O(1), 107.65(13)°; Cl(1)–Ti(1)–O(1)–O(2), –79.7(1)°.

**Table 1.** Products and Rate Constants for Reactions of Cp<sub>2</sub>Ti(OO<sup>t</sup>Bu)Cl with PR<sub>3</sub> in CD<sub>2</sub>Cl<sub>2</sub><sup>a</sup>

reaction	R <sub>3</sub> PO	R <sub>2</sub> PO <sup>t</sup> Bu	<sup>t</sup> BuOH <sup>b</sup>	Me <sub>2</sub> CCH <sub>2</sub>	<sup>t</sup> BuCl	Cp <sub>2</sub> TiCl <sub>2</sub>	k <sub>obs</sub> (×10 <sup>–4</sup> s <sup>–1</sup> )
<b>1</b>			96%	n/o	n/o	5%	4.1(2)
<b>1</b> + 1 PPh <sub>3</sub>	98%	n/o	24%	59%	12%	50%	5.9(3)
<b>1</b> + 5 PPh <sub>3</sub>	96%	n/o	29%	56%	11%	48%	7.2(6)
<b>1</b> + 20 PPh <sub>3</sub>	97%	n/o	26%	57%	11%	44%	6.9(5)
<b>1</b> + 20 <sup>t</sup> Bu <sub>3</sub> SnH			98%	n/o	n/d	4%	2.8(8)
<b>1</b> + 1 PPh <sub>3</sub> + 20 <sup>t</sup> Bu <sub>3</sub> SnH	23%	n/o	72%	n/o	n/d	4%	5.2(6)
<b>1</b> + 1 PEt <sub>3</sub>	n/o	~95% <sup>c</sup>	3%	tr	tr	27%	1.6(7)
<b>1</b> + 1 P(OPh) <sub>3</sub>	n/o	50%	28%	8%	3%	28%	3.2(7)
<b>1</b> + CBr <sub>4</sub>			95%	tr	tr	34%	3.5(5)

<sup>a</sup> n/d = not determined; n/o = not observed by <sup>1</sup>H or <sup>31</sup>P{<sup>1</sup>H} NMR; tr = detected in trace amount (<1%). <sup>b</sup> Hydroxyl resonance not observed in <sup>1</sup>H or <sup>2</sup>H NMR. <sup>c</sup> Et<sub>2</sub>PO<sup>t</sup>Bu grows in to a maximum of 70% yield but is concurrently consumed; the yield of Et<sub>2</sub>PO<sup>t</sup>Bu and its apparent decay products is ~95%.

results indicate that cyclohexane and norbornene do not react directly with **1** because the rate of decomposition is unchanged and epoxide products are not observed.


Complex **1** reacts with 1 equiv of PPh<sub>3</sub> in CD<sub>2</sub>Cl<sub>2</sub> to quantitatively form Ph<sub>3</sub>PO (98%), as expected for a peroxide complex.<sup>1</sup> Cp<sub>2</sub>TiCl<sub>2</sub> (50%) and a number of other Cp-containing species are formed, with the total integrated Cp intensity being roughly constant. The <sup>t</sup>Bu groups in **1** are converted to *tert*-butyl alcohol (23%), isobutylene (59%), and <sup>t</sup>BuCl (12%). In contrast, solutions of **1** with 1 equiv of PEt<sub>3</sub> do not form any Et<sub>3</sub>PO (by <sup>31</sup>P{<sup>1</sup>H} NMR). Instead, this reaction yields the phosphinite Et<sub>2</sub>PO<sup>t</sup>Bu, which was identified by <sup>31</sup>P NMR, mass spectrometry, and independent synthesis.<sup>15</sup>

The only reasonable pathway to form Et<sub>2</sub>PO<sup>t</sup>Bu from PEt<sub>3</sub> is by addition of <sup>t</sup>BuO• (eq 3). This reaction is very rapid (k<sub>3</sub> = 1.2 × 10<sup>9</sup> M<sup>–1</sup> s<sup>–1</sup>) and quantitatively forms Et<sub>2</sub>PO<sup>t</sup>Bu.<sup>16</sup> P<sup>n</sup>Bu<sub>3</sub> traps <sup>t</sup>BuO• to give 20% <sup>n</sup>Bu<sub>3</sub>PO and 80% <sup>n</sup>Bu<sub>2</sub>PO<sup>t</sup>Bu (eq 4).<sup>17</sup>



\* To whom correspondence should be addressed. E-mail: mayer@chem.washington.edu.

<sup>†</sup> University of Washington Crystallography Facility.

Scheme 1



Decomposition of **1** in the presence of 1 equiv of  $P^nBu_3$  in  $CD_2Cl_2$  yields this same ratio of phosphine oxide to phosphinite by  $^{31}P$  NMR integration, confirming the intermediacy of  $^tBuO\bullet$  in the decay of **1**.<sup>8</sup> Reaction of **1** with  $P(OPh)_3$  gives the phosphite  $^tBuOP(OPh)_2$  without any phosphate  $(PhO)_3PO$ . Given the known reactivity of  $P(OPh)_3$  with oxyl radicals (eq 5),<sup>18</sup> this shows that peroxy radicals are not present.



The reaction of **1** with  $PPh_3$  is also consistent with the involvement of  $^tBuO\bullet$ , as  $PPh_3 + ^tBuO\bullet$  gives exclusively  $Ph_3PO$  and  $^tBu\bullet$ .<sup>19</sup> The formation of isobutylene and  $^tBuCl$  supports the intermediacy of  $^tBu\bullet$ . As additional confirmation of this pathway,  $^nBu_3SnH$  was used as a competitive trap for  $^tBuO\bullet$ . The yield of  $Ph_3PO$  is reduced from 98 to 23% when 20 equiv of  $^nBu_3SnH$  is added to an equimolar solution of **1** and  $PPh_3$ . This is consistent with  $^nBu_3SnH + ^tBuO\bullet \rightarrow ^nBu_3Sn\bullet + ^tBuOH$  being 9 times slower than  $PPh_3 + ^tBuO\bullet \rightarrow Ph_3PO + ^tBu\bullet$  ( $2.2 \times 10^8$  vs  $1.9 \times 10^9 M^{-1} s^{-1}$ ).<sup>19,20</sup> The lower yield of  $PPh_3$  is not due to direct reaction of **1** and  $^nBu_3SnH$ , as the decay of **1** is not accelerated by  $^nBu_3SnH$  (Table 1).

The rate constant for decay of **1** is only slightly affected by the presence of additives (Table 1). Increasing the concentration of  $PPh_3$  causes small increases in  $k$ , but not linearly, and the presence of  $PEt_3$  or  $^nBu_3SnH$  actually slows the decomposition of **1**. The small changes in  $k$  are most likely a result of the differing stoichiometry of the reactions, changing the amount of **1** that reacts with product radicals such as  $^tBu\bullet$ .

The above results clearly show that  $^tBuO\bullet$  is an intermediate in the decomposition of **1**. This most likely occurs by rate-limiting O—O bond homolysis (Scheme 1). While a complete mechanistic description is hampered by the multiple titanium species formed, homolysis is supported by the positive  $\Delta S^\ddagger$  for decomposition. Other routes to  $^tBuO\bullet$  are difficult to reconcile with the data. Radical chain processes related to the Haber–Weiss mechanism<sup>1a</sup> are unlikely because of the simple first-order kinetic behavior and because decomposition is only marginally slowed by reductive ( $^nBu_3SnH$ ) or oxidative ( $CBR_4$ ) traps. Ti—O bond homolysis<sup>21</sup> to give  $^tBuOO\bullet$  is ruled out by the lack of formation of  $(PhO)_3PO$  or  $Et_3PO$ .<sup>22</sup> Intramolecular induced homolysis, such as attack of a Cp ligand on the peroxide, is unlikely because of the positive  $\Delta S^\ddagger$  and the apparent lack of consumption of Cp ligands.

Because titanium compounds are increasingly used as catalysts for peroxide oxidations,<sup>23</sup> the lack of oxygen atom transfer reactivity of **1** is unexpected. Most likely it is sterically difficult for **1** to adopt the reactive  $\eta^2$ -peroxide conformer. Surprisingly, the “ $Cp_2Ti(O\bullet)Cl$ ” generated by homolysis does not rapidly oxidize  $PEt_3$  to  $Et_3PO$ .

In summary, **1** undergoes O—O bond homolysis at 300 K. This is surprising because the Ti(IV) center is  $d^0$  and cannot be oxidized.

As noted above, all previous clear examples of homolytic cleavage of metal peroxide complexes are facilitated by oxidation of the metal center. It is not clear why decomposition of **1** is so much more facile than homolysis of  $^tBuOO^tBu$ : at 300 K,  $\Delta G^\ddagger = 22 \text{ kcal mol}^{-1}$  for **1** versus  $34 \text{ kcal mol}^{-1}$  for  $^tBuOO^tBu$ .<sup>24</sup> Homolytic decomposition of  $d^0$  peroxides may not be unique to **1**, as an isolated siloxo-ligated  $TiOO^tBu$  complex decomposes to  $^tBuOH$ , and a titanium silasesquioxane complex is reported to react with  $^tBuOOH$  to give  $^tBuOH$  and a trace of  $^tBuOO^tBu$ .<sup>25</sup> Work is continuing both on the chemistry of **1** and on generating more reactive peroxide compounds.

**Acknowledgment.** We thank Dr. Jeffery C. Yoder for valuable discussions and Dr. Joanna Long for assistance with NMR experiments. We are grateful to the National Institutes of Health for financial support.

**Supporting Information Available:** Synthetic, spectroscopic, kinetic, and X-ray crystallographic information for **1** (PDF and CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (a) Sheldon, R. A.; Kochi, J. K. *Metal-Catalyzed Oxidation of Organic Compounds*; Academic Press: New York, 1981. (b) *Organic Peroxides*; Ando, W., Ed.; Wiley & Sons: New York, 1992. (c) *Supplement E2: The Chemistry of Hydroxyl, Ether, and Peroxide Groups*; Patai, S., Ed.; John Wiley & Sons: New York, 1993. (d) Solomon, E. I.; Brunold, T. C.; Davis, M. I.; Kemsley, J. N.; Lee, S.-K.; Lehnert, N.; Neese, F.; Skulan, A. J.; Yang, Y.-S.; Zhou, J. *Chem. Rev.* **2000**, *100*, 235–349. (e) Que, L., Jr.; Ho, R. Y. N. *Chem. Rev.* **1996**, *96*, 2607–2624. (f) Klinman, J. P. *Chem. Rev.* **1996**, *96*, 2541–2561.
- Weissermel, K.; Arpe, H.-J. *Industrial Organic Chemistry*, 3rd ed.; VCH: New York, 1997; pp 267ff, 240ff.
- Katsuki, T.; Sharpless, K. B. *J. Am. Chem. Soc.* **1980**, *102*, 5976–5978.
- Biomimetic Oxidations Catalyzed by Transition Metal Complexes*; Meunier, B., Ed.; Imperial College Press: London, 1998.
- For a contrary view, see: Newcomb, M.; Toy, P. H. *Acc. Chem. Res.* **2000**, *33*, 449–455.
- cf., Lehnert, N.; Ho, R. Y. N.; Que, L., Jr.; Solomon, E. I. *J. Am. Chem. Soc.* **2001**, *123*, 8271–8290.
- Caution:** Solid samples of **1** can be explosive. NMR:  $^1H$  (499 MHz) ( $C_6D_6$ , 20 °C,  $\delta$ ), 5.98 (s, 10H,  $C_5H_5$ ), 1.17 (s, 9H,  $C(CH_3)_3$ ). HR-MS ( $EI^+$ ): calculated 302.05530  $m/z$ ; measured 302.05544. Anal. Calcd: C, 55.62%; H, 6.34%. Found: C, 55.71%; H, 6.33%.
- See Supporting Information for full experimental, synthetic, spectroscopic, and crystallographic data.
- Blackburn, T. F.; Labinger, J. A.; Schwartz, J. *Tetrahedron Lett.* **1975**, *35*, 3041–3044.
- van Asselt, A.; Santarsiero, B. D.; Bercaw, J. E. *J. Am. Chem. Soc.* **1986**, *108*, 8291–8293.  $Cp^* = \eta^5-C_5Me_5$ .
- Crystallographic data for **1**:  $C_{14}H_{19}ClO_2Ti$ ,  $M = 302.64$ , monoclinic,  $P2_1/c$ ,  $a = 6.4150(2) \text{ \AA}$ ,  $b = 12.0410(5) \text{ \AA}$ ,  $c = 19.0220(10) \text{ \AA}$ ,  $\beta = 108.7020(14)^\circ$ ,  $V = 1391.74(10) \text{ \AA}^3$ ,  $Z = 4$ ,  $\rho_c = 1.444 \text{ g cm}^{-3}$ ,  $\mu = 0.798 \text{ mm}^{-1}$ ,  $T = 130(2) \text{ K}$ . 5103 reflections collected, 3017 unique ( $R_{int} = 0.0447$ ). Refinement on all data converged at  $R1 = 0.039$ ,  $wR2 = 0.108$ .
- Huffman, J. C.; Moloy, K. G.; Marsella, J. A.; Caulton, K. G. *J. Am. Chem. Soc.* **1980**, *102*, 3009–3014.
- Reference 1c, p 150.
- Boche, G.; Möbus, K.; Harms, K.; Marsch, M. *J. Am. Chem. Soc.* **1996**, *118*, 2770–2771.
- $Et_2PCl + ^tBuOH + Et_3N$  in  $C_6H_6 \rightarrow Et_2PO^tBu$ ;  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  109.6 ppm.<sup>8</sup> MS ( $EI^+$ ): 162  $m/z$ .
- Davies, A. G.; Dennis, R. W.; Griller, D.; Roberts, B. P. *J. Organomet. Chem.* **1972**, *40*, C33–C35.
- Chatgililoglu, C.; Ingold, K. U.; Scaiano, J. C. *J. Am. Chem. Soc.* **1981**, *103*, 7739–7742.
- (a) Kochi, J. K.; Krusic, P. J. *J. Am. Chem. Soc.* **1969**, *91*, 3944–6. (b) Pobedimskii, D. G.; Kirpichnikov, P. A. *J. Polym. Sci., Polym. Chem. Ed.* **1980**, *18*, 815–825.
- Griller, D.; Ingold, K. U.; Patterson, L. K.; Scaiano, J. C.; Small, R. D., Jr. *J. Am. Chem. Soc.* **1979**, *101*, 3780–3785.
- Scaiano, J. C. *J. Am. Chem. Soc.* **1980**, *102*, 5399–5400.
- $Cp_2(Cl)Ti(TEMPO)$  undergoes Ti—O homolysis at 333 K: Huang, K. W.; Waymouth, R. M. *J. Am. Chem. Soc.* **2002**, *124*, 8200–8201.
- Buckler, S. A. *J. Am. Chem. Soc.* **1962**, *84*, 3093–3097.
- For recent examples and entries into this literature, see ref 25.
- Reference 1b, p 176.
- (a) Fugiwara, M.; Wessel, H.; Hyung-Suh, P.; Roesky, H. W. *Tetrahedron* **2002**, *58*, 239–242. (b) Crocker, M.; Herold, R. H. M.; Orpen, A. G.; Overgaard, M. T. A. *J. Chem. Soc., Dalton Trans.* **1999**, 3791–3804.

JA028500A