


Gold(I)-Catalyzed Intermolecular Addition of Phenols and Carboxylic Acids to Olefins

Cai-Guang Yang, and Chuan He

J. Am. Chem. Soc., **2005**, 127 (19), 6966-6967 • DOI: 10.1021/ja050392f • Publication Date (Web): 22 April 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 40 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Gold(I)-Catalyzed Intermolecular Addition of Phenols and Carboxylic Acids to Olefins

Cai-Guang Yang and Chuan He*


Department of Chemistry, The University of Chicago, 5735 South Ellis Avenue, Chicago, Illinois 60637

Received January 20, 2005; E-mail: chuanhe@uchicago.edu

Gold-catalyzed reactions have emerged as important synthetic methods.¹ Cationic gold(I) and gold(III) show exceptional activities to activate alkynes toward addition by a variety of functional groups both intra- and intermolecularly. This chemistry is also very valuable in constructing complex molecules.^{1,2} Despite the successes with alkynes, reactions involving nucleophilic addition to olefins catalyzed by gold are very limited.^{3,4} Recently, an interesting gold(III)-mediated addition of β -diketone to alkenes was shown, but the role of gold(III) was thought to mainly activate the nucleophile and the alkene scope is limited to styrene and norbornylene.⁴ To this date, there has not been a report of gold-catalyzed formation of carbon–heteroatom bonds through activation of inert olefins. Such reactions have been traditionally mediated by acids or stoichiometric amounts of toxic reagents.⁵ Platinum,⁶ ruthenium,⁷ and palladium⁸ have been shown to catalyze some of these reactions, but efficient intermolecular addition of various nucleophiles to inactivated olefins under mild conditions has yet to be achieved. In the case of palladium-catalyzed reactions, β -hydride elimination often occurs to afford unsaturated products.⁸

We report here that simple olefins can be activated by Ph_3PAuOTf .⁹ Intermolecular additions of phenols and carboxylic acids to alkenes can be catalyzed by this catalyst under relatively mild conditions. A reaction between *p*-methoxyphenol (1 mmol) and 4-allylanisole (4 mmol) was used to investigate activity of different catalysts (eq 1). Triflic acid, ZnOTf_2 , AgOTf , and $\text{AuCl}_3/3\text{AgOTf}$ all failed to give the desired product in good yields (Table 1). A combination of 2 mol % of Ph_3PAuCl and AgOTf gave a

Table 1. Effect of Catalysts on an Intermolecular Addition Reaction



entry	catalyst	yield ^a
1	15% HOTf	0%
2	5% ZnOTf_2	0%
3	5% AgOTf	trace
4	5% $\text{AuCl}_3/15\%$ AgOTf	<5%
5	2% $\text{Ph}_3\text{PAuCl}/2\%$ AgOTf	84%
6	2% Ph_3PAuOTf	78%

^a Yield based on the phenol and determined by crude ¹H NMR using an internal standard.

good yield of the Markovnikov product (entry 5, Table 1). If AgCl precipitates were filtered after mixing the same equivalents of Ph_3PAuCl and AgOTf , the filtrate still exhibited the same catalytic activity (entry 6, Table 1). Different solvents were screened, and toluene was found to be the best solvent for this reaction.

Both electron-rich and electron-deficient phenols serve as good substrates (Table 2). Different olefins including inactivated ones




Table 2. Intermolecular Addition of Phenols and Carboxylic Acids to Olefins^b

Entry	Nucleophile	Olefin	Product	Yield ^a
1	1a R = MeO	2a	3a R = MeO	78%
2	1b R = <i>tert</i> -Butyl	2a	3b R = <i>tert</i> -Butyl	70% ^b
3	1c R = CO ₂ Me	2a	3c R = CO ₂ Me	68%
4	1d R = NO ₂	2a	3d R = NO ₂	71% ^c
5	1a	2b	3e R = MeO	58%
6	1c	2b	3f R = CO ₂ Me	80%
7	1d	2b	3g R = NO ₂	81% ^c
8	1a	2c	3h R = MeO	3h' 71% ^d
9	1e R = Br	2c	3i R = Br	3i' 85% (7:1) ^b
10	1f R = H	2c	3j R = H	3j' 80% (15:1) ^b
11	1g BnCOOH	2a	3k	84% ^c
12	1g	2b	3l	95% ^c
13	1g	2c	3m 3m'	78% (10:1) ^{c,e}
14	1g	2d	3n 3n'	75% (5.5:1) ^c
15	1h	2b	3o	46% ^{c,f}
16	1i	2e	3p	>95% ^{c,g}

^a The isolated yields are reported. ^b NMR yield with an internal standard. ^c With 5 mol % of the catalyst. ^d Isolated yield for **3h**. ^e 5 mol % $\text{Ph}_3\text{PAuSbF}_6$ was used. ^f Conversion is 55% based on the alkene. ^g 1:1 ratio of acid and norbornylene were used. ^h Reactions were conducted with 1 mmol of nucleophiles, 4 mmol of olefins, and 2 mol % (for phenols) or 5 mol % (for carboxylic acids) of $\text{Ph}_3\text{PAuCl}/\text{AgOTf}$ in 2 mL of toluene at 85 °C.

work well in this reaction. Good yields can be obtained for all cases reported. Although excess amounts of olefins were used (4 equiv), usually 2 equiv of olefins could be recovered after the reaction. Carboxylic acids can also be added to alkenes to give esters under the same reaction conditions (Table 2), but a higher catalyst loading (5 mol %) is required for the completion of the reactions. A sterically bulky acid was employed, which still afforded a reasonable yield of the product (entry 15, Table 2).

We are exploring the scope of this gold(I)-based activity. Our data indicate that an intramolecular cyclization of a γ -hydroxyl

Scheme 1. Intramolecular Addition of Alcohol to Olefin**Scheme 2.** Terminal Olefin Migration Catalyzed by Au(I)**Scheme 3** Proposed Catalytic Mechanism

alkene can be mediated by gold(I) under the same conditions (Scheme 1). The product yield is comparable to that reported recently with a platinum-based system.^{6a} This type of intramolecular addition of a hydroxyl group to an alkene was suggested as a potential step in a gold(III)-catalyzed tandem reaction previously;^{3a} however, this has never been demonstrated experimentally. Preliminary results also indicate that acidic alcohols serve as good nucleophiles to add intermolecularly into olefins.

While investigating these reactions, we discovered that migration of double bonds occurred in some cases (entries 8–10, 13, and 14 in Table 2). To confirm the observation, we took 4-phenyl-1-butene and heated it at 85 °C for 20 h in the presence of 1 mol % of Ph_3PAuOTf without any nucleophiles in toluene. Seventy-five percent of the alkene was converted to 4-phenyl-2-butene in a 2.2:1 E/Z ratio (Scheme 2). Further migration to the conjugated system was not observed; the same phenomenon was also discovered with a ruthenium-based system.¹⁰

The reaction mechanism is proposed in Scheme 3. We believe the cationic gold(I) binds and activates alkene for a nucleophilic addition by the phenols or carboxylic acids,¹¹ a reaction similar to the Wacker process catalyzed by palladium(II).¹² A subsequent proton-transfer step affords the final product and regenerates gold(I) catalyst. The gold catalyst also promotes migration of double bonds, which gives rise to formation of small amounts of side products for some substrates (Scheme 3). The mechanism of the double bond migration mediated by gold(I) is unclear at this moment.

In summary, we report here the gold(I)-catalyzed intermolecular addition of phenols and carboxylic acids to olefins. The reaction is simple and runs under relatively mild conditions. To our knowledge, this is the first example of a gold(I)-mediated activation of inert alkenes toward nucleophilic addition. Experimental results support a proposed mechanism with the gold(I) directly activating the olefin. This study may open a new direction for alkene functionalization.

Acknowledgment. This research was supported by the University of Chicago. We acknowledge the donors of the American Chemical Society Petroleum Research Fund for support of this research (PRF 38848-G3). This research is also supported by a Research Innovation Award (RI1179) from Research Corporation.

Supporting Information Available: Experimental details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For selected reviews, see: (a) Hashmi, A. S. K. *Gold Bull.* **2004**, *37*, 51–65. (b) Hashmi, A. S. K. *Gold Bull.* **2003**, *36*, 3–9. (c) Dyker, G. *Angew. Chem., Int. Ed.* **2000**, *39*, 4237–4239. Selected examples: (d) Ito, Y.; Sawamura, M.; Hayashi, T. *J. Am. Chem. Soc.* **1986**, *108*, 6405–6406. (e) Teles, J. H.; Brode, S.; Chabanas, M. *Angew. Chem., Int. Ed.* **1998**, *37*, 1415–1418. (f) Fukuda, Y.; Utimoto, K. *J. Org. Chem.* **1991**, *56*, 3729–3731. (g) Asao, N.; Takahashi, K.; Lee, S.; Kasahara, T.; Yamamoto, Y. *J. Am. Chem. Soc.* **2002**, *124*, 12650–12651.
- (2) For selected more recent examples, see: (a) Nieto-Oberhuber, C.; Muñoz, M. P.; Buñuel, E.; Nevado, C.; Cárdenas, D. J.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2004**, *43*, 2402–2406. (b) Arcadi, A.; Bianchi, G.; Marinelli, F. *Synthesis* **2004**, 610–618. (c) Mamane, V.; Gress, T.; Krause, H.; Fürstner, A. *J. Am. Chem. Soc.* **2004**, *126*, 8654–8655. (d) Sherry, B. D.; Toste, F. D. *J. Am. Chem. Soc.* **2004**, *126*, 15978–15979. (e) Staben, S. T.; Kennedy-Smith, J. J.; Toste, F. D. *Angew. Chem., Int. Ed.* **2004**, *43*, 5350–5352. (f) Yao, T.; Zhang, X.; Larock, R. C. *J. Am. Chem. Soc.* **2004**, *126*, 11164–11165. (g) Zhang, L.; Kozmin, S. A. *J. Am. Chem. Soc.* **2004**, *126*, 11806–11807. (h) Hashmi, A. S. K.; Weyrauch, J. P.; Frey, W.; Bats, J. W. *Org. Lett.* **2004**, *6*, 4391–4394. (i) Morita, N.; Krause, N. *Org. Lett.* **2004**, *6*, 4121–4123. (j) Hashmi, A. S. K.; Weyrauch, J. P.; Rudolph, M.; Kurpejović, E. *Angew. Chem., Int. Ed.* **2004**, *43*, 6545–6547. (k) Straub, B. F. *Chem. Commun.* **2004**, 1726–1728. (l) Fürstner, A.; Hannen, P. *Chem. Commun.* **2004**, 2546–2547. (m) Milton, M. D.; Inada, Y.; Nishibayashi, Y.; Uemura, S. *Chem. Commun.* **2004**, 2712–2713.
- (3) (a) Intramolecular addition of a hydroxyl group to an alkene was proposed as part of a tandem reaction. See: Hashmi, A. S. K.; Schwarz, L.; Choi, J.-H.; Frost, T. M. *Angew. Chem., Int. Ed.* **2000**, *39*, 2285–2288. (b) Kobayashi, S.; Kakumoto, K.; Sugiura, M. *Org. Lett.* **2002**, *4*, 1319–1322.
- (4) Yao, X.; Li, C.-J. *J. Am. Chem. Soc.* **2004**, *126*, 6884–6885.
- (5) (a) Larock, R. C.; Leong, W. W. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon Press: New York, 1991; Vol. 4, p 297. (b) Robin, S.; Rousseau, G. *Tetrahedron* **1998**, *54*, 13681–13736. (c) Harding, K. E.; Tinger, T. H. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon Press: New York, 1991; Vol. 4, p 463. (d) Mulzer, J. In *Organic Synthesis Highlights*; Mulzer, J., Altenbach, H. J., Braun, M., Krohn, K., Reissing, H. U., Eds.; VCH: Weinheim, Germany, 1991; p 157. (e) Larock, R. C. *Solvolmercuriation/Demercuration Reactions in Organic Synthesis*; Springer-Verlag: Berlin, 1986; Chapter 3.
- (6) (a) Qian, H.; Han, X.; Widenhoefer, R. A. *J. Am. Chem. Soc.* **2004**, *126*, 9536–9537. (b) Wang, X.; Widenhoefer, R. A. *Chem. Commun.* **2004**, 660–661. (c) Liu, C.; Han, X.; Wang, X.; Widenhoefer, R. A. *J. Am. Chem. Soc.* **2004**, *126*, 3700–3701. (d) Wang, X.; Widenhoefer, R. A. *Organometallics* **2004**, *23*, 1649–1651.
- (7) (a) Oe, Y.; Ohta, T.; Ito, Y. *Chem. Commun.* **2004**, 1620–1621. (b) Oe, Y.; Ohta, T.; Ito, Y. *Synlett* **2005**, 179–181.
- (8) For a recent review and selected examples, see: (a) Stahl, S. S. *Angew. Chem., Int. Ed.* **2004**, *43*, 3400–3420. (b) Utsunomiya, M.; Kawatsura, M.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2003**, *42*, 5865–5868. (c) Arai, M. A.; Kuraishi, M.; Arai, T.; Sasai, H. *J. Am. Chem. Soc.* **2001**, *123*, 2907–2908. (d) Trend, R. M.; Ramtohl, Y. K.; Ferreira, E. M.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2003**, *42*, 2892–2895. (e) Uozumi, Y.; Kato, K.; Hayashi, T. *J. Am. Chem. Soc.* **1997**, *119*, 5063–5064. (f) Larock, R. C.; Hightower, T. R. *J. Org. Chem.* **1993**, *58*, 5298–5300. (g) Hegedus, L. S.; Allen, G. F.; Bozell, J. J.; Waterman, E. L. *J. Am. Chem. Soc.* **1978**, *100*, 5800–5807.
- (9) (a) Komiya, S.; Kochi, J. K. *J. Organomet. Chem.* **1977**, *135*, 65–72. (b) Canales, S.; Crespo, O.; Gimeno, M. C.; Jones, P. G.; Laguna, A.; Mendizabal, F. *Organometallics* **2000**, *19*, 4985–4994.
- (10) Arisawa, M.; Terada, Y.; Nakagawa, M.; Nishida, A. *Angew. Chem., Int. Ed.* **2002**, *41*, 4732–4734.
- (11) For selected gold-olefin complexes, see: (a) Huettel, R.; Reinheimer, H.; Dietl, H. *Chem. Ber.* **1966**, *99*, 2778–2781. (b) Dávila, R. M.; Staples, R. J.; Fackler, J. P., Jr. *Organometallics* **1994**, *13*, 418–420. (c) Cinelli, M. A.; Minghetti, G.; Stoccoro, S.; Zucca, A.; Manassero, M. *Chem. Commun.* **2004**, 1618–1619.
- (12) For examples for the Wacker process, see: Bäckvall, J. E.; Åkermark, B.; Ljunggren, S. O. *J. Am. Chem. Soc.* **1979**, *101*, 2411–2416 and references therein.

JA050392F