


Cobalt-Catalyzed Hydroazidation of Olefins: Convenient Access to Alkyl Azides

Jrme Waser, Hisanori Nambu, and Erick M. Carreira

J. Am. Chem. Soc., **2005**, 127 (23), 8294-8295 • DOI: 10.1021/ja052164r • Publication Date (Web): 21 May 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 6 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)




Cobalt-Catalyzed Hydroazidation of Olefins: Convenient Access to Alkyl Azides

Jérôme Waser, Hisanori Nambu, and Erick M. Carreira*

Laboratorium für Organische Chemie, ETH Hönggerberg, CH-8093 Zürich, Switzerland

Received April 5, 2005; E-mail: carreira@org.chem.ethz.ch


Amines and related functional groups are ubiquitous in natural products as well as pharmaceutical substances. Consequently, the development of methods for the introduction of nitrogen in simple organic compounds is an intense focus of research.¹ Azides are easily converted into either amines^{2a} or pharmaceutically relevant heterocycles.^{2b,c} Furthermore, they have recently generated much interest due to their stability under physiological conditions and their unique reactivity, which permits their use in bioconjugation via Staudinger ligation^{2d} or click chemistry.^{2e} The most common access route to alkyl azides involves the substitution reaction of primary or secondary alkyl halides with inorganic azides.³ The direct reaction of alkenes with hydrazoic acid sources is limited to alkenes that give rise to stabilized carbocations and require excess HN₃,^{4a} TMSN₃,^{4b} or zeolite-supported NaN₃,^{4c} unactivated monosubstituted olefins being described as unreactive in these processes.⁴ Herein we report the functionalization reaction of unactivated olefins to the corresponding azides with high Markovnikov selectivity using tosyl azide (TsN₃) and a silane. The reaction is mediated by a simple cobalt catalyst prepared in situ from Schiff base **1** and Co(BF₄)₂·6H₂O (eq 1).



We have recently documented the Co- and Mn-catalyzed hydrohydrazination reaction of alkenes (RO₂CN=NCO₂R + alkene → *N*-alkyl hydrazine).⁵ This reaction is the nitrogen equivalent of the olefin hydration reaction developed by Mukaiyama and Isayama^{6a} and displays broad substrate scope. In our ongoing search for other nitrogen sources that would be amenable for use in olefin functionalization processes, we have examined sulfonyl azides, as they are convenient, readily available, and widely handled reagents in the laboratory.⁷ Moreover, we speculated that sulfonyl azides could display parallel reactivity to that observed with azodicarboxylates in the cobalt- and manganese-mediated processes.

Given our prior success with Mn(dpm)₃⁸ and cobalt complexes in the hydrohydrazination reaction, we first examined their use together with 1.6 equiv of phenylsilane and 3 equiv of ethyl sulfonyl azide for the hydroazidation of 4-phenylbutene (**4**) as a test substrate. The manganese complex failed to provide azide product, and the cobalt complex obtained from Co(NO₃)₃·6H₂O, 2-aminoisobutyric acid, salicylaldehyde, NaOH, and H₂O₂ in ethanol⁹ afforded 4-phenyl-2-azidobutane in 50% yield after 48 h. Further extensive optimization of the reaction conditions did not lead to any improvements. Knowing that the catalyst used still contains impurities, we re-examined the preparation of the complex, with the hope that greater purity would result in increased reactivity. Using Co(OH)₃ as catalyst precursor,⁹ we succeeded in isolating

Chart 1



complex **5** (Chart 1), which we thought to be the active catalyst for the hydrohydrazination reaction; however, this complex was inactive in olefin hydroazidation. Analysis of the mass spectrum revealed complex **6** to be present in the mixture obtained via the in situ oxidation procedure en route to **5**. Complex **6** could be prepared by refluxing a 1:1:1 mixture of salicylaldehyde, 2-aminoisobutyric acid, and Co(OAc)₂·4H₂O in water under argon, collecting the precipitating solid and oxidizing it in air in the presence of 1 equiv of 2-amino-isobutyric acid in ethanol. The activity of the complex obtained was comparable to that observed with the mixture isolated using our original protocol.

The identification of a cobalt complex incorporating a single tridentate ligand in the active catalyst presented new opportunities for catalyst design. We speculated that increasing the steric bulk around the tridentate ligand could lead to a complex with enhanced stability and perhaps greater activity. With this design strategy in mind, **7** was synthesized. Preformation of Schiff base **1** derived from 3,5-di-*tert*-butyl salicylaldehyde and α,α-diphenyl glycine followed by reaction with Co(NO₃)₃·6H₂O and oxidation with H₂O₂ proved crucial to the isolation of **7**. Interestingly, coordination of a second molecule of α,α-diphenyl glycine to the cobalt center was never observed by NMR and mass spectroscopy. Nonetheless, we observed that **7** was indeed a better catalyst, converting 4-phenylbutene (**4**) to the desired azide in 12 h in 60% yield. Further investigations revealed a batch-to-batch dependency with some catalyst preparations showing lower reactivity and long (up to 12 h) initiation period. Inspired by Mukaiyama's work in olefin hydration,⁶ we used 30% *t*-BuOOH as an additive to facilitate initiation of the reaction. We were gratified to observe full conversion to product in 2–8 h under otherwise identical conditions. In the optimal procedure, in situ preparation of the catalyst from Co(BF₄)₂·6H₂O and **1** gave 70% of the desired azide in 1.5 h, allowing reproducible yields and reaction times.

Analysis of the reaction mixture led to the identification of 4-phenylbutane as the major side product (28% yield). At this point, we decided to vary the structure of both silane and sulfonyl azide, hoping to increase the ratio of products in favor of the alkyl azide. Tosyl azide proved to be best, increasing the azide/alkane ratio from 3:1 to 9:1 (as determined by GC). Finally, using tetramethyldisiloxane (TMDSO) and tosyl azide, we found that the ratio was increased to 20:1, albeit with slight attenuation of reactivity (reaction time 3 h instead of 2 h). Using the optimized procedure, we routinely obtained 4-phenyl-2-azido-butane in 85–90% yield.

With a reliable procedure in hand, we proceed to examine the scope of the hydroazidation reaction, with both phenylsilane and

Table 1. Hydroazidation Reaction of Olefins (Eq 1)


| Entry | Alkene | Product | Yield ^{a)} with PhSiH ₃ | Yield ^{b)} with TMDSO |
|-------|--------|---------|---|--------------------------------------|
| 1 | | | 90% | 86% |
| 2 | | | 72% | 69% |
| 3 | | | 65% | 62% |
| 4 | | | 73% | 85% ^{c)} |
| 5 | | | 55% | 67% |
| 6 | | | 35% | 39% ^{c)} |
| 7 | | | 75% | 77% |
| 8 | | | 49% ^{d)} | 46% ^{d)} |
| 9 | | | 86% | 90% |
| 10 | | | 89% dr: 4:1 | 76% ^{c)} dr: 4:1 |
| 11 | | | 73% ^{d)} | 58% ^{c),d)} |
| 12 | | | 63% ^{d)} | 48% ^{c),d)} |
| 13 | | | 66% | 48% ^{c),d)} |

^a General procedure A: 0.5 mmol alkene, 0.8 mmol PhSiH₃, 1.5 mmol TsN₃, 30 mol % *t*-BuOOH, 6 mol % ligand **1**, 6 mol % Co(BF₄)₂·6H₂O, 2.5 mL of ethanol at 23 °C under argon. ^b General procedure B: 1.0 mmol TMDSO was used instead of PhSiH₃. ^c 2.0 mmol TMDSO were used. ^d Starting material was partially recovered (see Supporting Information for further details).

TMDSO (Table 1). All the terminal olefins tested (entries 1–8) showed excellent Markovnikov selectivity. TMDSO is generally better for this class of olefins as less alkane side products are formed; when this observation is coupled with the fact that it is also inexpensive, it is rendered the more convenient reagent for the reaction. An aromatic ring in allylic or homoallylic position was well-tolerated (entries 1–3), but styrene derivatives (not shown) were not active, although they had proven to be excellent substrates for the related hydrohydrazination reaction. Homoallylic and allylic ethers were also good substrates (entries 4–6), provided a bulky silyl protecting group was used. The free alcohols proved to be inactive. Esters and ketones were tolerated (entries 7 and 8). Chemoselectivity was excellent, as no reduction of the ketone was observed. Geminally disubstituted and trisubstituted olefins were also good substrates for this reaction, furnishing the tertiary azide exclusively (entries 9–13). For these substrates, phenylsilyl proved to be the reductant of choice, due to its higher reactivity.

The mild conditions of the olefin hydroazidation reaction permitted us to examine various processes that effect the conversion of the product to other useful compounds without the need of isolation and purification of the azides themselves. We have observed that the one-pot conversions of olefins to amines using *in situ* reduction^{2a} (eq 2) and to 1,4-disubstituted triazoles using

Sharpless' procedure^{2b} (eq 3) were feasible in useful yields for 4-phenylbutene (**4**) as the prototypical substrate. In the reduction reaction, a simple extraction procedure sufficed to allow isolation of the free amine in 76% yield and 95% purity, as determined by NMR.



We have documented the cobalt-catalyzed hydroazidation reaction as a new entry to alkyl azides directly from unactivated olefins. The method complements existing approaches based on substitution of alkyl halides or hydrazoic acid addition to activated olefins. In addition to the azides that can be generated, the reaction can be coupled to processes such as reduction to give rise to amines and cycloaddition giving useful heterocycles. Further studies are being conducted to better understand the similarities and differences between this process and the hydrohydrazination reaction, with the aim of discovering more efficient and versatile catalytic systems.

Acknowledgment. We thank the Swiss National Foundation for support of this research and the Roche Research Foundation for the support of J.W. H.N. thanks the Japan Society for the Promotion of Science (JSPS) for a postdoctoral fellowship.

Supporting Information Available: Experimental procedures and spectral data for all products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) Brown, B. R. *The Organic Chemistry of Aliphatic Nitrogen Compounds*; Oxford University Press: New York, 1994. (b) Ricci, A. *Modern Amination Methods*; Wiley-VCH: Weinheim, Germany, 2000. (c) Roesky, P. W.; Muller, T. E. *Angew. Chem., Int. Ed.* **2003**, *42*, 2708. (d) Espino, C. G.; Fiori, K. W.; Kim, M.; Du Bois, J. *J. Am. Chem. Soc.* **2004**, *126*, 15378. (e) Utsunomiya, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2003**, *125*, 14286. (f) Brice, J. L.; Harang, J. E.; Timokhin, V. I.; Anastasi, N. R.; Stahl, S. S. *J. Am. Chem. Soc.* **2005**, *127*, 9, 2868.
- (a) Rao, H. S. P.; Siva, P. *Synth. Commun.* **1994**, *24*, 549. (b) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596. (c) Demko, Z. P.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2110. (d) Kohn, M.; Breinbauer, R. *Angew. Chem., Int. Ed.* **2004**, *43*, 3106. (e) Kolb, H. C.; Sharpless, K. B. *Drug Discovery Today* **2003**, *8*, 1128.
- (a) Scriven, E. F. V.; Turnbull, K. *Chem. Rev.* **1988**, *88*, 297. For an approach using radicals, see: (b) Ollivier, C.; Renaud, P. *J. Am. Chem. Soc.* **2001**, *123*, 4717.
- (a) Hassner, A.; Fibiger, R.; Andisik, D. *J. Org. Chem.* **1984**, *49*, 4237. (b) Breton, G. W.; Daus, K. A.; Kropp, P. *J. Org. Chem.* **1992**, *57*, 6646. (c) Sreekumar, R.; Padmakumar, R.; Rugmini, P. *Chem. Commun.* **1997**, 1133. For silyl enolate azidation, see: (d) Magnus, P.; Lacour, J.; Evans, P. A.; Roe, M. B.; Hulme, C. *J. Am. Chem. Soc.* **1996**, *118*, 3406. For olefin iodoazidation, see: (e) Hassner, A.; Levy, L. A. *J. Am. Chem. Soc.* **1965**, *87*, 4203. For diazidation, see: (f) Fristad, W. E.; Brandvold, T. A.; Peterson, J. R.; Thompson, S. R. *J. Org. Chem.* **1985**, *50*, 3647. (g) Snider, B. B.; Lin, H. *Synth. Commun.* **1998**, *28*, 1913. For olefin selenoazidation, see: (h) Tiecco, M.; Testaferri, L.; Sand, C.; Tomassini, C.; Marini, F.; Bagnoli, L.; Temperini, A. *Angew. Chem., Int. Ed.* **2003**, *42*, 3131. For conjugate addition to unsaturated carbonyl compounds, see: (i) Myers, J. K.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1999**, *121*, 8959. (j) Horstmann, T. E.; Guerin, D. J.; Miller, S. *J. Angew. Chem., Int. Ed.* **2000**, *39*, 3635. For olefin aziridination using azides, see: (k) Mahoney, J. M.; Smith, C. R.; Johnston, J. N. *J. Am. Chem. Soc.* **2005**, *127*, 1354.
- (a) Waser, J.; Carreira, E. M. *J. Am. Chem. Soc.* **2004**, *126*, 5676. (b) Waser, J.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2004**, *43*, 4099.
- (a) Isayama, S.; Mukaiyama, T. *Chem. Lett.* **1989**, 1071. For a discussion of the effect of peroxides on the reaction rate, see: (b) Tokuyasu, T.; Kunikawa, S.; Masuyama, A.; Nojima, M. *Org. Lett.* **2002**, *4*, 3595.
- El-Sayed, R. A. *Phosphorus, Sulfur Silicon Relat. Elem.* **2004**, *179*, 237.
- dpm = dipivaloylmethanato.
- Burrows, R. C.; Bailar, J. C. *J. Am. Chem. Soc.* **1966**, *88*, 4150.

JA052164R