


Communication

Redox Couples of Inducible Nitric Oxide Synthase

Andrew K. Udit, Wendy Belliston-Bittner, Edith C. Glazer, Yen Hoang Le Nguyen, James M. Gillan, Michael G. Hill, Michael A. Marletta, David B. Goodin, and Harry B. Gray

J. Am. Chem. Soc., **2005**, 127 (32), 11212-11213 • DOI: 10.1021/ja0520114 • Publication Date (Web): 20 July 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 3 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Redox Couples of Inducible Nitric Oxide Synthase

Andrew K. Udit,[†] Wendy Belliston-Bittner,[‡] Edith C. Glazer,^{‡,||} Yen Hoang Le Nguyen,[‡]
James M. Gillan,[†] Michael G. Hill,[†] Michael A. Marletta,[§] David B. Goodin,^{||} and Harry B. Gray*[‡]

Department of Chemistry, Occidental College, Los Angeles, California 90041, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, The Scripps Research Institute, La Jolla, California 92037, and Department of Chemistry, University of California, Berkeley, California 94720

Received March 29, 2005; E-mail: hbgray@caltech.edu

The critical role of nitric oxide (NO) in cellular signaling is firmly established.¹ In higher animals, NO is synthesized by nitric oxide synthase (NOS), which converts L-arginine to citrulline and NO with NADPH and O₂ as co-substrates.² The enzyme consists of a reductase domain where electrons from NADPH are shuttled through bound flavins FAD and FMN, a calmodulin binding region that controls electron transfer (ET) to the heme, and a heme domain that contains a cysteine-ligated heme and tetrahydrobiopterin (H₄B). Investigations herein utilize the heme domain of inducible NOS (iNOS), an immune system isoform implicated in several diseases in humans.³

The complexity of NOS and its impact on human health have thrust work on the NOS mechanism into the spotlight. It is well established that the catalytic cycle involves two turnovers of the enzyme. The first turnover converts L-arginine to *N*-hydroxyarginine; although envisioned as a P450-like hydroxylation, the reaction is dependent on one-electron oxidation of H₄B.⁴ The second turnover is thought to involve a ferric peroxide nucleophile, but H₄B is also required for this turnover in an as-yet undetermined role. Although many details of the NOS mechanism remain to be elucidated, it is certain that ET reactions are key steps in the catalytic cycle.

Direct electrochemistry of iNOS can be achieved by confining the protein in didodecyltrimethylammonium bromide (DDAB) films on the surface of basal plane graphite electrodes (BPG).^{5–7} We report reduction potentials for Fe^{III/II} and Fe^{II/I} couples, Fe^{III/II} ET kinetics, catalytic reduction of dioxygen, and evidence for water-free and water-bound forms of the iNOS heme based on scan rate and pH dependence data.

DDAB films were formed on BPG (0.07 cm²) by depositing 5 μ L of a 10 mM aqueous solution of DDAB on the electrode surface, followed by slow drying overnight. iNOS was incorporated into the film by soaking the coated electrode in a solution of enzyme (\sim 20 μ M in 50 mM KP_i, 50 mM KCl, pH 7 buffer) for 30 min. A voltammogram of iNOS in DDAB on BPG is shown in Figure 1. We have assigned *E*₁ (–191 mV) and *E*₂ (–1049 mV) to heme Fe^{III/II} and Fe^{II/I} couples, respectively, consistent with other studies of heme proteins in DDAB films.^{8,9} Notably, a couple similar to *E*₂ was observed for neuronal nitric oxide synthase (nNOS), but was not assigned.⁷ Our assignments of *E*₁ and *E*₂ are supported by voltammetry in the presence of carbon monoxide; *E*₁ shifts approximately +62 mV (consistent with other studies^{10,11}), while *E*₂ is not observed (presumably beyond the solvent window). For comparison, redox titrations of iNOS in its resting state (six-coordinate heme, low spin) conducted in solution yield a potential of –544 mV (vs Ag/AgCl) for Fe^{III/II}.¹² Empirically, potential shifts




Figure 1. Cyclic voltammogram of iNOS in DDAB on BPG (0.07 cm²) at 200 mV/s in 50 mM KP_i/50 mM KCl/pH7.




Figure 2. Cyclic voltammograms of iNOS in DDAB on BPG in 50 mM KP_i/50 mM KCl (a) at 0.05 (black), 1 (gray), and 8 (dashed) V/s, pH 7; (b) at 1 V/s, pH 5; (c) at 1 V/s, pH 9. (d) Variation of the average Fe^{III/II} midpoint potential with pH at 200 mV/s.

are usually observed for heme proteins in surfactant films;¹³ the hydrophobic film likely leads to partial heme dehydration, causing the positive potential shift.¹⁴

Besides CO, other molecules in solution also bind the heme. In the presence of 500 mM imidazole, *E*₁ shifts +20 mV. When dioxygen is added, large catalytic reduction currents at the onset of *E*₁ are observed (Supporting Information, Figure S1).

For *E*₁, the peak current is linear with scan rate (surface bound) up to 16.7 V/s, after which it is linear with the square root of the scan rate (diffusive).¹⁵ This behavior is characteristic of thin-film electrochemistry and indicates finite diffusion of the protein within the film.¹⁶ Thus, up to 16.7 V/s, we treated the redox system as surface-confined.

Closer inspection of Figure 1 reveals that the *E*₁ cathodic wave is slightly broader than the other waves. Voltammetry at variable scan rates allowed us to resolve this peak into two distinct cathodic processes, *E*_{p,c}(1) and *E*_{p,c}(2). Figure 2a shows voltammograms recorded in pH 7 buffer at different scan rates. At high and low scan rates (8 vs 0.05 V/s), only one cathodic peak (*E*_{p,c}(1) vs *E*_{p,c}(2)) is present, while both cathodic processes are observed at intermedi-

[†] Occidental College.

[‡] California Institute of Technology.

^{||} The Scripps Research Institute.


[§] University of California, Berkeley.

ate (1 V/s) scan rates. In addition to being scan rate dependent, these cathodic processes are also pH dependent. At pH 5 and 1 V/s (Figure 2b), only one cathodic peak ($E_{p,c}(1)$) is present. Conversely, at pH 9, two distinct cathodic peaks are visible at 1 V/s (Figure 2c); in fact, at higher pH, $E_{p,c}(2)$ is even more prominent. Notably, during the first cathodic sweep, $E_{p,c}(2)$ predominates, while in subsequent cathodic sweeps, both $E_{p,c}(1)$ and $E_{p,c}(2)$ are resolved.

This pH and scan rate dependences led us to believe that the iNOS heme axial water ligand is involved in an equilibrium that gives rise to $E_{p,c}(1)$ and $E_{p,c}(2)$ (Scheme 1). At pH 7 and 1 V/s, the first cathodic sweep results in $E_{p,c}(2)$; this correlates with initial reduction of $\text{Fe}^{\text{III}}-\text{OH}_2$. The lability of the $\text{Fe}^{\text{II}}-\text{OH}_2$ bond results in rapid conversion to a five-coordinate heme.^{17,18} Oxidation of Fe^{II} to Fe^{III} , followed by a second cathodic sweep at intermediate scan rates

(1 V/s, Figure 2a), gives rise to two reduction peaks, corresponding to reduction of water-free ($E_{p,c}(1)$) and water-bound ($E_{p,c}(2)$) Fe^{III} hemes.

Scheme 1



Our interpretation is supported by the following observations. First, $E_{p,c}(1)$ and $E_{p,c}(2)$ differ by approximately 133 mV, a value consistent with the potential difference between five- and six-coordinate iNOS hemes.¹² Second, voltammetry at low pH results in a single cathodic peak attributable to $E_{p,c}(1)$. Under acidic conditions, disruption of hydrogen bonding in the heme pocket likely leads to dissociation of the axial water, resulting in a five-coordinate heme. Third, voltammetry at high pH (Figure 2c), where the axial water is partially deprotonated, shows that $E_{p,c}(2)$ is more prominent. Hydroxide will have greater affinity for the cationic metal center, thereby shifting the cathodic peak distribution further to $E_{p,c}(2)$. Fourth, voltammetry in the presence of imidazole yields a single cathodic process, consistent with constant heme ligation as it cycles between oxidation states (Figure S2).¹⁹

The dependence of the cathodic peak distribution on scan rate can be explained in terms of Scheme 1. At high scan rates, after oxidation of Fe^{II} , reduction of water-free Fe^{III} occurs faster than water ligation to Fe^{III} , yielding only $E_{p,c}(1)$. Conversely, scanning slowly allows water enough time to ligate Fe^{III} before reduction, shifting the cathodic peak distribution to $E_{p,c}(2)$. Thus, intermediate scan rates yield both $E_{p,c}(1)$ and $E_{p,c}(2)$.

Using $E_{p,c}(1)$, k° ($\Delta G^\circ = 0$) for ET to the five-coordinate heme evaluated at 16.7 V/s and pH 7 is 370 s^{-1} .²⁰ With this value, we used digital simulation to effectively model the experimental results in Figure 2a (see Supporting Information). From the simulation, we estimated values for k_1 , k_2 , and k_3 in Scheme 1 to be 1, 0.5, and $> 100 \text{ s}^{-1}$, respectively. As expected,^{17,18} water dissociation from Fe^{II} is rapid (k_3), while the kinetics of the Fe^{III} equilibrium with water (k_1 , k_2) appear slow. In the catalytic cycle of iNOS, the first ET event is proposed to be the rate-limiting step for substrate turnover, occurring at 1 s^{-1} .²¹ k° for the five-coordinate heme indicates that well-coupled pathways do exist between the protein surface and the heme; thus, electron tunneling is unlikely to be

rate-limiting in catalysis. Rather, our simulation suggests that water dissociation from Fe^{III} may function as a potential gating mechanism for the catalytic cycle.²²

Proton-coupled ET was observed for nNOS in DDAB films, as demonstrated by the variation of $E_{1/2}$ with pH.⁷ For comparison, we performed voltammetry on iNOS at 200 mV/s and plotted the variation of $E_{1/2}$ with pH for E_1 (Figure 2d).²³ The data reveal no variation of $E_{1/2}$ for $\text{pH} < 5$, implying simple conversion of five-coordinate Fe^{III} to Fe^{II} . At $\text{pH} > 7$, $E_{1/2}$ varies linearly with pH according to -53 mV/pH unit, as is often found for heme proteins.^{24,25} The observation of proton-coupled ET in a pH range where the heme is water-ligated further indicates that the heme axial water ligand gates ET in iNOS.

Acknowledgment. Supported by NSERC (Canada) (A.K.U.), HHMI (A.K.U.), NIH (DK19038; GM070868), the Packard Foundation (M.G.H.), the Ellison Medical Foundation (Senior Scholar Award in Aging to H.B.G.), and the Parsons Foundation (W.B.B.); S. Luzzi (Berkeley) assisted with protein expression and purification.

Supporting Information Available: Details of electrode preparation, protein purification, digital simulations, voltammograms in the presence of dioxygen and imidazole, cytochrome *c* voltammetry, and an iNOS absorption spectrum in a DDAB film. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Culotta, E.; Koshland, D. E. *Science* **1992**, *258*, 1862–1865.
- Alderton, W. K.; Cooper, C. E.; Knowles, R. G. *Biochem. J.* **2001**, *357*, 593–615.
- Aktan, F. *Life Sci.* **2003**, *75*, 639–657.
- Hurshman, A. R.; Krebs, C.; Edmondson, D. E.; Marletta, M. A. *Biochemistry* **2003**, *42*, 13287–13303.
- Rusling, J. F. *Acc. Chem. Res.* **1998**, *31*, 363–369.
- Studies have shown that heme proteins retain their solution spectroscopic properties within surfactant films. See, for example: (a) Ma, H.; Hu, N. *Anal. Lett.* **2001**, *34*, 339–361. (b) Rusling, J. F.; Nassar, A. F. *J. Am. Chem. Soc.* **1993**, *115*, 11891–11897.
- Direct electrochemistry of the neuronal nitric oxide synthase heme domain utilizing the surfactant film methodology has been reported: Bayachou, M.; Boutros, J. A. *J. Am. Chem. Soc.* **2004**, *126*, 12722.
- Immoos, C. E.; Chou, J.; Bayachou, M.; Blair, E.; Greaves, J.; Farmer, P. J. *J. Am. Chem. Soc.* **2004**, *126*, 4934–4942.
- Ma, H.; Hu, N. *Anal. Lett.* **2001**, *34*, 339–361.
- Zhang, Z.; Nassar, A.-E.; Lu, Z.; Schenkman, J. B.; Rusling, J. F. *J. Chem. Soc., Faraday Trans.* **1997**, *93*, 1769–1774.
- Fleming, B. D.; Tian, Y.; Bell, S. G.; Wong, L.; Urlacher, V.; Hill, H. A. O. *Eur. J. Biochem.* **2003**, *270*, 4082–4088.
- Presta, A.; Weber-Main, A. M.; Stankovich, M. T.; Stuehr, D. J. *J. Am. Chem. Soc.* **1998**, *120*, 9460–9465.
- Lvov, Y. M.; Lu, Z.; Schenkman, J. B.; Zu, X.; Rusling, J. F. *J. Am. Chem. Soc.* **1998**, *120*, 4073–4080.
- Tezcan, F. A.; Winkler, J. R.; Gray, H. B. *J. Am. Chem. Soc.* **1998**, *120*, 13383–13388.
- Bard, A. J.; Faulkner, L. R. *Electrochemical Methods*, 2nd ed.; John Wiley & Sons: New York, 2001.
- Lin, R.; Immoos, C. E.; Farmer, P. J. *J. Biol. Inorg. Chem.* **2000**, *5*, 738–747.
- Wilker, J. J.; Dmochowski, I. J.; Dawson, J. H.; Winkler, J. R.; Gray, H. B. *Angew. Chem., Int. Ed.* **1999**, *38*, 90–92.
- Crutchley, R. J.; Ellis, W. R.; Gray, H. B. *J. Am. Chem. Soc.* **1985**, *107*, 5002–5004.
- Dawson, J. H.; Andersson, L. A.; Sono, M. *J. Biol. Chem.* **1983**, *258*, 13637–13645.
- Laviron, E. *J. Electroanal. Chem.* **1979**, *101*, 19–28.
- Presta, A.; Siddhanta, U.; Wu, C.; Sennequier, N.; Huang, L.; Abu-Soud, H. M.; Erzurum, S.; Stuehr, D. J. *Biochemistry* **1998**, *37*, 298–310.
- For a discussion of gated ET in proteins, see: Armstrong, F. A. *J. Chem. Soc., Dalton Trans.* **2002**, 661–671.
- At 200 mV/s, the two cathodic peaks in E_1 collapse into a single peak, yielding a potential representative of the population distribution between $E_{p,c}(1)$ and $E_{p,c}(2)$ and permitting us to estimate a midpoint potential ($E_{1/2}$) for E_1 .
- Munge, B.; Estavillo, C.; Schenkman, J. B.; Rusling, J. F. *ChemBioChem* **2003**, *4*, 82–89.
- Aguey-Zinsou, K.; Bernhardt, P. V.; Voss, J. J. D.; Slessor, K. E. *Chem. Commun.* **2003**, 418–419.

JA0520114