


Engineered Biosynthesis of Aklanonic Acid Analogues

Taek Soon Lee, Chaitan Khosla, and Yi Tang

J. Am. Chem. Soc., **2005**, 127 (35), 12254-12262 • DOI: 10.1021/ja051429z • Publication Date (Web): 10 August 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 7 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Engineered Biosynthesis of Aklanonic Acid Analogues

Taek Soon Lee,[†] Chaitan Khosla,^{*,†,‡,§} and Yi Tang[¶]

Contribution from the Departments of Chemistry, Chemical Engineering, and Biochemistry, Stanford University, Stanford, California 94305, and Department of Chemical Engineering, University of California at Los Angeles, Los Angeles, California 90095

Received March 6, 2005; E-mail: khosla@stanford.edu

Abstract: Aklanonic acid, an anthraquinone natural product, is a common advanced intermediate in the biosynthesis of several antitumor polyketide antibiotics, including doxorubicin and aclacinomycin A. Intensive semisynthetic and biosynthetic efforts have been directed toward developing improved analogues of these clinically important compounds. The primer unit of such polyfunctional aromatic polyketides is an attractive site for introducing novel chemical functionality, and attempts have been made to modify the primer unit by precursor-directed biosynthesis or protein engineering of the polyketide synthase (PKS). We have previously demonstrated the feasibility of engineering bimodular aromatic PKSs capable of synthesizing unnatural hexaketides and octaketides. In this report, we extend this ability by preparing analogues of aklanonic acid, a decaketide, and its methyl ester. For example, by recombining the R1128 initiation module with the dodecaketide-specific pradimicin PKS, the isobutyryl-primed analogue of aklanonic acid (YT296b, **10**) and its methyl ester (YT299b, **12**) were prepared. In contrast, elongation modules from dodecaketide-specific spore pigment PKSs were unable to interact with the R1128 initiation module. Thus, in addition to revealing a practical route to new anthracycline antibiotics, we also observed a fundamental incompatibility between antibiotic and spore pigment biosynthesis in the actinomycetes bacteria.

Introduction

Aklanonic acid, a member of the polyketide family of natural products,^{1,2} is an anthraquinone compound and a biosynthetic precursor of several important antibiotics, including daunorubicin, baumycin A1, nogalamycin, and the anticancer drugs, doxorubicin and aclacinomycin A (Figure 1A and Figure 2). The primary differences between these natural products lie in their substitution patterns, especially their glycosylation patterns. Over the past few decades, there have been intensive attempts to prepare semisynthetic analogues of this family of compounds³ (Figure 1B). For example, anthracyclines containing a disaccharide chain with substitutions at 4'-hydroxy group or a monosaccharide with a methoxymorpholino ring (PNU152243) have shown promising activity.⁴⁻⁶ Similarly, removal of the 4-methoxy group from certain anthracyclines resulted in increased drug activity.^{7,8} The C-9 side chain has also been

modified with or without reduction of the carbonyl group.⁹⁻¹¹ At least one of these analogues (MEN10755) is currently in clinical trials.³

Much has been learned within the past two decades about the biosynthesis of anthracycline natural products through the combined exploitation of chemical, biochemical, and genetic approaches.^{4,12,13} These insights have started to yield practical benefits; for example, a bacterium has been engineered to synthesize epirubicin, a clinically important semisynthetic analogue of doxorubicin with reduced side effects¹⁴ (Figure 1B). Notwithstanding this progress, an even greater diversity of biosynthetic anthracyclines can be realized when the polyketide backbones of these natural products are systematically modified. Here, we describe a new strategy for achieving this goal.

The primer unit in aromatic polyketide biosynthesis is an attractive site for introducing novel chemical functionality into existing natural products.¹⁵ In the absence of a dedicated mechanism for introducing non-acetyl primer units, the minimal PKS, comprising a ketosynthase (KS), a chain length factor (CLF), an acyl carrier protein (ACP), and a malonyl-CoA:ACP

[†] Department of Chemistry, Stanford University.

[‡] Department of Chemical Engineering, Stanford University.

[§] Department of Biochemistry, Stanford University.

[¶] University of California at Los Angeles.

- (1) O'Hagan, D. *The Polyketide Metabolites*; Harwood: Chichester, UK, 1991.
- (2) Hopwood, D. A. *Chem. Rev.* **1997**, *97*, 2465.
- (3) Binaschi, M.; Bigioni, M.; Cipollone, A.; Rossi, C.; Goso, C.; Maggi, C. A.; Capranico, G.; Animati, F. *Curr. Med. Chem.: Anti-Cancer Agents* **2001**, *1*, 113.
- (4) Arcamone, F.; Cassinelli, G. *Curr. Med. Chem.* **1998**, *5*, 391.
- (5) Arcamone, F. M. *Biochimie* **1998**, *80*, 201.
- (6) Grandi, M.; Pezzoni, G.; Ballinari, D.; Capolongo, L.; Suarato, A.; Bargiotti, A.; Faiardi, D.; Spreafico, F. *Cancer Treat. Rev.* **1990**, *17*, 133.
- (7) Capranico, G.; Supino, R.; Binaschi, M.; Capolongo, L.; Grandi, M.; Suarato, A.; Zunino, F. *Mol. Pharmacol.* **1994**, *45*, 908.
- (8) Capranico, G.; Zunino, F.; Kohn, K. W.; Pommier, Y. *Biochemistry* **1990**, *29*, 562.

- (9) Bachur, N. R. *J. Pharmacol. Exp. Ther.* **1971**, *177*, 573.
- (10) Cassinelli, G.; Grein, A.; Masi, P.; Suarato, A.; Bernardi, L.; Arcamone, F.; Di Marco, A.; Casazza, A. M.; Pratesi, G.; Soranzo, C. *J. Antibiot.* **1978**, *31*, 178.
- (11) Acton, E. M.; Tong, G. L.; Mosher, C. W.; Smith, T. H.; Henry, D. W. *J. Med. Chem.* **1979**, *22*, 922.
- (12) Hutchinson, C. R. *Chem. Rev.* **1997**, *97*, 2525.
- (13) Kantola, J.; Kunnari, T.; Hautala, A.; Hakala, J.; Ylihonko, K.; Mantsala, P. *Microbiology* **2000**, *146* (Pt 1), 155.
- (14) Madduri, K.; Kennedy, J.; Rivola, G.; Inveni-Solari, A.; Filippini, S.; Zanuso, G.; Colombo, A. L.; Gewain, K. M.; Occi, J. L.; MacNeil, D. J.; Hutchinson, C. R. *Nat. Biotechnol.* **1998**, *16*, 69.
- (15) Moore, B. S.; Hertweck, C. *Nat. Prod. Rep.* **2002**, *19*, 70.




Figure 1. (A) Examples of natural anthracycline polyketides. (B) Examples of clinically useful semisynthetic analogues of daunorubicin and doxorubicin.




Figure 2. Biosynthesis of daunorubicin, doxorubicin, and aclacinomycin A.

transacylase (MAT), decarboxylates one malonyl equivalent and initiates polyketide assembly with an acetyl group.^{16–18} However, certain natural products, such as doxorubicin, wailupe-mycin, and the R1128, estrogen receptor antagonists, can incorporate various non-acetyl primers, such as propionate, benzoate, and a range of short-chain fatty acids, into their

backbones (Figure 3).^{12,19–21} In some cases, such as the R1128 PKS and the daunorubicin PKS, the synthase includes a dedicated module of enzymes for chain initiation.^{22,23} Recent studies have demonstrated that these initiation modules can collaborate with elongation modules from heterologous octa-

(16) McDaniel, R.; Ebert-Khosla, S.; Hopwood, D. A.; Khosla, C. *Science* **1993**, *262*, 1546.

(17) McDaniel, R.; Ebert-Khosla, S.; Hopwood, D. A.; Khosla, C. *Nature* **1995**, *375*, 549.

(18) Hopwood, D. A.; Sherman, D. H. *Annu. Rev. Genet.* **1990**, *24*, 37.

(19) Hori, Y.; Abe, Y.; Ezaki, M.; Goto, T.; Okuhara, M.; Kohsaka, M. *J. Antibiot.* **1993**, *46*, 1055.

(20) Sitachitta, N.; Gadepalli, M.; Davison, B. S. *Tetrahedron* **1996**, *52*, 8073.

(21) Piel, J.; Hertweck, C.; Shipley, P. R.; Hunt, D. M.; Newman, M. S.; Moore, B. S. *Chem. Biol.* **2000**, *7*, 943.




Figure 3. Non-acetate-primed polyketide biosynthesis. (A) R1128 biosynthesis, (B) doxorubicin biosynthesis, (C) wailupemycin A biosynthesis. Primer units are in bold.

ketide synthases and decaketide synthases to produce non-acetyl-primed hexaketide and octaketide natural products, respectively.²⁴ In each case, chain length of the polyketide backbone is dictated by the elongation module (i.e., the minimal PKS). Since the anthracycline antitumor antibiotics shown in Figure 1A are decaketides, it was therefore hypothesized that primer unit modifications of this class of natural products would require the use of a dodecaketide PKS. Specifically, we targeted the engineering of aklanonic acid and aklanonic acid methyl ester analogues since these compounds are the first common stable intermediates in the biosynthesis of the antitumor compounds shown in Figure 1A.

Several dodecaketide synthases have been reported in the literature. For example, the spore pigment PKSs from various *Streptomyces* species, including *whiE* from *S. coelicolor*,²⁵ *sch* from *S. halstedii*, and *cur* from *S. curaco*, synthesize dodecaketide backbones.^{26,27} The pradimicin PKS from *Actinomadura hibisca* has also been identified as a dodecaketide synthase.²⁸ Although our initial efforts to prepare aklanonic acid analogues attempted to harness the spore pigment PKSs, we observed a fundamental incompatibility between the R1128 initiation module and the spore pigment elongation modules. We therefore

harnessed the pradimicin KS-CLF to successfully produce a range of aklanonic acid analogues in good yield. In addition to providing a route to new antitumor compounds, our results highlight the biochemical logic of PKS specificity during contemporaneous antibiotic and spore pigment synthesis in the life cycle of an actinomycete bacterium.

Experimental Methods

Materials. Silica gel plates Si250F for thin layer chromatography were obtained from J. T. Baker. All other chemicals and solvents were purchased from Sigma, Aldrich, and Fisher Scientific and were of the highest available grade.

Bacterial Strain and General Methods for DNA Manipulation. *S. coelicolor* strain CH999 was used as the host for transformation by shuttle vectors. Protoplast preparation and PEG-assisted transformation were performed as described by Hopwood et al.²⁹ All cloning steps were performed in *Escherichia coli* strain XL-1 Blue (Stratagene). PCR was performed using *PfuTurbo* polymerase (Stratagene). PCR products were first cloned into a pCR-Blunt vector (Invitrogen) followed by DNA sequencing (Stanford PAN Facility). Unmethylated DNA was obtained using the methylase-deficient strain GM2163 (New England Biolabs). T4 DNA ligase (Invitrogen) was used for ligation of restriction fragments.

Construction of Plasmids. The following primers were used to amplify the individual genes: *Sch* KS: 5'-GATTAATTAAGGAG-GAGCCATCATCATGAGCCGCCGGGTCTCGTCACCGG-3', 5'-CGATGCATTCATCTCAGCCTCTCTCCGGTCCGG-3' (PacI/NsiI); *Sch* CLF: 5'-GAATGCATGGAGGAGCCATCATGAGCG-

(22) Meadows, E. S.; Khosla, C. *Biochemistry* **2001**, *40*, 14855.
(23) Bao, W.; Sheldon, P. J.; Wendt-Pienkowski, E.; Hutchinson, C. R. *J. Bacteriol.* **1999**, *181*, 4690.
(24) Tang, Y.; Lee, T. S.; Khosla, C. *PLoS Biol.* **2004**, *2*, E31.
(25) Davis, N. K.; Chater, K. F. *Mol. Microbiol.* **1990**, *4*, 1679.
(26) Blanco, G.; Pereda, A.; Mendez, C.; Salas, J. A. *Gene* **1992**, *112*, 59.
(27) Bergh, S.; Uhlen, M. *Gene* **1992**, *117*, 131.
(28) Dairi, T.; Hamano, Y.; Igarashi, Y.; Furumai, T.; Oki, T. *Biosci. Biotechnol. Biochem.* **1997**, *61*, 1445.

(29) Kieser, T.; Bibb, M. J.; Buttner, M. J.; Charter, K. F.; Hopwood, D. A. *Practical Streptomyces Genetics*; The John Innes Foundation: Norwich, UK, 2000.

Table 1. Plasmid Constructions and Resulting Polyketide Products^a

plasmid	minimal PKS ^b	KR (actIII)	aromatase, cyclase	R1128 LM	polyketide products	ref
pYT227	<i>sch</i> with <i>whiE ACP</i>	—	—	—	TW93c-h	ref 30
pYT226	<i>sch</i> with <i>whiE ACP</i>	+	—	—	TW94c-d	ref 30
pYT232	<i>sch</i> with <i>whiE ACP</i>	—	—	+	TW93c-h	this work
pYT231	<i>sch</i> with <i>whiE ACP</i>	+	—	+	TW94c-d Y231a,b (1, 2)	this work
pYT236	<i>sch</i> with <i>whiE ACP</i>	—	<i>whiE CYC</i>	+	TW95a,b	ref 30, this work
pYT210	<i>sch</i> with <i>zhuN</i>	—	<i>zhuI, zhuJ</i>	+	no polyketide	this work
pYT209	<i>sch</i> with <i>zhuN</i>	+	<i>actVII, actIV</i>	+	No polyketide	this work
pYT243	<i>pms</i> with <i>zhuN</i>	—	—	+	YT243a,b (3, 4) (SEK15-like)	this work
pYT266	<i>pms</i> with <i>zhuN</i>	+	—	+	YT266a,b (5, 6) (RM20b-like)	this work
pYT253	<i>pms</i> with <i>zhuN</i>	+	<i>pms CYC</i>	+	TW95a,b	this work
pYT259	<i>pms</i> with <i>pms ACP</i>	+	<i>pms CYC</i>	+	TW95a,b	this work
pYT292	<i>pms</i> with <i>zhuN</i>	+	<i>grisARO/CYC</i>	+	YT292a,b (7, 8) (SEK43-like)	this work
pYT296	<i>pms</i> with <i>zhuN</i>	+	<i>grisARO/CYC dpsY</i>	+	YT296a,b (9, 10) (aklanonic acid-like)	this work
pYT299 ^c	<i>pms</i> with <i>zhuN</i>	+	<i>grisARO/CYC dpsY</i>	+	YT299a,b (11, 12) (methyl ester)	this work

^a The *Streptomyces coelicolor* strain CH999 is used as the host for polyketide biosynthesis. Each plasmid is derived from pRM5.¹⁶ Each construct contains genes for the minimal PKS, cyclases, and the R1128 initiation module, which consists of *zhuG*, *zhuH*, and *zhuC*. Products are analyzed by LC/MS and NMR.
^b The minimal PKS consists of the indicated KS—CLF, ACP, and the endogenous MAT. ^c pYT299 has methyl transferase (*dnrC*) gene, which is different from pYT296.

CACCCCGGCGAGCCGTCG-3' (NsiI); *whiE ACP*: 5'-CCTCTAGAG-GAGGAGCCCATATGACCGATCAGCAGCTGGAC-3', 5'-GGAC-TAGTCCACTCCCCTCTTGAGGGCGCC-3' (XbaI, NdeI/SpeI); *whiE* cyclase: 5'-GCACTAGTGCCTCAAGACGGGAGTGTGACATG-3' (SpeI); *Pms* KS: 5'-GATTAATTAAGGAGGAGCCATCATGAGC-CGACCCAGGGGGGGCGG-3', 5'-GGATGCATCATGCGGCCCTCCGACGACGG-3' (PacI/NsiI); *Pms* CLF: 5'-GGATGCATGGAG-GAGCCATCATGAGCGTCTCGACGGCGGACGCGCCGG-3', 5'-GGTCTAGAGGTCAGGTGTTCCGGCCGCGCAGCACC-3' (NsiI/XbaI); *Pms* ACP: 5'-GGTCTAGAGGAGGAGCCCATATGGCAA-CCCAGAACGCACCATCG-3', 5'-GGACTAGTCATCGCTCGGC-CACCGCCCGCCG-3' (XbaI, NdeI/SpeI); *Pms* CYC: 5'-GGAC-TAGTGGAGGAGCCATCATGACGCAGCTGGCGCACCGACAGCG-3', 5'-GGTCACCGCGCCCGGGCCCGCCG-3' (SpeI); *gris ARO/CYC*: 5'-GGTITAAACGAGGAGCCAGCATGTCGACGCCCGCCT-GCGCGAGG-3', 5'-GGACTAGTCAGCCCGCCGCGCCGCGGCACG-3' (PmeI/SpeI); *dpsY* CYC: 5'-GGACTAGTGGAGGAGCCAGCAT-GTGCTGTGGTGGCCCGAGAGGAG-3', 5'-CATCCTAGGAATT-AGCTAGCTCATTCGTCGACGAAGCGACCG-3' (SpeI/NheI/AvrII); *dnrG* oxidase: 5'-GATGCTAGCGGAGGAGCCAGCATGCCA-CAGCCAGAACC GAACGACG-3', 5'-CATCCTAGGAATAT-CACGTCGTCGCCCTCCGGCCCG-3' (NheI/AvrII); *dnrC* methyl esterase: 5'-GGCCTAGGAGGAGCCAGCATGCAGGAT-TCGTCTACAAAGAG-3', 5'-GGTCAGTGGGCGACCCGGGCG-GTGACG-3' (AvrII); *dnrD* cyclase: 5'-GGCCTAGGAGGAGCCAG-CATGAGCACGCAGATCGATCTGGTCCG-3', 5'-GGACTAGT-CAGGGGCGCCAGCCCTCGGGGGTGGG-3' (AvrII/SpeI). The introduced restriction sites are shown in italics and indicated in parentheses. Genes encoding *sch* KS and CLF were amplified from pIJ4301,³⁰ and genes encoding *whiE* ACP and cyclase were amplified from pIJ4296.³⁰ Genes encoding *pms* KS, CLF, ACP, and CYC were amplified from the genomic DNA of *Actinomadura hibisca* P157-2.²⁸ Genes encoding *gris ARO/CYC* were amplified from pIJ5216.³¹

Genes encoding *dpsY* CYC, *dnrG* oxidase, and *dnrC* methyl esterase were amplified from the genomic DNA of *S. peuceitius*.^{32,33} Different combinations of cassettes, as shown in Table 1, were introduced into either pYT127 (KR—R1128 initiation module) or pYT128 (R1128 initiation module without KR) to yield the corresponding constructs.²⁴ The physical maps of plasmids pYT266, pYT292, pYT296, and pYT299 are described in Figure 1 of the Supporting Information.

- (30) Yu, T. W.; Shen, Y.; McDaniel, R.; Floss, H. G.; Khosla, C.; Hopwood, D. A.; Moore, B. S. *J. Am. Chem. Soc.* **1998**, *120*, 7749.
(31) Yu, T. W.; Bibb, M. J.; Revill, W. P.; Hopwood, D. A. *J. Bacteriol.* **1994**, *176*, 2627.
(32) Lomovskaya, N.; Doi-Katayama, Y.; Filippini, S.; Nastro, C.; Fonstein, L.; Gallo, M.; Colombo, A. L.; Hutchinson, C. R. *J. Bacteriol.* **1998**, *180*, 2379.
(33) Madduri, K.; Hutchinson, C. R. *J. Bacteriol.* **1995**, *177*, 3879.

Culture Conditions, Extractions, and Small-Scale Analysis.

Strains were grown on R5 media plates²⁹ containing 50 mg/L thioestrepton at 30 °C for 7–10 days. For LC/MS and analytical HPLC analysis, a well-pigmented plate was chopped into fine pieces and extracted with 30 mL of ethyl acetate containing 1% acetic acid. The solvent was removed in vacuo, and the residue was dissolved in 1 mL of methanol. The polyketide products were separated by analytical reverse-phase HPLC and detected at 280 and 410 nm using the Alltech Econosphere C18 column (250 mm × 4.6 mm); linear gradient: 10% acetonitrile (MeCN) in water (0.1% TFA) to 60% MeCN in water (0.1% TFA) over 30 min with a flow rate of 1 mL/min. HPLC retention times were as follows: **1**: 25.3 min; **2**: 21.8 min; **3**: 24.1 min; **4**: 25.6 min; **5**: 30.5 min; **6**: 32.6 min; **7**: 24.6 min; **8**: 27.7 min; **9**: 38.2 min; **10**: 38.7 min; **11**: 40.5 min; **12**: 40.8 min; linear gradient: 40% MeCN in water (0.1% TFA) to 80% MeCN in water (0.1% TFA) over 30 min with the flow rate of 1 mL/min; **9**: 22.3 min; **10**: 25.3 min; **11**: 33.8 min; **12**: 35.1 min (data not shown). LC/MS was performed at the Vincent Coates Foundation Mass Spectrometry Laboratory at Stanford University using a ThermoFinnigan quadrupole ion trap LC/MS system either by electrospray ionization or by atmospheric pressure chemical ionization (both positive and negative ionization).

Large-Scale Production and Isolation. A sufficient number of R5 plates (40–120 plates depending on the yield of the product) streaked with the desired CH999 strains were incubated at 30 °C for 7–10 days. The plates were chopped into fine pieces and extracted with 1–3 L of ethyl acetate containing 1% acetic acid according to the number of plates. The extract was dried over sodium sulfate, and the solvent was removed in vacuo. The residue was purified by short path silica gel column chromatography (eluted with 50% ethyl acetate in hexane with 1% acetic acid), and the product-containing fractions were collected for preparative reverse-phase HPLC. The solvent was removed in vacuo, and the residue was dissolved in 5 mL of methanol and filtered for injection into the preparative reverse-phase HPLC column (250 mm × 22.5 mm, C18 column; Alltech Econosil). A 10–60% MeCN in water (0.1% TFA) gradient was used over 50 min with a flow rate of 3 mL/min. The solvent was removed in vacuo from the collected fractions containing the expected biosynthetic product, and the residue was dissolved in 1 mL of acetone to be loaded onto a preparative TLC plate. Preparative TLCs were developed in various compositions of ethyl acetate in hexane with or without 1% acetic acid, and the desired bands were excised from the TLC plate and washed with ethyl acetate for the recovery of the compound.

Spectroscopic Analysis. The ¹H and ¹³C NMR spectra were obtained on Varian Inova-500 or Mercury-400 spectrometers. The ¹H NMR spectra were referenced at 0 ppm with internal tetramethylsilane in CDCl₃ and at 3.31 ppm in methanol-*d*₄. The ¹³C NMR spectra were




Figure 4. Biosynthesis by spore pigment producing dodecaketide synthases.

referenced at 77.1 ppm in CDCl_3 and at 49.1 ppm in methanol- d_4 . Multiple-bond ^1H - ^{13}C connectivity was determined by proton-detected HMBC on Varian Inova-600 NMR instruments. The ^1H and ^{13}C NMR assignments are summarized in Tables 2–4. Long-range ^{13}C - ^1H HMBC correlation data for the compounds. **1:** H4 (C2, C3, C5, C6), H8 (C6, C7, C9, C10, C12), H10 (C8, C9, C11), H14 (C6, C12, C13, C16), H20 (C18, C22), H22 (C18, C20, C24), H24 (C18, C22, C23). **2:** H4 (C1, C2, C3, C6), H8 (C6, C10, C12), H9 (C7, C11), H10 (C11, C12), H14 (C6, C12, C13, C16), H20 (C18, C22), H22 (C18, C20, C24), H24 (C18, C22, C23). **10:** H2 (C1, C3, C4, C16), H4 (C2, C5, C6, C14), H8 (C6, C10), H9 (C7, C11), H10 (C8, C12), H20 (C18, C19, C21, C22, C23), H21 (C19, C20, C22, C23), H22 (C20, C21, C23), H23 (C21, C22, C23). **12:** H2 (C1, C3, C4, C16), H4 (C2, C5, C6, C14), H8 (C6, C10), H9 (C7, C11), H10 (C8, C12), H20 (C17, C18, C19, C22), H21 (C19, C20, C22, C23), H22 (C20, C21, C23), H23 (C21, C22, C23), H25 (C1). LR-ESIMS and HR-ESIMS were collected under positive or negative ionization mode as follows: (+)-ESI mass spectra are obtained as $[\text{M} + \text{H}]^+$ or $[\text{M} + \text{Na}]^+$, and all (–)-ESI mass spectra are obtained as $[\text{M} - \text{H}]^-$. LR-ESIMS m/z : (**1**) 451.2 ($[\text{M} - \text{H}]^-$, $\text{C}_{24}\text{H}_{19}\text{O}_9$); (**2**) 433.0 ($[\text{M} - \text{H}]^-$, $\text{C}_{24}\text{H}_{17}\text{O}_8$); (**3**) 426.9 ($[\text{M} + \text{H}]^+$, $\text{C}_{23}\text{H}_{23}\text{O}_8$); (**4**) 440.9 ($[\text{M} + \text{H}]^+$, $\text{C}_{24}\text{H}_{25}\text{O}_8$). HR-ESIMS m/z : (**5**) 427.1389 (calcd for $\text{C}_{23}\text{H}_{23}\text{O}_8$, 427.1393); (**6**) 441.1549 (calcd for $\text{C}_{24}\text{H}_{25}\text{O}_8$, 441.1549); (**7**) 409.1293 (calcd for $\text{C}_{23}\text{H}_{21}\text{O}_7$, 409.1287); (**8**) 423.1442 (calcd for $\text{C}_{24}\text{H}_{23}\text{O}_7$, 423.1444); (**9**) 419.1125 (calcd for $\text{C}_{22}\text{H}_{20}\text{O}_7\text{Na}$, 419.1107); (**10**) 461.1225 (calcd for $\text{C}_{24}\text{H}_{22}\text{O}_8\text{Na}$, 461.1212); (**11**) 433.1255 (calcd for $\text{C}_{23}\text{H}_{22}\text{O}_7\text{Na}$, 433.1263); (**12**) 475.1364 (calcd for $\text{C}_{25}\text{H}_{24}\text{O}_8\text{Na}$, 475.1369). HR-ESIMS was performed at the Vincent Coates Foundation Mass Spectrometry Laboratory at Stanford University using a Micromass QToF API US (Manchester, United Kingdom) quadrupole-time-of-flight mass spectrometer, which was operated in MS-only mode with an internal standard (polyalanine, Sigma P9003).

Results

The plasmids constructed for this study are shown in Table 1. The plasmids were derived from the *E. coli/S. coelicolor*

shuttle vector pRM5¹⁶ (Figure 1 in Supporting Information). Each of the plasmids was transformed into the engineered *S. coelicolor* host CH999,¹⁶ and the production of polyketides was initially assayed by using LC/MS.

Evaluation of KS–CLF Enzymes from Spore Pigment Biosynthetic Pathways as Catalysts for Novel Decaketide Biosynthesis. For the biosynthesis of new aklanonic acid derivatives, we first examined the ability of dodecaketide-specific Sch KS–CLF from the spore pigment biosynthetic pathway of *S. halstedii*²⁶ to productively interact with the R1128 initiation PKS module. The biosynthetic products of both the Sch minimal PKS alone and the Sch minimal PKS with *acIII* KR have been characterized³⁰ (Figure 4, and Figure 2 in Supporting Information). When the Sch KS–CLF was recombined with the WhiE ACP (pYT227), we were able to observe the unreduced dodecaketides (TW93c-d). Similarly, addition of KR to pYT227 (resulting in pYT226) gave rise to the reduced dodecaketides TW94c-d. The genes encoding the R1128 initiation module were therefore added to pYT227 and pYT226 to yield pYT232 and pYT231, respectively. Constructs pYT232 and pYT231 were introduced into *S. coelicolor* CH999 via transformation; however, LC/MS analysis of their biosynthetic products did not reveal the presence of isobutyryl or propionyl-primed polyketides, as observed earlier for minimal PKSs with shorter chain length specificity.²⁴ Instead, the transformants only produced acetyl-primed products, such as TW93c-d (CH999/pYT232) and TW94c-d (CH999/pYT231). Two new acetyl-primer-derived biosynthetic products were isolated from CH999/pYT231, with masses of 452 and 434. The yields of these two compounds were comparable to that of TW94c-d. The two compounds were characterized using ^1H and ^{13}C NMR (Table 2) along with COSY and HMBC analysis. We anticipated that the compound with a mass of 434 (YT231b, **2**) may be a

Table 2. ^1H and ^{13}C NMR Assignments for YT231a (**1**) and YT231b (**2**)^a

No. ^b	YT231a		YT231b	
	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))
1	164.3		162.6	
2	90	5.44 (d, 1H, 3)	96	5.46 (d, 1H, 2)
3	172.8		173.5	
4	107.5	6.11 (s, 1H)	108.0	6.09 (d, 1H, 2)
5	160.5		157.3	
6	125.8		122.8	
7	141.1		136.1	
8	37	3.09 (dd, 1H, 4, 17), 2.86 (dd, 1H, 6, 17)	117.4	7.05 (dd, 1H, 1, 8)
9	66.6	4.32 (m, 1H)	129.2	7.28 (t, 1H, 8)
10	30.8	2.97 (dd, 1H, 4, 17), 2.75 (dd, 1H, 6, 17)	109.9	6.75 (dd, 1H, 1, 8)
11	203.9		155.8	
12	116.7		114.9	
13	161.9		159.9	
14	119.2	6.80 (s, 1H)	111.9	6.71 (s, 1H)
15	143.1		141.7	
16	47.3	4.25 (d, 2H, 19)	49.9	4.30 (s, 2H)
17	204.9		205.4	
18	120.0		120.6	
19	161.2		161.5	
20	101.4	6.14 (s, 1H)	101.3	6.13 (d, 1H, 2)
21	168.1		168.9	
22	111.1	6.14 (s, 1H)	110.8	6.15 (d, 1H, 2)
23	145.8		140.5	
24	21.4	2.16 (s, 3H)	21.0	2.08 (s, 3H)

^a Spectra were obtained at 500 MHz for proton and 125 MHz for carbon and were recorded in CD_3OD . ^b Carbons are labeled as shown in Figure 4.

dehydration product of the compound with a mass of 452 (YT231a, **1**); this was confirmed by ^1H NMR spectroscopic analysis, which shows that the two pairs of *dd* peaks of H-8 and H-10 in the latter compound were replaced by aromatic proton peaks in the former. The two compounds were eventually characterized as having distinct cyclization patterns compared to TW94d (Figure 4 and Table 2).

We further examined the biosynthetic capacity of Sch KS–CLF by addition of the WhiE cyclase to pYT232 (resulting in pYT236). As we expected from the previous study by Moore et al.,³⁰ this construct resulted in abundant production of TW95a and TW95b (the isolated yields were 50 and 8 mg/L, respectively), but it still did not produce any non-acetyl-primed polyketide.

To attempt improving crosstalk between the R1128 initiation module and the Sch KS–CLF, we replaced the WhiE ACP with ZhuN, the ACP from the elongation module of the R1128 PKS.²⁴ This modified bimodular PKS was constructed in the absence (pYT210) and presence (pYT209) of the ketoreductase gene. However, neither construct yielded any polyketide, not even the dodecaketides observed with CH999/pYT231 and CH999/pYT232. Thus, there appears to be a fundamental incompatibility between initiation PKS modules from antibiotic pathways and elongation PKS modules from spore pigment pathways.

Biosynthesis of Non-Acetyl-Primed Decaketide Backbones Using the Pradimicin KS–CLF. To find a dodecaketide synthase that could communicate successfully with the R1128 initiation module, we selected the antifungal pradimicin polyketide synthase (*pms*) as a possible partner for the R1128 initiation module. The *pms* KS and CLF genes were amplified from genomic DNA of *Actinomadura hibisca* P157-2 by PCR.²⁸ We replaced *tcm* KS and *tcm* CLF cassettes in pYT128 with the 1.3 kb *PacI*–*NsiI* cassette of *pms* KS and 1.3 kb *NsiI*–*XbaI* cassette of *pms* CLF to yield pYT243 (Table 1). This construct


was introduced into *S. coelicolor* CH999 by PEG-mediated transformation. LC/MS analysis of the crude extract from the resulting transformant showed two peaks with strong absorption at 280 nm and masses of 426 and 440, corresponding to the expected masses of SEK15 analogues primed by propionyl and isobutyryl groups, respectively (Scheme 1). The more abundant isobutyryl-primed compound was purified by preparative HPLC and TLC. ^1H NMR analysis of this compound showed similar peak assignments for most backbone protons to SEK15³⁴ (Table 3). This compound, however, had three more sets of peaks in the aliphatic proton region than SEK15 had, and the most characteristic peak was a doublet with six protons at $\delta = 0.8$ ppm, which corresponds to two terminal methyl groups in the isobutyryl-primed compound. Further spectroscopic analysis with ^{13}C NMR and COSY confirmed the structure of this compound as the expected SEK15 analogue (YT243b, **4**) along with the comparison with spectroscopic data for SEK15.³⁴

To produce decaketide analogues reduced at C-9, we inserted the *act KR* gene into pYT243, yielding pYT266 (Table 1). The expected biosynthetic products were the propionyl- and isobutyryl-primed analogues of RM20b/c³⁵ (Scheme 1). Indeed, LC/MS analysis revealed two peaks with the expected masses of propionyl- (YT266a, **5**) and isobutyryl (YT266b, **6**)-primed RM20b ($m/z = 427$ and 441 by (–)-ESIMS, respectively), both of which were confirmed by HR-ESIMS. The above results demonstrated that the compatible interactions between the R1128 initiation module and the elongation module from the *pms* PKS can indeed produce non-acetyl-primed decaketides.

In a further step toward engineered biosynthesis of aklanonic acid analogues, we inserted the gene encoding the pradimicin cyclase into plasmids carrying the R1128 initiation module, the

(34) Fu, H.; Ebert-Khosla, S.; Hopwood, D.; Khosla, C. *J. Am. Chem. Soc.* **1994**, *116*, 4166.

(35) Fu, H.; McDaniel, R.; Hopwood, D. A.; Khosla, C. *Biochemistry* **1994**, *33*, 9321.

Scheme 1. Biosynthesis of Non-Acetate-Primed Decaketides by the Bimodular Dodecaketide Polyketide Synthases**Table 3.** ^1H and ^{13}C NMR Assignments for Y1243b (**4**) and Y1292b (**8**)^a

No. ^b	Y1243b		Y1292b	
	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))
1	164.0		166.8	
2	86.6	4.90 (s, 1H)	88.0	4.96 (s, 1H)
3	172.8		172.3	
4	101.8	5.49 (s, 1H)	101.8	5.51 (s, 1H)
5	163.1		162.3	
6	39.0	3.53 (s, 2H)	36.6	3.53 (s, 2H)
7	135.4		133.6	
8	109.1	6.27 (s, 1H)	130.7	6.76 (d, 1H, 8)
9	159.8		120.8	7.18 (t, 1H, 8)
10	101.6	6.16 (s, 1H)	114.7	6.72 (d, 1H, 8)
11	157.4		154.5	
12	121.1		130.0	
13	200.1		199.4	
14	117.2		116.8	
15	163.6		163.0	
16	100.6	6.16 (s, 1H)	100.4	6.09 (s, 1H)
17	162.2		162.7	
18	111.0	6.30 (s, 1H)	109.3	6.11 (s, 1H)
19	141.9		147.4	
20	36.8	2.23 (t, 2H, 8)	32.3	2.27 (t, 2H, 8)
21	27.7	1.37–1.32 (m, 2H)	27.8	1.06–1.01 (m, 2H)
22	31.0	1.37–1.32 (m, 1H)	39.4	1.28–1.24 (m, 1H)
23	22.2	0.78 (d, 6H, 6.5)	22.2	0.70 (d, 6H, 6.5)

^a Spectra were obtained at 500 MHz for proton and 125 MHz for carbon and were recorded in CD_3OD . ^b Carbons are labeled as shown in Scheme 1.

act KR, *pms KS-CLF*, and either *zhuN* or *pms ACP*, yielding plasmids pYT253 and pYT259, respectively. However, neither construct produced the expected C-7/C-12-cyclized polyketide. Instead, these constructs produced only the acetyl-primed compounds, TW95a and TW95b, as evidenced by LC/MS

analysis (data not shown). This result suggests that Pms CYC is a C-9 unreduced polyketide-specific cyclase like ZhuI and ZhuJ in R1128 PKS²⁴ and the cyclases in WhiE PKS.³⁰ We could not observe Y1266 compounds peaks in LC/MC analysis, and this suggests that Pms CYC works faster than act KR. We therefore replaced the *pms* CYC with another bifunctional aromatase/cyclase, the *gris* ARO/CYC from the griseusin gene cluster, which has been previously used for synthesizing SEK43.³⁶ The resulting construct (pYT292) was transformed into *S. coelicolor* CH999. The recombinant strain appeared to produce both the propionyl- (Y1292a, **7**) and the isobutyryl (Y1292b, **8**)-primed analogues of SEK43, as confirmed by LC/MS ($m/z = 411$ and 425 , respectively, by (+)-ESIMS; 409 and 423 , respectively, by (–)-ESIMS) with a combined yield over 30 mg/L (Figure 4). The compound with $m/z = 425$ from LC/MS, which was more abundant, was purified by successive purification steps and identified as the expected isobutyryl-primed SEK43 (Y1292b, **8**) by ^1H and ^{13}C NMR (Table 3), COSY, and comparison with SEK43.¹⁷ These results provide a basis for attempting the engineered biosynthesis of new aklanonic acid analogues.

Biosynthesis of Aklanonic Acid Analogues. For the synthesis of aklanonic acid analogues with non-acetyl primer units, the monocyclic polyketide precursors of Y1292a and Y1292b must be cyclized in a controlled fashion into a tricyclic anthraquinone. For this, we selected the *dpsY* cyclase from the daunorubicin biosynthetic pathway.^{12,37} The 820 bp *dpsY* gene was amplified from genomic DNA of *S. peuceetius* by PCR and inserted as an *SpeI*-*AvrII* cassette into pYT292 to yield the

(36) Zawada, R. J.; Khosla, C. *J. Biol. Chem.* **1997**, *272*, 16184.

(37) Hautala, A.; Torkkell, S.; Raty, K.; Kunnari, T.; Kantola, J.; Mantsala, P.; Hakala, J.; Ylihonko, K. *J. Antibiot.* **2003**, *56*, 143.

Table 4. ^1H and ^{13}C NMR Assignments for YT296a,b (**9**, **10**), YT299a,b (**11**, **12**)^a

No. ^b	YT296a		YT296b		YT299a		YT299b	
	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))	^{13}C δ (ppm)	^1H δ (ppm) (m, area, J_{HH} (Hz))
1	172.4		172.6		170.3		170.3	
2	39.4	3.80 (s, 2H)	40.7	3.86 (s, 2H)	38.9	3.791 (s, 2H)	39.5	3.86 (s, 2H)
3	141.1		141.3		141.5		142.7	
4	122.5	7.78 (s, 1H)	122.3	7.85 (s, 1H)	122.6	7.78 (s, 1H)	122.4	7.77 (s, 1H)
5	133.9		134.2		133.5		133.7	
6	181.0		179.6		192.8		181.2	
7	133.3		132.8		133.4		132.5	
8	125.2	7.86 (dd, 1H, 8, 1.2)	120.6	7.86 (dd, 1H, 8, 1.2)	120.4	7.86 (dd, 1H, 8, 1.2)	120.4	7.86 (dd, 1H, 8, 1.2)
9	137.8	7.73 (t, 1H, 8)	137.9	7.73 (t, 1H, 8)	137.7	7.73 (t, 1H, 8)	137.7	7.72 (t, 1H, 8)
10	136.0	7.35 (dd, 1H, 8, 1.2)	125.2	7.34 (dd, 1H, 8, 1.2)	125.1	7.35 (dd, 1H, 8, 1.2)	125.1	7.33 (dd, 1H, 8, 1.2)
11	160.3		160.6		160.0		160.3	
12	115.5		115.6		115.3		115.8	
13	192.8		193.6		181.2		192.7	
14	115.8		115.8		115.8		115.8	
15	162.8		162.8		162.8		162.8	
16	130.4		133.0		136.6		132.5	
17	206.7		184.4		205.8		184.2	
18	42.4	3.04 (t, 2H, 7)	103.4	6.06 (s, 1H)	42.3	3.04 (t, 2H, 7)	103.2	5.92 (s, 1H)
19	32.5	1.65–1.60 (m, 2H)	196.0		32.3	1.67–1.58 (m, 2H)	195.3	
20	27.8	1.78–1.70 (m, 1H)	37.4	2.44 (t, 2H, 8)	27.8	1.78–1.69 (m, 1H)	36.6	2.41 (t, 2H, 8)
21	22.5	0.94 (d, 6H, 6)	34.8	1.60–1.56 (m, 2H)	22.5	0.94 (d, 6H, 6)	33.6	1.59–1.53 (m, 2H)
22			28.1	1.70–1.62 (m, 1H)	52.5	3.714 (s, 3H)	28.1	1.66–1.61 (m, 1H)
23			22.4	0.94 (d, 6H, 6)			22.8	0.93 (d, 6H, 6.5)
25							52.5	3.71 (s, 3H)

^a Spectra were obtained at 500 MHz for proton and 125 MHz for carbon and were recorded in CDCl_3 . ^b Carbons are labeled as shown in Scheme 1.

construct pYT296. On the basis of earlier precedents, we anticipated that an endogenous oxygenase in *S. coelicolor* CH999 would be able to oxidize the tricyclic core into anthracyclic aklanonic acid analogues.^{17,24} The HPLC trace of the crude extract of CH999/pYT296 showed several characteristic anthraquinone-type peaks with absorption at 280 and 410 nm. LC/MS analysis demonstrated that the two most abundant anthraquinone-type peaks have masses of 396 and 438, corresponding to the expected isobutyryl-primed aklanonic acid analogue and perhaps a related compound that lacks one acetyl elongation unit (Figure 3 in Supporting Information). Production of these metabolites was scaled up. Preparative TLC yielded three major yellow products when developed with 25% ethyl acetate in hexane plus 1% acetic acid. ^1H NMR analysis of one of the products revealed peaks to the aromatic hydrogen atoms and the C-2 benzylic hydrogen of aklanonic acid, but this compound lacked the enolizable methylene in the aklanonic acid side chain ($\delta = 6.0$ ppm) (Table 4). MS analysis of this compound was also consistent with its identity as an isobutyryl-primed nonaketide. Further spectroscopic analysis of the compound via ^{13}C NMR, COSY, and HMBC confirmed its structure, as shown in Scheme 1 (YT296a, **9**, 2 mg/L). The ^1H and ^{13}C NMR spectra of the second compound suggested it was more closely related to aklanonic acid,³⁸ with additional peaks corresponding to the alkyl side chain, including a doublet corresponding to terminal methyl groups, analogous to YT243b (**4**) and YT292b (**8**). COSY and HMBC analyses confirmed the incorporation of isobutyryl primer into this polyketide backbone, and the compound was identified as the isobutyryl-primed deca-ketide (YT296b, **10**, 3 mg/L), the expected analogue of aklanonic acid (Scheme 1). These findings demonstrate the feasibility of engineered biosynthesis of aklanonic acid ana-

logues by harnessing the initiation module from the R1128 PKS, the elongation module from the pradimicin PKS, the *act* ketoreductase, the bifunctional *gris* aromatase/cyclase, and the DpsY cyclase from the daunorubicin biosynthetic pathway.

Biosynthesis of Aklanonic Acid Methyl Ester Analogues.

In daunorubicin biosynthesis (and presumably in the biosynthesis of other anthracyclines, as well), the carboxylic acid of aklanonic acid is converted into a methyl ester, presumably by the DnrC protein.³³ Therefore, we cloned the 850 bp *dnrC* gene from genomic DNA of *S. peuceitius* by PCR and inserted the corresponding SpeI-AvrII cassette into pYT296, yielding pYT299. The recombinant *S. coelicolor* strain CH999/pYT299 was found to produce anthraquinone-like compounds, with expectedly longer retention times via reverse-phase HPLC. LC/MS analysis of crude extracts from this culture identified the two most abundant anthraquinone-type products as compounds with masses of 410 and 452. Corresponding $[\text{M} - \text{OMe}]^+$ peaks (379 and 421, respectively) were also observed, suggesting that these compounds were the isobutyryl-primed nonaketide and deca-ketide anthraquinones, respectively. Large-scale production led to their isolation and identification as methyl esters of YT296a (YT299a, **11**, 2 mg/L) and YT296b (YT299b, **12**, 1 mg/L) by spectroscopic comparison with authentic aklanonic acid methyl ester.³⁸

Discussion

Polyketides are a pharmacologically important class of natural products, having yielded many clinically useful drugs. A number of improved analogues of therapeutically promising polyketides have been prepared by synthetic and, more recently, biosynthetic means. In particular, engineering biosynthetic pathways of

(38) Eckardt, K.; Tresselt, D.; Schumann, G.; Ihn, W.; Wagner, C. *J. Antibiot.* **1985**, *38*, 1034.

polyketide antibiotics to incorporate novel primer units is an effective way to generate analogues of existing natural products.^{15,39,24}

In recent reports, we and others have shown that bacterial aromatic PKSs can be engineered to synthesize rationally modified polycyclic aromatic polyketides.^{24,40–42} Specifically, we have harnessed bimodular PKSs to produce engineered hexaketides and octaketides.²⁴ Since some of the most pharmacologically interesting polyketides, including important anticancer drugs, such as doxorubicin and aclacinomycin, are decaketides, we wished to extend our methodology for primer unit modification to decaketide natural products. In doing so, however, our attempt to harness the dodecaketide PKS from spore pigment pathways failed. Analysis of this failure has highlighted the possible existence of a fundamental incompatibility between initiation modules from antibiotic pathways and elongation modules from spore pigment pathways. Fortunately, the dodecaketide synthase from the pradimicin pathway was able to productively interact with the R1128 initiation module to produce non-acetyl-primed decaketides, including novel aklanonic acid analogues. Aklanonic acid is a key biosynthetic intermediate in the formation of several pharmacologically intriguing anthracyclines. The 4-methylpentanoyl-primed aklanonic acid analogue (YT296b, **10**) and its methyl ester (YT299b, **12**) were biosynthesized by an engineered bimodular aromatic polyketide synthase, together with a nonaketide product (YT296a, **9**) and its methyl ester (YT299a, **11**).

As has been recently demonstrated, the biosynthetic potential of this new source of anthracycline antibiotics can be considerably expanded by feeding natural and unnatural amino acids to the producer organisms.⁴³ For example, we recently produced the 2-cyclopropylacetyl-primed aromatic octaketide, 3,8-dihydroxy-1-(3-cyclopropylpropyl)anthraquinone-2-carboxylic acid, from a culture of CH999/pYT127²⁴ by feeding 1 g/L of cyclopropyl alanine to this organism. The amino acid is converted into 2-cyclopropylacetyl-CoA through amino acid

catabolism, which in turn is incorporated into the polyketide backbone.⁴⁴ The cyclopropyl functionality is useful for a variety of synthetic modifications, highlighting the potential for chemobiosynthetic expansion of the aglycone library.⁴⁵

Further biosynthetic elaboration of these new aklanonic acid analogues can be achieved by reconstituting in *S. coelicolor* CH999 a variety of downstream enzymes, such as the fourth ring cyclase (dnrD), the ketoreductase (dnrE), and the glycosyltransferases from the daunorubicin pathway (Figure 2).^{33,46,47} Analogous glycosyltransferases have been identified in the daunorubicin⁴⁸ and aclacinomycin A⁴⁹ biosynthetic gene clusters. Although the tolerance of these tailoring enzymes for our new aklanonic acid analogues remains to be established, recent studies suggest that the L-2-deoxyfucoyltransferase (AknK) from the aclacinomycin A pathway has broad substrate specificity.⁴⁹

In conclusion, engineering of bimodular aromatic polyketide synthases has opened the door to qualitatively novel anthracycline antibiotics through modifications at a position not readily accessible by synthetic methods alone.

Acknowledgment. We thank Dr. Allis Chien at the Stanford Mass Spectrometry Facility and Dr. Stephen Lynch at the Stanford Chemistry Department NMR Facility for providing help with the mass spectrometric and the NMR spectroscopic analysis of the compounds. This work was supported by a grant from the National Institutes of Health (NIH) (CA 77248 to C.K.).

Supporting Information Available: Physical maps of plasmids pYT266, pYT292, pYT296, and pYT299. HPLC traces for CH999/pYT227, CH999/pYT226, CH999/pYT236, CH999/pYT231, CH999/pYT243, CH999/pYT266, CH999/pYT292, CH999/pYT296, CH999/pYT299, with peak assignment by LC/MS. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA051429Z

- (39) Denoya, C. D.; Fedechko, R. W.; Hafner, E. W.; McArthur, H. A.; Morgenstern, M. R.; Skinner, D. D.; Stutzman-Engwall, K.; Wax, R. G.; Wernau, W. C. *J. Bacteriol.* **1995**, *177*, 3504.
(40) Hutchinson, C. R.; Colombo, A. L. *J. Ind. Microbiol. Biotechnol.* **1999**, *23*, 647.
(41) Metsa-Ketela, M.; Palmu, K.; Kunnari, T.; Ylihanko, K.; Mantsala, P. *Antimicrob. Agents Chemother.* **2003**, *47*, 1291.
(42) Shen, Y.; Yoon, P.; Yu, T. W.; Floss, H. G.; Hopwood, D.; Moore, B. S. *Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96*, 3622.
(43) Tang, Y.; Lee, T. S.; Lee, H. Y.; Khosla, C. *Tetrahedron* **2004**, *60*, 7659.

- (44) The manuscript is prepared.
(45) Wessjohann, L. A.; Brandt, W. *Chem. Rev.* **2003**, *103*, 1625.
(46) Dickens, M. L.; Ye, J.; Strohl, W. R. *J. Bacteriol.* **1995**, *177*, 536.
(47) Kendrew, S. G.; Katayama, K.; Deutsch, E.; Madduri, K.; Hutchinson, C. R. *Biochemistry* **1999**, *38*, 4794.
(48) Otten, S. L.; Liu, X.; Ferguson, J.; Hutchinson, C. R. *J. Bacteriol.* **1995**, *177*, 6688.
(49) Lu, W.; Leimkuhler, C.; Oberthur, M.; Kahne, D.; Walsh, C. T. *Biochemistry* **2004**, *43*, 4548.