


## Metathesis and Metallotropy: A Versatile Combination for the Synthesis of Oligoenynes

Mansuk Kim, and Daesung Lee

*J. Am. Chem. Soc.*, **2005**, 127 (51), 18024-18025 • DOI: 10.1021/ja057153c • Publication Date (Web): 06 December 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009



### More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 8 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)


## Metathesis and Metallotropy: A Versatile Combination for the Synthesis of Oligoenynes

Mansuk Kim and Daesung Lee\*

Department of Chemistry, University of Wisconsin, Madison, Wisconsin 53706

Received October 20, 2005; E-mail: dlee@chem.wisc.edu


The bond reorganization processes, defined as metallotropic shift, of various alkynyl carbene complexes with Rh,<sup>1</sup> Mn,<sup>2</sup> Re,<sup>3</sup> Cr,<sup>4</sup> Mo,<sup>4</sup> and W<sup>4</sup> metals have been reported. The rearrangement involving Rh, Cr, Mo, and W is a [1,3]-shift (eq 1), while that with Mn and Re is formally defined as a [1,1.5]-shift (eq 2). However, the metallotropic shift of ruthenium alkynyl carbene complexes has not been observed until recently.<sup>5</sup> Because the metallotropic [1,3]-shift of a transient ruthenium carbene complex was involved in the enyne ring-closing metathesis (RCM) of diyne-containing substrates (eq 3),<sup>6–8</sup> we surmised that these two bond



reorganization processes may share a close mechanistic relationship. Also, due to their tandem bond-forming and -breaking nature, both processes can be incorporated into a synthetic design in multiple combinations such that an efficient access to remarkably complex networks can arise. The prerequisite for this supposition is that the equilibrium composition or the reactivity of the equilibrating alkynyl ruthenium carbene species in eq 1 or 3 can be controlled to allow only one of the regioisomeric carbene complexes to react selectively. In this vein, the controlled [1,3]-shift behavior of an alkylidene derived from ruthenium complex **1**<sup>9</sup> depending on the substituent is highly instrumental (eq 3).<sup>10</sup> On the basis of this observation, we envisioned that, if properly designed, an alternating metallotropic [1,3]-shift and enyne RCM could produce a single propagating carbene species that would travel along a linear poly-1,3-diyne ( $-[XCH_2CCCCCH_2]_n-$ ), generating an enediyne and  $\pi$ -conjugated oligoenynes (Scheme 1).<sup>11</sup> In this sequential enyne metathesis and metallotropic [1,3]-shift (M&M) event, the overall catalytic process initiated from one of the terminal alkenes would relay the metal carbene to the proximate alkyne.<sup>12,13</sup> Suppressed by strain from effecting a RCM reaction, the newly formed alkynyl carbene would undergo a [1,3]-shift, creating a new carbene effectively positioned for a facile metathetical closure to a five-membered ring. Herein we report one-step construction<sup>14</sup> of enediynes and oligoenynes enabled by the uniquely controlled repetitive metallotropic [1,3]-shift of ruthenium carbene species and its RCM.

First, the planned M&M sequence was examined in a general sense with the lowest homologue **2a** (Table 1, entry 1).<sup>15</sup> Treatment

**Scheme 1.** Sequential Metathesis and Metallotropy (M&M)




**Table 1.** Enyne Metathesis and Metallotropic [1,3]-Shift<sup>a</sup>

entry	ene-1,3-diyne	M&M product	yield (%) <sup>b</sup>
1	<b>2a</b>	<b>3a</b>	86 (67) <sup>c</sup>
2	<b>2b</b> , X = O	<b>3b</b> , X = O	63
3	<b>2c</b> , X = NTs	<b>3c</b> , X = NTs	47
4	<b>2d</b> , X = O, Y = O	<b>3d</b> , X = O, Y = O	51
5	<b>2e</b> , X = O, Y = NTs	<b>3e</b> , X = O, Y = NTs	45
6	<b>2f</b> , X = O, Y = O	<b>3f</b> , X = O, Y = O	32
7	<b>2g</b> , X = NTs, Y = O	<b>3g</b> , X = NTs, Y = O	36


<sup>a</sup> With catalyst **1** (5 mol %) in CH<sub>2</sub>Cl<sub>2</sub> (0.02 M) under reflux for 4–8 h.  
<sup>b</sup> Isolated yield. <sup>c</sup> Yield under ethylene atmosphere.

of **2a** with **1** (5 mol %) in CH<sub>2</sub>Cl<sub>2</sub> (40 °C) provided the desired product in 86% yield. The presence of ethylene during the reaction was found to be deleterious, which is contrary to the well-known beneficial ethylene effect in normal enyne metathesis.<sup>16</sup> It is also



**Figure 1.** X-ray structure of **3c**.

**Scheme 2.** M&M with an Alkyne-Terminating Group



surprising that the 1,3-diyne moieties in the substrate did not undergo CM with ethylene,<sup>17</sup> indicating that enyne RCM between the terminal alkene followed by the metallotropic [1,3]-shift and the terminating second RCM should be faster than CM of the propagating ruthenium carbene with ethylene. Next, a longer homologue with three diyne units between two terminal alkenes was tested. The all-oxygen-containing chain **2b** provided 63% yield of M&M product **3b** (entry 2), whereas the *N*-tosyl-group-bearing substrate **2c** afforded a lower yield (47%) of **3c** (entry 3). The symmetric nature of these product structures is apparent in their <sup>1</sup>H and <sup>13</sup>C NMR spectra, which was further confirmed by X-ray diffraction analysis of **3c** (Figure 1).

The effectiveness of this serial M&M sequence for the formation of oligoenynes is further demonstrated by using longer substrate homologues. Treatment of substrates **2d** and **2e** with catalyst **1** under typical RCM conditions provided *cis*-oligoenynes **3d** (X = O, Y = O) in 51% yield (entry 4) and **3e** (X = O, Y = NTs) in 45% yield (entry 5), respectively. The overall isolated yields of these longer homologue oligoenynes are marginal, and yet the theoretical yield for each bond-forming step is still higher than 90%. The longest homologues in the series, **2f** and **2g**, provided single isolated products **3f** and **3g** in 32% and 36% yield, respectively (entries 6 and 7), which remarkably involves the formation of five new triple bonds and six new double bonds in a single operation. Oligoenynes **3a–g** are stable even at room temperature over many weeks.<sup>18</sup>

An alkyne-terminating group for the metathesis can be accommodated to install a functionalizable group onto an enediyne product (Scheme 2). When **4a** was treated with **1** under ethylene,<sup>19</sup> RCM–[1,3]-shift–RCM reaction occurred to give **5a** (49%)<sup>20</sup> and its dimer **6a** (30%). Similarly, RCM of **4b** gave **5b** and **6b**. However, due to the instability of **5b**, the reaction mixture was directly treated with *N*-bromosuccinimide in pyridine, which allowed the isolation and characterization of **6b** (27%) and stable epoxide **7** (34%), respectively.

In conclusion, we have demonstrated that the combined use of enyne metathesis and metallotropic [1,3]-shift of alkynyl ruthenium carbenes is a powerful synthetic tool to construct enediyne and oligoenynes. The facile streamlining of the enyne metathesis and metallotropic [1,3]-shift of alkynyl ruthenium carbene may indicate

that the electronic and geometrical constraints for both processes are similar, yet further study is necessary to provide a more accurate mechanistic picture for the latter.<sup>21</sup> The reaction mechanism and the utility of this bond-forming process will be further explored.

**Acknowledgment.** We thank NSF (CHE-0401783) for financial support of this work as well as the NSF and NIH for NMR and mass spectrometry instrumentation. We also thank Prof. Wender for helpful discussions and Dr. Guzei for the X-ray structure determination of compound **3c**.

**Supporting Information Available:** General procedures and characterization of represented compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

**References**

- (1) Padwa, A.; Austin, D. J.; Gareau, Y.; Kassir, J. M.; Xu, S. L. *J. Am. Chem. Soc.* **1993**, *115*, 2637.
- (2) (a) Casey, C. P.; Dzwiniel, T. L. *Organometallics* **2003**, *22*, 5285. (b) Casey, C. P.; Dzwiniel, T. L.; Kraft, S.; Guzei, I. A. *Organometallics* **2003**, *22*, 3915. (c) Ortin, Y.; Sournia-Saquet, A.; Lugan, N.; Mathieu, R. *Chem. Commun.* **2003**, 1060.
- (3) (a) Casey, C. P.; Kraft, S.; Powell, D. R. *J. Am. Chem. Soc.* **2000**, *122*, 3771. (b) Casey, C. P.; Kraft, S.; Powell, D. R. *Organometallics* **2001**, *20*, 2651. (c) Casey, C. P.; Kraft, S.; Kavana, M. *Organometallics* **2001**, *20*, 3795. (d) Casey, C. P.; Kraft, S.; Powell, D. R. *J. Am. Chem. Soc.* **2002**, *124*, 2584.
- (4) Barluenga, H.; de la Ruá, R. B.; de Sáa, D.; Ballesteros, A.; Tomás, M. *Angew. Chem., Int. Ed.* **2005**, *44*, 4981.
- (5) (a) Kim, M.; Miller, R. L.; Lee, D. *J. Am. Chem. Soc.* **2005**, *127*, 12818. (b) van Otterlo, W. A. L.; Ngidi, E. L.; de Koning, C. B.; Fernandes, M. A. *Tetrahedron Lett.* **2004**, *45*, 659.
- (6) Reviews on metathesis: (a) Grubbs, R. H.; Chang, S. *Tetrahedron* **1998**, *54*, 4413. (b) Fürstner, A. *Angew. Chem., Int. Ed.* **2000**, *39*, 3012. (c) Schrock, R. R.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2003**, *42*, 4592. (d) Deiters, A.; Martin, S. F. *Chem. Rev.* **2004**, *104*, 2199.
- (7) Reviews on enyne metathesis: (a) Giessert, A. J.; Diver, S. T. *Chem. Rev.* **2004**, *104*, 1317. (b) Poulsen, C. S.; Madsen, R. *Synthesis* **2003**, 1. (c) Mori, M. In *Handbook of Metathesis*; Grubbs, R. H., Ed.; Wiley-VCH: Weinheim, Germany, 2003; Vol. 2, pp 176–204.
- (8) Review on cross metathesis: (a) Connon, S. J.; Blechert, S. *Angew. Chem., Int. Ed.* **2003**, *42*, 1900. A leading reference: Chatterjee, A. K.; Choi, T.-L.; Sanders, D. P.; Grubbs, R. H. *J. Am. Chem. Soc.* **2003**, *125*, 11360 and references therein.
- (9) Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. *Org. Lett.* **1999**, *1*, 953.
- (10) A stoichiometric reaction between **1** and diyne: Trnka, T. M.; Day, M. W.; Grubbs, R. H. *Organometallics* **2001**, *20*, 3845.
- (11) (a) Grissom, J. W.; Gunawardena, G. U.; Klingberg, D.; Huang, D. *Tetrahedron* **1996**, *52*, 6453. (b) Diederich, F. *Chem. Commun.* **2001**, 219. (c) Kosinski, C.; Hirsch, A.; Heinemann, F. W.; Hampel, F. *Eur. J. Org. Chem.* **2001**, 3879. (d) Tour, J. M. *Chem. Rev.* **1996**, *96*, 537. (e) Diederich, F. *Nature* **1994**, *369*, 199.
- (12) (a) Hansen, E. C.; Lee, D. *Org. Lett.* **2004**, *6*, 2035. (b) Hoyer, T. R.; Jeffrey, C. S.; Tennakoon, M. A.; Wang, J.; Zhao, H. *J. Am. Chem. Soc.* **2004**, *126*, 10210.
- (13) Computational studies of enyne metathesis mechanism: Lippstreu, J. J.; Straub, B. F. *J. Am. Chem. Soc.* **2004**, *127*, 7444.
- (14) Wender, P. A.; Miller, B. L. In *Organic Synthesis: Theory & Applications*; Hudlicky, T., Ed.; JAI: Greenwich, CT, 1993; Vol. 2, pp 27–66.
- (15) Preparation of diynes: Marino, J. P.; Nguyen, H. N. *J. Org. Chem.* **2002**, *67*, 6841. Details of substrate preparation are described in the Supporting Information.
- (16) (a) Kinoshita, A.; Sakakibara, N.; Mori, M. *J. Am. Chem. Soc.* **1997**, *119*, 12388. (b) Mori, M.; Sakakibara, N.; Kinoshita, A. *J. Org. Chem.* **1998**, *63*, 6082.
- (17) Selective cross metathesis of a *cis*-disubstituted double bond was observed in the presence of 1,3-diyne in the same molecule with Grubbs catalyst **1** under ethylene: Ratnayake, A. S.; Hemscheidt, T. *Org. Lett.* **2002**, *4*, 4667.
- (18) The origin of the low overall chemical yields for the longer homologues is not clear. However, on the basis of the appearance of a polar baseline material on the TLC even at the early stage of the reaction, we believe the starting material polymerizes under the reaction conditions.
- (19) Although ethylene was found to be deleterious for the RCM of substrate **2a**, it was used to minimize the formation of dimeric product **6a**.
- (20) Compound **5a** contains roughly 5% of styryl-group-incorporated product.
- (21) Putative intermediates in the metallotropic [1,3]-shift should have close analogy to the dehydrotungstenacyclobutadiene: McCullough, L. G.; Listemann, M. L.; Schrock, R. R.; Churchill, M. R.; Ziller, J. W. *J. Am. Chem. Soc.* **1983**, *105*, 6729.

JA057153C