



**Room Temperature Ring-Opening Metathesis of Pyridines by a Transient Ti≡C Linkage** [*J. Am. Chem. Soc.* 2006, 128, 6798–6799]. Brad C. Bailey, Hongjun Fan, John C. Huffman, Mu-Hyun Baik,\* and Daniel J. Mindiola\*

Page 6799. An incorrect dielectric constant was mistakenly used to simulate the pyridine solvent. When the correct dielectric constant  $\epsilon = 12.3$  is used, the reported solution phase energies



**Figure 2.** Proposed mechanism for the ROM of pyridine. Only core structures are shown for illustration. All energies are solution phase free energies, and all bond lengths are given in Å. [Ti] represents the (PNP)Ti framework, and solvation calculations were carried out at the optimized gas-phase geometry employing the dielectric constant of  $\epsilon = 12.3$  (pyridine).

change slightly. The corrected energies of all species discussed in the paper are summarized in the revised Figures 2 and 3. While these changes do not affect the conclusions and observed trends in a meaningful way, the authors apologize for the error.



**Figure 3.** Proposed mechanism of the ROM of pyridine. All energies are solution phase free energies. The dotted line represents the conversion of 7 directly to 8, as also depicted in Figure 2.

JA071777W

10.1021/ja071777w

Published on Web 05/16/2007