


## A Ferrous Center as Reaction Site for Hydration of a Nitrile Group into a Carboxamide in Mild Conditions

Nasser K. Thallaj, Juliette Przybilla, Richard Welter, and Dominique Mandon

*J. Am. Chem. Soc.*, **2008**, 130 (8), 2414-2415 • DOI: 10.1021/ja710560g

Downloaded from <http://pubs.acs.org> on February 8, 2009



### More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

## A Ferrous Center as Reaction Site for Hydration of a Nitrile Group into a Carboxamide in Mild Conditions

Nasser K. Thallaj,<sup>§</sup> Juliette Przybilla,<sup>§</sup> Richard Welter,<sup>||</sup> and Dominique Mandon\*<sup>§</sup>

Laboratoire de Chimie Biomimétique des Métaux de Transition and Laboratoire DECOMET, Institut de Chimie, UMR CNRS no. 7177-LC 3, Université Louis Pasteur, 4 rue Blaise Pascal, B.P. 1032, F-67070 Strasbourg cedex, France

Received November 28, 2007; E-mail: mandon@chimie.u-strasbg.fr


The mild and efficient hydration of nitriles in ecological-friendly conditions, a challenging research area, still far from being achieved by synthetic chemists,<sup>1–7</sup> is known in biological systems: Nature has designed the so-called nitrile hydratases (NHases), which are metal-containing proteins that are key players in the process of assimilation of nitriles upon conversion to amides.<sup>8,9</sup> The presence of Co(III) or Fe(III) ions centers is essential to the activity of the enzyme, and a generally accepted hypothesis involves the interaction of the substrate and reagent with the metal site.<sup>10–14</sup> The detailed mechanism of the reaction, however, is far from being completely understood, especially the question as to whether water molecule or the nitrile, or both, coordinates to the metal in order to be activated.

In our efforts to develop electron-deficient iron complexes with tripodal tetraamine ligands,<sup>15–17</sup> we synthesized the cyanide- $\alpha$ -substituted tripod CNTPA,  $L_1 = (6\text{-cyano } 2\text{-pyridylmethyl})\text{bis}(2\text{-pyridylmethyl})\text{amine}$ , and report that under anaerobic and wet conditions, the dichloroferrous complex  $L_1\text{FeCl}_2$  converts to  $L_2\text{-FeCl}_2$ , with  $L_2 = \text{H}_2\text{NCOTPA}$ , (6-carboxamido 2-pyridylmethyl)-bis(2-pyridylmethyl)amine. Decomplexation of  $L_2\text{FeCl}_2$  affords the free  $L_2$  ligand, that has incorporated  $^{18}\text{O}$  when  $\text{H}_2^{18}\text{O}$  is used. This reaction represents to our knowledge the first example of the hydration of a nitrile carried out by a ferrous ( $\text{Fe}^{\text{II}}$ ) center to yield the corresponding carboxamide. We also demonstrate that the nitrile group does not need to be coordinated to react with water.


The preparation of  $L_1$  and  $L_1\text{FeCl}_2$  follows straightforward procedures.<sup>18,15</sup>  $L_1\text{FeCl}_2$  is a red thermally stable and moderately oxygen-sensitive compound displaying an  $\text{Fe}^{\text{II}} \Rightarrow$  ligand MLCT transition at  $\lambda = 389 \text{ nm}$  ( $\epsilon = 1.96 \times 10^3 \text{ mmol}\cdot\text{cm}^2$ ) in its UV–vis spectrum in line with a tetradentate coordination mode of the ligand.<sup>15,19</sup> Both its  $^1\text{H}$  NMR (well resolved paramagnetically shifted signals) and the molecular conductivity data ( $\Lambda = 13.8 \text{ S}\cdot\text{mol}^{-1}\cdot\text{cm}^2$  at  $10^{-3} \text{ M}$ ) support a pseudo-octahedral geometry in solution.<sup>15</sup> IR spectroscopy shows  $\nu_{\text{CN}}$  at  $2238 \text{ cm}^{-1}$ . Single crystals of  $L_1\text{FeCl}_2$  could be obtained. As expected from solution studies, the ligand coordinates in a tetradentate fashion around the metal, and the two chloride ions complete the coordination sphere. The ORTEP diagram is displayed in Figure 1.

Contacts between the nitrile group and the iron and equatorial chloride are evidenced by  $d_{\text{C18-Fe}} = 3.54 \text{ \AA}$ ,  $d_{\text{C19-Cl1}} = 3.70 \text{ \AA}$ , and  $d_{\text{N5-Cl1}} = 3.69 \text{ \AA}$ . The  $\text{C}_{18}\text{C}_{19}\text{N}_5$  segment is bent with  $\angle\text{C}_{18}\text{C}_{19}\text{N}_5 = 173.8^\circ$ . This provides evidence for a repulsive interaction between the equatorial chloride ligands and the CN group, the latter being probably destabilized.

$L_1\text{FeCl}_2$  is thermally stable in anaerobic solution for days. The addition of 100 equiv of degassed water to a solution of  $L_1\text{FeCl}_2$  resulted in minor yet detectable changes in the UV–visible



**Figure 1.** ORTEP drawing of  $L_1\text{FeCl}_2$ . Selected bond lengths ( $\text{\AA}$ ) and angles (deg): Fe–N3 2.202(4); Fe–N1 2.215(4); Fe–N2 2.266(4); Fe–N4 2.312(4); Fe–Cl1 2.3496(15); Fe–Cl2 2.4324(15); N3–Fe–Cl2 166.97(12); N1–Fe–N4 146.20(17); N2–Fe–Cl1 171.22(12); C18–C19–N5 178.8(6). The insert shows contacts between the iron center and C<sub>19</sub>, equatorial chloride Cl<sub>1</sub> and the nitrile substituent, and tilting of the C<sub>18</sub>C<sub>19</sub>N<sub>5</sub> segment.



**Figure 2.** ORTEP drawing of the  $[L_2\text{FeCl}]^+$  cation. Selected bond lengths ( $\text{\AA}$ ) and angles (deg): Fe–N2 2.1339(16); Fe–O1 2.1443(16); Fe–N4 2.1913(18); Fe–N5 2.2195(19); Fe–N3 2.2430(18); Fe–Cl1 2.3328(6); O1–C1 1.259(3); N1–C1 1.329(3); N2–Fe–Cl1 173.00(5); N5–Fe–N4 151.37(7); N3–Fe–O1 150.97(6); N1–C1–O1 121.9(2).


spectrum over 24–40 h, the  $\text{Fe}^{\text{II}} \Rightarrow$  ligand MLCT being shifted to  $\lambda = 396 \text{ nm}$  ( $\epsilon = 1.82 \times 10^3 \text{ mmol}\cdot\text{cm}^2$ ), with an isobestic point at  $\lambda = 412 \text{ nm}$ . When the reaction temperature was lowered to  $0^\circ\text{C}$ , no change could be detected, even in the presence of more added water.<sup>20</sup> In  $^1\text{H}$  NMR, the formation of a new high-spin ferrous species over 24 h was observed. An amido carbonyl group was detected in IR with  $\nu_{\text{C=O}} = 1679 \text{ cm}^{-1}$ . The new  $L_2\text{FeCl}_2$  complex could be obtained in a single-crystalline form. Its ORTEP diagram is displayed on Figure 2.

Obviously, the nitrile substituent has been converted into an amide group, with  $d_{\text{C1O1}} = 1.258 \text{ \AA}$  and  $d_{\text{C1N1}} = 1.329 \text{ \AA}$ . Unlike in  $L_1\text{FeCl}_2$ , the substituted pyridyl arm lies trans to the chloride, and the oxygen of the amide group coordinates to the iron center trans to the tertiary amine of the tripod with  $d_{\text{O1-Fe}} = 2.14 \text{ \AA}$ . The

<sup>§</sup>Laboratoire de Chimie Biomimétique des Métaux de Transition.



<sup>||</sup>Laboratoire DECOMET.

**Scheme 1.** Proposed Mechanism for the Hydration of the Nitrile Substituent upon Addition of Water to  $L_1FeCl_2$ <sup>a</sup>



<sup>a</sup> Crystal structures are provided for  $L_1FeCl_2$ ,  $L_2FeCl_2$ .

**Scheme 2.** Hydration of the Nitrile Substituent of  $L_1 = CNTPA$  Leading to  $L_2 = H_2NCOTPA$ , upon Temporary Complexation to Iron Dichloride in Wet Conditions<sup>a</sup>



<sup>a</sup> Insert: ESMS, when  $H_2^{18}O$  is used;  $m/z = 358.14$  for  $L_2(^{18}O) + Na^+$ .

complex is thus cationic, the uncoordinated chloride ion lying far away from the metal center with  $d_{FeCl} = 6.28 \text{ \AA}$ .

$L_1$  as a free ligand is indefinitely stable in the presence of water, and  $L_1FeCl_2$  also is indefinitely stable, but in the absence of water. Thus, coordination of water to the ferrous center must occur prior to any further reaction, as displayed in Scheme 1. The first step corresponds to the dissociation of the chloride from the metal in the presence of water, followed by coordination of water. In  $L_2FeCl_2$  the fact that the substituted pyridyl arm lies trans to the chloride, that is, at a different position than in the starting material  $L_1FeCl_2$ , strongly suggests rearrangement of the coordination polyhedron, and coordination of water at the metal site as a necessary step in the conversion of  $L_1$  to  $L_2$ . The detection of an isobestic point in UV-vis during hydration reaction impeded detection of any intermediate.

Treatment of  $L_1FeCl_2$  with  $H_2O$  during 48 h at room temperature followed by acidification with  $CF_3CO_2H$  under inert atmosphere resulted in complete bleaching of the medium. Extraction of the organic phase under standard conditions with  $CH_2Cl_2$  afforded  $L_2$  as a white solid together with variable amounts (however, never more than 20% based on NMR) of  $L_1$ .  $L_2$  was characterized by  $^1H$  and  $^{13}C$  NMR, IR, and mass spectroscopy. In the  $^{13}C$  NMR, the  $\delta_{CN} = 117.3$  ppm resonance was replaced by a new signal at  $\delta_{CO} = 167.0$  ppm, and IR displayed a new absorption at  $\nu_{CO} = 1685$

$cm^{-1}$ . MS yielded the molecular ion at  $m/z = 334.15$  ( $L_2H^+$ ) and  $356.14$  ( $L_2Na^+$ ) as the main signal. As displayed in Scheme 2, the use of  $H_2^{18}O$  resulted in incorporation of  $^{18}O$  in  $L_2$  with  $m/z = 336.16$  ( $L_2[^{18}O]H^+$ ) and  $358.14$  ( $L_2[^{18}O]Na^+$ ).

The involvement of a ferrous center in non-redox chemistry to promote the conversion of functional groups is unusual.<sup>21,22</sup> To our knowledge, the present communication is the first reported example of hydration of a nitrile function by a ferrous complex. In addition, we demonstrate that the smooth hydration reaction does not necessarily require the nitrile to be coordinated, but activated in the vicinity of a metal-coordinated water molecule. Enhanced Lewis acidity at the metal center might also contribute to what could be described as an anchimer effect, leading to nitrile hydration by a simple "outer-sphere mechanism".<sup>9,14</sup> Comparative studies with complexes of the same ligand with other transition metals are currently under investigation in our laboratory.

**Acknowledgment.** We wish to thank Dr Remy Louis, head of the Institut de Chimie in Strasbourg, for constant encouragements and support, and the CNRS and ULP. The Conseil Scientifique de l'ULP is acknowledged for specific support no. AO CS ULP 2006.

**Supporting Information Available:** All synthetic details, including the preparation and characterization of all compounds mentioned; experimental details for the hydration reaction. For the two reported structures, the crystallographic files in CIF format have been deposited. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- Zil'berman, E. N. *Russ. Chem. Rev.* **1984**, *53*, 900–912.
- Dopp, D.; Dopp, H., Eds. *Methoden der Organischen Chemie (Houben-Weyl)*; Thieme: Stuttgart, Germany, 1985; Vol. E5 (2), pp 1024–1031.
- Brown, R. R., Ed.; *The Organic Chemistry of Aliphatic Nitrogen Compounds*; Oxford University Press: Oxford, 1994; pp 217–221; 342–346.
- Bauer, W., Jr. In *Ullmann's Encyclopedia of Industrial Chemistry*, 5th ed.; John Wiley & Sons: New York, 1990; Vol. A16, p 441.
- Kukushkin, V. Y.; Pombeiro, A. J. L. *Chem. Rev.* **2002**, *102*, 1771–1802.
- Yamaguchi, K.; Matsushita, M.; Mizuno, N. *Angew. Chem., Int. Ed.* **2004**, *43*, 1576–1580 and references therein.
- Jiang, X. B.; Minnaard, A.; Feringa, B. L.; De Vries, J. G. *J. Org. Chem.* **2004**, *69*, 2327–2331 and references therein.
- Kobayashi, M.; Shimizu, S. *Curr. Opin. Chem. Biol.* **2000**, *4*, 95–102.
- Mascharak, P. K. *Coord. Chem. Rev.* **2002**, *225*, 201–214.
- Miyayaga, A.; Fushinobu, S.; Ito, K.; Wakagi, T. *Biochem. Biophys. Res. Commun.* **2001**, *288*, 1169–1174.
- Arakawa, T.; Kawano, Y.; Kataoka, S.; Katayama, Y.; Kamiya, N.; Yohda, M.; Odaka, M. *J. Mol. Biol.* **2007**, *366*, 1497–1509.
- Nagashima, S.; Nakasako, M.; Dohmae, N.; Tsujimura, M.; Takio, K.; Odaka, M.; Yohda, M.; Kamiya, N.; Endo, I. *Nat. Struct. Biol.* **1998**, *5*, 347–351.
- Song, L.; Wang, M.; Shi, J.; Xue, Z.; Wang, M.-X.; Qian, S. *Biochem. Biophys. Res. Commun.* **2007**, *362*, 319–324.
- Huang, W.; Jia, J.; Cummings, J.; Nelson, M.; Schneider, G.; Lindqvist, Y. *Structure* **1997**, *5*, 691–699.
- Mandon, D.; Machkour, A.; Goetz, S.; Welter, R. *Inorg. Chem.* **2002**, *41*, 5364–5372.
- Machkour, A.; Mandon, D.; Lachkar, M.; Welter, R. *Inorg. Chem.* **2004**, *43*, 1545–1550.
- Thallaj, N. K.; Machkour, A.; Mandon, D.; Welter, R. *New. J. Chem.* **2005**, *29*, 1555–1558.
- Nelson, M. S.; Rodgers, J. *J. Chem. Soc. A* **1968**, 272–276.
- Machkour, A.; Mandon, D.; Lachkar, M.; Welter, R. *Inorg. Chim. Acta* **2005**, *358*, 839–843.
- Addition of a large excess of water resulted in precipitation of the starting material in the UV-vis cuvette.
- See for instance: Jain, R.; Hao, B.; Liu, R.-P.; Chan, M. K. *J. Am. Chem. Soc.* **2005**, *127*, 4558–4559.
- With a diferric complex, see: Hazell, A.; Jensen, K.; McKenzie, C. J.; Toftlund, H. *Inorg. Chem.* **1994**, *33*, 3127–3134.

JA710560G