


Ni(II) and Co(II) Sensing by Escherichia coli RcnR

Jeffrey S. Iwig, Sharon Leitch, Robert W. Herbst, Michael J. Maroney, and Peter T. Chivers
J. Am. Chem. Soc., **2008**, 130 (24), 7592-7606 • DOI: 10.1021/ja710067d • Publication Date (Web): 28 May 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Ni(II) and Co(II) Sensing by *Escherichia coli* RcnRJeffrey S. Iwig,[†] Sharon Leitch,[‡] Robert W. Herbst,[‡] Michael J. Maroney,[‡] and Peter T. Chivers^{*†}*Washington University School of Medicine, Department of Biochemistry and Molecular Biophysics, Saint Louis, Missouri 63110, and University of Massachusetts, Department of Chemistry, Amherst, Massachusetts 01003*

Received November 6, 2007; E-mail: chivers@wustl.edu

Abstract: *Escherichia coli* RcnR and *Mycobacterium tuberculosis* CsoR are the founding members of a recently identified, large family of bacterial metal-responsive DNA-binding proteins. RcnR controls the expression of the metal efflux protein RcnA only in response to Ni(II) and Co(II) ions. Here, the interaction of Ni(II) and Co(II) with wild-type and mutant RcnR proteins is examined to understand how these metals function as allosteric effectors. Both metals bind to RcnR with nanomolar affinity and stabilize the protein to denaturation. X-ray absorption and electron paramagnetic resonance spectroscopies reveal six-coordinate high-spin sites for each metal that contains a thiolate ligand. Experimental data support a tripartite N-terminal coordination motif (NH₂-Xaa-NH-His) that is common for both metals. However, the Ni(II)- and Co(II)-RcnR complexes are shown to differ in the remaining coordination environment. Each metal coordinates a conserved Cys ligand but with distinct M-S distances. Co(II)-thiolate coordination has not been observed previously in Ni(II)/Co(II)-responsive metalloregulators. The ability of RcnR to recruit ligands from the N-terminal region of the protein distinguishes it from CsoR, which uses a lower coordination geometry to bind Cu(I). These studies facilitate comparisons between Ni(II)-RcnR and NikR, the other Ni(II)-responsive transcriptional regulator in *E. coli*, to provide a better understanding how different nickel levels are sensed in *E. coli*. The characterization of the Ni(II)- and Co(II)-binding sites in RcnR, in combination with bioinformatics analysis of all RcnR/CsoR family members, identified a four amino acid fingerprint that likely defines ligand-binding specificity, leading to an emerging picture of the similarities and differences between different classes of RcnR/CsoR proteins.

Introduction

Metal-responsive DNA-binding proteins that control transporter and enzyme gene expression in bacteria are well-known.¹ These metalloregulators have been divided into several structural families, named after their founding members: MerR, ArsR/SmtB, DtxR, Fur, NikR, and CopY.¹ Each family, with the exception of NikR, contains subfamilies that respond to distinct metals or other small molecules. For example, the MerR family includes the Cu-responsive CueR,² the Zn-responsive ZntR,³ and the Pb-responsive PbrR.⁴ Less well-understood are the structural bases for metal-sensing and metal-specific function by these proteins in the context of the complex intracellular milieu. Specificity is important to avoid promiscuous activation by noncognate metals. An additional structural challenge is present when two regulatory proteins that control opposing functional pathways must respond to the same metal ion, but at different concentrations, such as the zinc-responsive regulatory proteins Zur and ZntR⁵ that control influx and efflux pathways, respectively. Structural characterization of the metal-binding

sites provides insight into their thermodynamic and kinetic properties, which provide a basis for understanding their function in the cell.

The present understanding of metalloregulator structure-function relationships indicates that regulation of DNA-binding activity by the cognate metal is generally allosteric, so that the proper conformational change required for function is linked to the metal-binding specificity. The metal ion selectivity is at least partly achieved by the coordination number/geometry of the metal-protein complex,⁶ and by ligand selection, as not all metals that bind *in vitro* elicit a biological response. These features have been demonstrated for the Ni(II)-responsive transcriptional repressor, NikR, from *Escherichia coli*, which has been shown to bind to a number of metals with a high affinity⁷ but responds *in vivo* only to the presence of Ni(II) ions.⁸

Recently, we identified a nickel- and cobalt-responsive metalloregulator, *E. coli* RcnR,⁹ that represses the transcription of *rcnA*, which encodes a nickel and cobalt efflux protein.¹⁰ RcnR exhibits only minimal sequence identity with the previously described families of transcriptional regulators, leading

[†] Washington University School of Medicine.[‡] University of Massachusetts.(1) Giedroc, D. P.; Arunkumar, A. I. *Dalton Trans.* **2007**, (29), 3107–20.(2) Outten, F. W.; Outten, C. E.; Hale, J.; O'Halloran, T. V. *J. Biol. Chem.* **2000**, *275* (40), 31024–9.(3) Xiong, A.; Jayaswal, R. K. *J. Bacteriol.* **1998**, *180* (16), 4024–9.(4) Borremans, B.; Hobman, J. L.; Provoost, A.; Brown, N. L.; van Der Lelie, D. *J. Bacteriol.* **2001**, *183* (19), 5651–8.(5) Outten, C. E.; O'Halloran, T. V. *Science* **2001**, *292* (5526), 2488–92.(6) Pennella, M. A.; Shokes, J. E.; Cospoer, N. J.; Scott, R. A.; Giedroc, D. P. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100* (7), 3713–8.(7) Wang, S. C.; Dias, A. V.; Bloom, S. L.; Zamble, D. B. *Biochemistry* **2004**, *43* (31), 10018–28.(8) Leitch, S.; Bradley, M. J.; Rowe, J. L.; Chivers, P. T.; Maroney, M. J. *J. Am. Chem. Soc.* **2007**, *129*, 5085–95.

to the proposal of a new metalloregulator family.⁹ RcnR homologues are found throughout the eubacterial kingdom, emphasizing their widespread role in regulating the cell physiology of microbes from diverse growth niches. Despite the lack of a known DNA-binding motif, RcnR represses transcription by directly binding to the *rcnA* promoter. This interaction is negatively affected by Ni(II) or Co(II) binding, a pattern functionally similar to that of the structurally distinct SmtB/ArsR family of transcription factors (reviewed in ref 11). The subsequent functional and structural characterization of *Mycobacterium tuberculosis* CsoR, a Cu(I)-responsive homologue of RcnR, confirmed the discovery of a new structural class of metalloregulators (RcnR/CsoR family) characterized by an all α -helical structure anchored by a four-helix bundle.¹² The discovery of this new family has been reviewed recently.¹

The nickel and cobalt requirements of *E. coli* are distinct. The nickel physiology of *E. coli* is dominated by the anaerobic expression of Ni-Fe hydrogenases,^{13,14} which obtain their nickel via the NikABCDE nickel permease.^{15,16} The cobalt physiology of this organism is less well-understood. *E. coli* cannot synthesize adenosylcobalamin (vitamin B₁₂) *de novo* but has a specific transport pathway for this essential cofactor.^{17–20} Cobalt ions are thought to enter the cell only through transporters with a low specificity such as the magnesium transporter, CorA.^{20–22} These observations suggest that Co(II) ions are regarded by the cell as toxic, one manifestation of which is its adverse effect on Fe-S clusters in enzymes such as aconitase and ferrichrome reductase.²³

Physiologically, the *rcnR-rcnA* module is functionally linked to other nickel-dependent activities of the cell.⁹ At low extracellular nickel levels, low or transient expression of RcnA is required to prevent premature *nikABCDE* repression by NikR.⁹ This effect likely results from a competition between nickel efflux by RcnA and nickel-binding and transcriptional repression by NikR. Under steady state growth conditions, the NikR repressor responds to lower ambient Ni(II) concentrations than RcnR, as *nikABCDE* expression is completely repressed at nickel levels that exhibit minimal *rcnA* expression.⁹ As Ni(II) levels increase, RcnR repression of *rcnA* transcription is

diminished, allowing full induction of RcnA expression to prevent excess metal accumulation in the cell. The complementary responses for the nickel-dependent regulators in *E. coli* raise the question as to how these two proteins are structurally tuned to respond to different nickel levels in the cell.

To better understand ligand recognition and allosteric regulation in this new family of metalloregulators, we investigated binding of the cognate metals, Ni(II) and Co(II), to the RcnR protein. Ni(II) and Co(II) are similar in size and ligand preference, but as to whether they bind to RcnR at the same site and with the same coordination environment is unknown. Ni(II) and Co(II) do not commonly adopt a trigonal planar geometry (two cysteines and one histidine) observed in the Cu(I)-CsoR structure,¹² suggesting that the RcnR protein must supply additional metal-binding ligands. X-ray absorption (XAS) and UV-vis spectroscopies revealed a six-coordinate metal-binding site for Ni(II) or Co(II) containing one S-donor and five N/O-donor atoms. Studies of RcnR mutants show that Ni(II) and Co(II) use overlapping but distinct binding sites to inhibit RcnR DNA binding.

Experimental Procedures

Molecular Biology. All chemicals were purchased from Fisher Scientific (Pittsburgh, PA) except where indicated. Oligonucleotides (Operon Biotechnologies, Huntsville, AL), plasmids, and bacterial strains used in this study are listed in the Supporting Information (Table S8). Several *rcnR-P_{rcnA}-lacZ* mutations and pRcnR mutations were constructed using the following methods: (1) The QuikChange protocol (Stratagene, La Jolla, CA) was used with slight modification. For each 50 μ L reaction, 200 ng of plasmid DNA and 150 ng of each primer were used, with 2.5 units of *Pfu* Turbo DNA polymerase (Stratagene, La Jolla, CA). DNA was then subjected to digestion with 20 units of *Dpn* I (New England Biolabs, Beverly, MA) for 1.5 h at 37 °C. After heat inactivation for 10 min, 5 μ L of the reaction was transformed into *E. coli* JM109 cells following the TSS protocol.²⁴ (2) Additional *rcnR-P_{rcnA}-lacZ* mutations and pRcnR mutations were constructed by overlap extension PCR²⁵ and ligation into pPC163 or pET-22b as described previously.⁹ (3) Mutations near the 5' and 3' regions of *rcnR* were constructed by simple PCR amplification of the *rcnR* gene using a mutant primer (Supporting Information Table S7), followed by subcloning into the appropriate plasmid. All mutant constructs were sequenced (Seqwright, Houston, TX) to verify that only the desired mutation was present.

RcnR Overexpression and Purification. *E. coli* DL41 (DE3)²⁶ cells harboring plasmid encoding wild-type or mutant RcnR proteins were grown in LB media containing ampicillin (200 μ g/mL, 538 μ M) at 37 °C to an OD₆₀₀ of ~1.0, prior to the addition of IPTG (120 mg/L, 0.5 mM) to induce RcnR expression. The cells were harvested after 3 h by centrifugation (4600g, 25 min), resuspended in residual media (~5 mL/L culture), and frozen at -80 °C. Thawed cell pellets were lysed by sonication, at 4 °C in the presence of 5 mM tris(2-carboxyethyl)-phosphine hydrochloride (TCEP) (Pierce) and 1.5 mM phenylmethylsulfonylfluoride (Sigma), a serine protease inhibitor. Subsequent steps were all performed at 20 °C. After centrifugation, the lysate supernatant was applied to an SP Sepharose column (18 mL) equilibrated with 20 mM Hepes (pH 7.0), 1.0 mM TCEP, 5.0 mM EDTA, and 10% glycerol (buffer A), with 50 mM NaCl. The column was then washed with 40 mL of buffer A with 50 mM NaCl followed by a linear gradient of 0.05–1

- (9) Iwig, J. S.; Rowe, J. L.; Chivers, P. T. *Mol. Microbiol.* **2006**, *62* (1), 252–62.
- (10) Rodrigue, A.; Effantin, G.; Mandrand-Berthelot, M. A. *J. Bacteriol.* **2005**, *187* (8), 2912–6.
- (11) Busenlehner, L. S.; Pennella, M. A.; Giedroc, D. P. *FEMS Microbiol. Rev.* **2003**, *27* (2–3), 131–43.
- (12) Liu, T.; Ramesh, A.; Ma, Z.; Ward, S. K.; Zhang, L.; George, G. N.; Talaat, A. M.; Sacchettini, J. C.; Giedroc, D. P. *Nat. Chem. Biol.* **2007**, *3* (1), 60–8.
- (13) Ballantine, S. P.; Boxer, D. H. *J. Bacteriol.* **1985**, *163* (2), 454–9.
- (14) Sawers, R. G.; Ballantine, S. P.; Boxer, D. H. *J. Bacteriol.* **1985**, *164* (3), 1324–31.
- (15) Wu, L. F.; Mandrand-Berthelot, M. A.; Waugh, R.; Edmonds, C. J.; Holt, S. E.; Boxer, D. H. *Mol. Microbiol.* **1989**, *3* (12), 1709–18.
- (16) Wu, L. F.; Navarro, C.; Mandrand-Berthelot, M. A. *Gene* **1991**, *107* (1), 37–42.
- (17) Lawrence, J. G.; Roth, J. R. *Genetics* **1996**, *142* (1), 11–24.
- (18) Roth, J. R.; Lawrence, J. G.; Bobik, T. A. *Annu. Rev. Microbiol.* **1996**, *50*, 137–81.
- (19) Cadieux, N.; Bradbeer, C.; Reeger-Schneider, E.; Koster, W.; Mohanty, A. K.; Wiener, M. C.; Kadner, R. J. *J. Bacteriol.* **2002**, *184* (3), 706–17.
- (20) de Veaux, L. C.; Clevenson, D. S.; Bradbeer, C.; Kadner, R. J. *J. Bacteriol.* **1986**, *167* (3), 920–7.
- (21) Gibson, M. M.; Bagga, D. A.; Miller, C. G.; Maguire, M. E. *Mol. Microbiol.* **1991**, *5* (11), 2753–62.
- (22) Wu, L. F.; Navarro, C.; de Pina, K.; Quenard, M.; Mandrand, M. A. *Environ. Health Perspect.* **1994**, *102*, 297–300.
- (23) Ranquet, C.; Ollagnier-de-Choudens, S.; Loiseau, L.; Barras, F.; Fontecave, M. *J. Biol. Chem.* **2007**, *282* (42), 30442–51.

- (24) Chung, C. T.; Niemela, S. L.; Miller, R. H. *Proc. Natl. Acad. Sci. U.S.A.* **1989**, *86* (7), 2172–5.
- (25) Ho, S. N.; Hunt, H. D.; Horton, R. M.; Pullen, J. K.; Pease, L. R. *Gene* **1989**, *77* (1), 51–9.
- (26) Hendrickson, W. A.; Horton, J. R.; LeMaster, D. M. *EMBO J.* **1990**, *9* (5), 1665–72.

M NaCl in buffer A (110 mL total volume, flow rate 2 mL/min, collected in 2 mL fractions). RcnR eluted at ~400 mM NaCl, and these fractions were pooled and concentrated at ~4000g to 1 mL using 2 mL 5 kDa MW cutoff Vivaspin concentrators (Vivascience, Hanover, Germany) and applied to a HiLoad 26/60 Superdex 200 column equilibrated with buffer A containing 300 mM NaCl. RcnR eluted as a single peak, at a volume most consistent with a tetramer. Fractions containing RcnR were pooled and diluted to 25 mL and applied to an S6 ion exchange column (2 cm × 6 cm) (Biorad, Hercules, CA) running on a Biorad Biologic FPLC. The column was then washed with 20 mL of buffer A with 50 mM NaCl, and the protein was eluted using a linear gradient of 0.05–1 M NaCl (48 mL total volume, 4 mL/min flow rate). RcnR eluted between 490 and 620 mM NaCl. Purified RcnR was stored at 4 °C in 20 mM Hepes (pH 7.0), 500 mM NaCl, 1 mM TCEP, 1 mM EDTA, and 5% glycerol for up to 4 weeks; thereafter, the DNA-binding specificity and affinity began to decrease. The identity of the purified protein was confirmed by N-terminal sequencing using Edman degradation (Midwest Analytical, St. Louis, MO). Prior to use in different experiments, RcnR samples were desalted 2 times into buffer M [10 mM Hepes (pH 7.0), 150 mM NaCl, and 5% glycerol] using Microbiospin 6 columns (Biorad) to remove EDTA and TCEP present in the storage buffer. Modifications to this approach are noted where relevant.

RcnR Characterization. An experimentally determined extinction coefficient of $\epsilon_{276} = 2530 \text{ M}^{-1} \text{ cm}^{-1}$ was determined by measuring the absorbance of an aliquot of purified denatured RcnR and determining its concentration by amino acid analysis (Texas A&M Protein Chemistry Laboratory, College Station, TX). The free sulfhydryl content of native RcnR was quantified by a standard thiol titration method²⁷ using 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB). Samples contained RcnR (5–10 μM) in 100 mM sodium phosphate (pH 7.3). Quantification of the sulfhydryl concentration was obtained using a TNB extinction coefficient of $14\,150 \text{ M}^{-1} \text{ cm}^{-1}$ at 412 nm using a Shimadzu UV-2401PC spectrophotometer.²⁸

Metal Content Assays. For PAR assays, 3–4 monomer equivalents of NiCl_2 or CoCl_2 was added to RcnR protein previously desalted into buffer M, and the samples were incubated at 20 °C for 2 h. Weakly bound and excess free metal ions were removed by a second passage over a Microbiospin 6 column equilibrated with buffer M.

PAR Assay. 4-(2-Pyridylazo) resorcinol (PAR) experiments followed an established protocol^{29,30} with minor modifications. Standard curves for Ni(II) and Co(II) in complex with PAR were generated from samples containing 1–9 μM metal and 20 μL of 1 mg/mL PAR in 120 μL total volume [50 mM Hepes (pH 7.3) and 6 M guanidine hydrochloride]. The absorbance was measured for each sample (Ni, 505 nm and Co, 540 nm) after equilibration for 1 h at 20 °C. RcnR samples were prepared similarly, using two samples at two different protein concentrations between 1 and 9 μM . The metal content reported was the average of all four samples.

Atomic Absorption Spectroscopy (AAS). Samples for AAS were prepared in a similar manner to those stated previously except that protein was desalted into buffer M lacking glycerol to remove excess metal ions. Aliquots (100 μL) were then incubated overnight (20 °C) in 2% nitric acid, then diluted 10-fold with H_2O to a final volume of 1 mL. Samples were analyzed on a PerkinElmer AAnalyst 600 graphite furnace atomic absorption spectrophotometer equipped with a nickel hollow cathode lamp (Department of Biology, Washington University). Nickel content was quantified

by comparison to a standard curve of 2–100 $\mu\text{g/L}$ nickel. Each sample was quantified 3 times and averaged.

UV–vis Spectroscopy. Metal-bound RcnR UV–vis spectra were measured on a Shimadzu UV-2401PC spectrophotometer. Metals were added from a stock solution to individual protein samples (previously desalted) for each nickel or cobalt concentration. Samples (100 μL volume) were allowed to equilibrate for 15 h at 20 °C under aerobic conditions after metal addition before spectral collection (1 cm path length, 0.2 nm step size).

For iodoacetamide (IAM) (Sigma, St. Louis, MO) treated samples, 5 mM IAM was added to RcnR in storage buffer (2 h at 20 °C). Control samples with water added instead of IAM were prepared in parallel. The samples were then desalted twice into buffer M before metal addition.

Mag-Fura 2 and Fura 2 Metal Titrations. Fluorescence measurements were all taken on a Cary Eclipse fluorescence spectrophotometer in a 150 μL , 1 cm path length Hellma quartz cuvette. A range of concentrations of NiCl_2 or CoCl_2 was incubated with 0.5 μM mag-fura 2 (Molecular Probes, Carlsbad, CA) or 0.3 μM fura 2, respectively, for 3 h in the presence or absence of the indicated concentrations of RcnR. Fluorescence measurements were made using $\lambda_{\text{ex}} = 380 \text{ nm}$ and $\lambda_{\text{em}} = 505 \text{ nm}$ for mag-fura 2 and $\lambda_{\text{ex}} = 360 \text{ nm}$ and $\lambda_{\text{em}} = 500 \text{ nm}$ for fura 2, with both excitation and emission slit widths set to 10 nm. Each reported value was the average of three readings. Data were fit to a four independent-binding sites model using Scientist version 2.01 (Micromath, St. Louis, MO).

Analytical Ultracentrifugation. Sedimentation equilibrium data were obtained on a Beckman XL-A ultracentrifuge using six-cell sample holders (1.2 cm path length). Samples of 110 μL of RcnR (5 and 25 μM in buffer M plus 200 μM EDTA) were centrifuged at 25 000 and 35 000 rpm (20 °C) with a 120 μL buffer blank. A_{230} or A_{276} measurements were obtained after 10 h and every 2 h thereafter until three consecutive scans overlapped (20 h total) to ensure that equilibrium had been achieved. The partial specific volume for RcnR (0.7504 mL/g at 20 °C) was determined from the amino acid content using SEDNTERP.³¹ A buffer density of 1.019 g/mL was also calculated from SEDNTERP. The SEDPHAT software package was used to globally analyze data obtained at multiple rotor speeds and protein concentrations.³²

Circular Dichroism (CD). CD experiments were carried out on a JASCO 710 spectropolarimeter using 1 cm (urea melts) or 1 mm (wavelength scans) path length cylindrical quartz cuvettes. Wavelength scans (20 °C, speed 20 nm/min, 1 nm step) were performed on samples containing 15 μM RcnR in 10 mM Hepes (pH 7.0), 100 mM NaCl, and 5% glycerol. Thirty micromolar Ni(II) or Co(II) was added where indicated. Samples were equilibrated for 3 h (20 °C) before scans. Each spectrum was the average of three accumulations. The raw, blank subtracted signal (mdeg) was converted to mean residue ellipticity (MRE) and fraction helix, as previously described.^{33,34}

Urea denaturation samples were performed as described previously.³⁵ A freshly prepared stock of 10 M urea in buffer M was used to make individual samples (900 μL) containing different urea concentrations, which were allowed to equilibrate for 12–15 h at 20 °C. Ellipticity data for each urea concentration was collected for 60 s at 230 nm in 1 s intervals and averaged. Because the pathway for unfolding could not be unambiguously determined with respect to the oligomeric state, urea concentrations that caused half-

(27) Creighton, T. E. *Protein Structure: A Practical Approach*; Oxford University Press: Oxford, 1989; pp 155–67.

(28) Riddles, P. W.; Blakeley, R. L.; Zerner, B. *Methods Enzymol.* **1983**, *91*, 49–60.

(29) Siemann, S.; Brewer, D.; Clarke, A. J.; Dmitrienko, G. I.; Lajoie, G.; Viswanatha, T. *Biochim. Biophys. Acta* **2002**, *1571* (3), 190–200.

(30) Hunt, J.; Neece, S.; Ginsburg, A. *Anal. Biochem.* **1985**, *146*, 150–7.

(31) Laue, T. M.; Shah, B. D.; Ridgeway, T. M.; Pelletier, S. L. *Analytical Ultracentrifugation in Biochemistry and Polymer Science*; Royal Society of Chemistry: Cambridge, U.K., 1992; pp 90–125.

(32) Vistica, J.; Dam, J.; Balbo, A.; Yikilmaz, E.; Mariuzza, R. A.; Rouault, T. A.; Schuck, P. *Anal. Biochem.* **2004**, *326* (2), 234–56.

(33) Myers, J. K.; Pace, C. N.; Scholtz, J. M. *Biochemistry* **1997**, *36* (36), 10923–9.

(34) Rohl, C. A.; Chakrabarty, A.; Baldwin, R. L. *Protein Sci.* **1996**, *5* (12), 2623–37.

(35) Chivers, P. T.; Sauer, R. T. *Chem. Biol.* **2002**, *9* (10), 1141–8.

maximal unfolding are reported based on fits to a model of folded monomer to unfolded monomer.³⁶

X-ray Absorption Spectroscopy (XAS). Samples of RcnR were desalted and metallated as described previously. The samples were then concentrated to 1.5–3.2 mM in 20 mM Hepes (pH 7.0), containing either 300 mM NaCl or NaBr, prior to data collection. XAS data for all RcnR samples were collected at beamline 9-3 at the Stanford Synchrotron Radiation Laboratory (SSRL) under dedicated ring conditions. Each sample was syringed into a polycarbonate cuvette that was wrapped in kapton tape and frozen in liquid nitrogen. Data were collected at 10 K using a liquid helium cryostat (Oxford Instruments). The ring conditions were 3 GeV, and the current was 80–100 mA for each sample. The beamline optics consist of a Si(220) double crystal monochromator ($\phi = 90^\circ$), a flat rhodium coated mirror upstream of the monochromator for harmonic rejection and vertical collimation, and a toroidal focusing mirror after the monochromator. X-ray fluorescence was detected with a 30 element Ge detector (Canberra). The energy of each K_α edge was calibrated to the first inflection point of the corresponding foil: Co, 7709.5 eV and Ni, 8331.6 eV. X-ray absorption near edge spectroscopy (XANES) data were collected from ± 200 eV relative to the edge. Analysis of the XANES region was carried out by fitting a cubic function to the baseline in the pre-edge region of the spectrum and using a 75% Gaussian and 25% Lorentzian function to fit the rise in fluorescence occurring at the edge. Gaussian functions were added to fit each pre-edge transition, and the areas of the inserted Gaussians were taken to be the peak area.⁸ Extended X-ray absorption fine structure (EXAFS) images were collected to 13.5 k above the absorption edge (E_0) for WT Co–RcnR and 16.3 k above the absorption edge energy (E_0) for both C35A RcnR complexes and each wild-type Ni–RcnR. XAS data reduction was carried out using EXAFS123 with graphics produced in Igor Pro (WaveMetrics version 5.0) as previously described.⁸ For EXAFS analysis, four to seven scans were averaged to generate the summed EXAFS spectrum. Each summed spectrum was normalized and background corrected using a three-section cubic spline for the pre-edge and postedge regions. The data were converted to k -space using the relationship $[2m_e(E - E_0)/\hbar^2]^{1/2}$ (where m_e is the mass of the electron and E_0 is the threshold energy above the absorption edge: 7723 eV for Co and 8340 eV for Ni). Least-squares fitting of EXAFS arising from the first coordination sphere was performed using Fourier-filtered spectra with transform limits of 2.0–12.5 \AA^{-1} for WT RcnR samples and 2.0–14.0 \AA^{-1} for C35A RcnR samples, as determined by the signal/noise ratio in the data and to allow appropriate comparisons. A back-transform window of 1.0–2.3 \AA (uncorrected for phase shifts) was used to model scattering atoms in the first coordination sphere. Fits were calculated for coordination numbers four, five, and six, holding the number of scattering atoms in a shell to integer values and using all combinations of S- and N-donors (see Supporting Information). Best fits were judged by minimizing the goodness-of-fit parameter (GOF) and deviation of σ^2 values from typical values. The back-transform window was expanded to 1.0–4.0 \AA (uncorrected for phase shifts) to accommodate scattering from imidazole rings, which were fit using multiple scattering parameters as previously described.⁸ Histidine ligand counting was achieved by adding integer numbers of imidazole rings to the fit and observing the effect on GOF and σ^2 and provides an estimate of the number of imidazole ligands in model compounds with an error of $\pm 25\%$ (about the same as the error in N , the number of first shell atoms). The final fits included in the text were selected from the best fits of Fourier-filtered data and were further refined using unfiltered data. For all fits, r (metal–scatterer distance), σ^2 (disorder parameter), and E_0 (phase shift) were allowed to refine; however, they were constrained so that only one value of E_0 was used and only one value of σ^2 per type of scattering atom was employed.




Figure 1. Divalent metal induction of P_{rcnA} expression. RZ4500 cells were grown with pJ114 in the presence of the indicated divalent metal ions as described in the Experimental Procedures. LacZ activity was assayed as previously described.⁹ Induction from P_{rcnA} only occurs in the presence of Ni or Co. Relative LacZ activity of 1 corresponds to 136 Miller units. Error bars show the SD of the LacZ activity of three separate samples.

EPR Spectroscopy. EPR spectra were obtained with a Bruker Elexsys 500 spectrometer equipped with an ER 4102 single cavity operating at the X-band (9.46 GHz). Measurements were performed at 12 K using an Oxford Instruments EPR-9000 liquid helium continuous flow cryostat. EPR spectra were simulated using the matrix diagonalization program Xsophe (Bruker Biospin). The spin Hamiltonian employed was: $\mathbf{H} = \beta g \mathbf{H} \mathbf{S} + \mathbf{S} \mathbf{D} \mathbf{S}$.

β -Galactosidase Reporter Experiments. β -Galactosidase reporter experiments were set up as described previously with the following modification.⁹ Starter cultures of *E. coli* strain PC563 ($\Delta rcnR$) cells containing WT or mutant $rcnR$ genes on the $rcnR$ - P_{rcnA} - $lacZ$ plasmid (pJ114) were grown aerobically in Luria–Bertani broth (LB) with chloramphenicol (34 mg/L, 0.1 mM). These cultures were used to inoculate triplicate cultures of 1.87 mL of LB with chloramphenicol (34 mg/L, 0.1 mM) in capped microfuge tubes (starting OD₆₀₀ of 0.000001). Cultures were then grown for 12–15 h at 37 °C and assayed as previously described.⁹

For metal induction experiments, cells were treated in a similar manner except pJ114 was in the $\Delta lacZ$ *E. coli* strain RZ4500.³⁷ Cells were grown anaerobically at the maximal concentration for each individual divalent metal that resulted in <10% inhibition of growth (measured by final OD₆₀₀), which was as follows: 30 μ M MnCl₂, 1.5 mM FeSO₄, 150 μ M CoCl₂, 500 μ M NiCl₂, 100 μ M CuCl₂, 300 μ M ZnCl₂, and 10 μ M CdSO₄.

Results

RcnR Metal Specificity *in vivo*. Previous studies demonstrated that Ni(II) and Co(II) induced transcription from a P_{rcnA} - $lacZ$ fusion construct.⁹ To test the effect of other divalent transition metal ions on P_{rcnA} - $lacZ$ expression, cells were grown anaerobically in LB media containing the maximal metal ion concentrations that inhibited growth by <10%. As shown in Figure 1, Mn(II), Fe(II), Cd(II), Cu(II), and Zn(II) all had no detectable effect on $lacZ$ expression. Divalent metal concentrations that caused up to 50% growth inhibition failed to activate expression from P_{rcnA} more than 2-fold (data not shown). These results establish Ni(II) and Co(II) as the only two physiologically relevant metal ions for $rcnA$ induction under these growth conditions and led us to study their interactions with the RcnR protein in more detail.

RcnR Purification and Characterization. Native *E. coli* RcnR was purified to homogeneity (~ 5 mg/L culture) by a combination of anion exchange and gel filtration chromatographies (see Experimental Procedures). The N-terminal amino acid sequence

(36) Santoro, M. M.; Bolen, D. W. *Biochemistry* **1988**, *27* (21), 8063–8.

(37) Choe, M.; Reznikoff, W. S. *J. Bacteriol.* **1991**, *173* (19), 6139–46.




Figure 2. Secondary structure and stability of *E. coli* RcnR measured by CD spectroscopy. (A) Far UV CD spectra of 15 μM RcnR without metal (black line) or with 30 μM NiCl_2 (orange line) or CoCl_2 (blue line). Protein samples were in 10 mM Hepes (pH 7.0), 100 mM NaCl, and 5% glycerol. Data were converted to mean residue ellipticity as described.^{33,34} (B) Urea denaturation (monitored at 230 nm) of apo-, Ni(II)-, or Co(II)-substituted RcnR (2.5 μM). NiCl_2 or CoCl_2 were added to a 5 μM final concentration. Protein samples were in buffer M.

of the protein (SHTIR; see also Supporting Information Figure S10) revealed the expected cleavage of the N-terminal methionine,³⁸ resulting in an 89-residue native protein (the numbering used here is based on the *rcnR* gene sequence). Using the PAR assay for transition metal ion detection,³⁰ purified RcnR contained <0.04 metal equivalents per monomer. Thiol titration with DTNB indicated that the free sulfhydryl content of natively folded RcnR was 0.9 ± 0.1 (expected was 1.0).

The CD spectrum of purified RcnR (Figure 2A) showed minima at 222 and 208 nm, consistent with a folded, α -helical protein. Conversion of the mean residue ellipticity to fraction helix gives 61% helix, in reasonable agreement with calculations of 68–78% helix from secondary structure prediction algorithms.³³ The protein secondary structure showed little perturbation in the presence of NiCl_2 or CoCl_2 (Figure 2A), indicating a minimal net change in secondary structure due to metal binding.

The RcnR protein unfolded cooperatively as a function of increasing urea concentration (Figure 2B). RcnR showed a concentration-dependent stability, with transition midpoints of 2.7 M urea and 3.2 M urea for 2 and 12 μM protein, respectively, indicating that oligomer dissociation and RcnR unfolding are coupled (Supporting Information Figure S1). Metal addition increased RcnR stability, shifting the midpoint of the denaturation curve from 2.7 to 5.0 M urea for Ni and 4.4 M urea for Co (Figure 2B).

Assembly State of RcnR. Sedimentation equilibrium experiments were performed to rigorously characterize the oligomeric state of the protein and to determine as to whether metal binding caused any changes in the RcnR assembly state. For the apoprotein, data sets were fit globally to a single species model, yielding a molecular mass of 39 399 Da (Supporting Information Figure S2), consistent with an RcnR tetramer (expected mass 40 010 Da). Residuals for this fit did not show systematic deviations, and inclusion of a dimeric species in the model did not improve the quality of the fit because the best fit for the tetramer–dimer dissociation constant was >2 orders of magnitude less than the protein concentrations used. Thus, negligible amounts of dimer ($<10\%$) were present under these solution conditions. Likewise, the addition of an octameric species did not improve data fitting. Experiments also were carried out in the presence of 200 μM NiCl_2 or 200 μM CoCl_2 instead of EDTA, and under these conditions, RcnR remained tetrameric (see Supporting Information Figure S2). Similarly, the addition of 1 mM DTT (see Supporting Information Figure S2), higher pH (7.5), or increased salt (300 mM NaCl) did not affect the species distribution. Thus, under these experimental conditions, RcnR formed a stable tetramer, and this tetrameric state was not perturbed by Ni(II) or Co(II) binding, suggesting that the tetrameric form is the active biological species. However, dimer or monomer species may occur under solution conditions different than those tested here.

Electronic Absorption Spectroscopy. The Ni(II)– and Co(II)–RcnR complexes were analyzed by UV–vis spectroscopy (Figure 3A) and exhibited features consistent with six-coordinate metal centers featuring thiolate ligation. The difference spectra (holo–apo) for 1:1 M(II)–RcnR (monomer equivalents) complexes exhibited intense absorptions for each metal [$M = \text{Co(II)}$, $\epsilon_{314} = 2664 \text{ M}^{-1} \text{ cm}^{-1}$ and $\epsilon_{238} = 4675 \text{ M}^{-1} \text{ cm}^{-1}$ and $M = \text{Ni(II)}$, $\epsilon_{280} = 1778 \text{ M}^{-1} \text{ cm}^{-1}$ and $\epsilon_{235} = 3937 \text{ M}^{-1} \text{ cm}^{-1}$]. The feature at 310 nm in the Co(II)–RcnR spectrum is not present if RcnR is preincubated with the thiol modifying agent iodoacetamide (5 mM), while the peak at 235 nm is unchanged, indicating that the longer wavelength feature is due to thiolate to Co(II) ligand-to-metal charge transfer (LMCT) (Supporting Information Figure S3). Similarly, for Ni(II)–RcnR, the 280 nm feature is nearly absent when RcnR is preincubated with iodoacetamide, while the 231 nm peak maintains a constant intensity, indicating that the peak at 280 nm is due to $\text{CysS} \rightarrow \text{Ni(II)}$ LMCT (Supporting Information Figure S3). These assignments also are supported by EXAFS analyses of Co(II)– and Ni(II)–RcnR and mutagenesis data that confirm S coordination (*vide infra*), as well as the high molar extinction coefficients observed for the UV–vis absorptions and by similar absorptions in Co-substituted proteins^{39,40} and the few examples of six-coordinate Ni(II) thiolate complexes that are known.^{41,42}

Thus, a preliminary interpretation of the observed LMCT spectra is that the Co(II)–RcnR complex exhibits a $\text{S} \rightarrow \text{M(II)}_{\pi}$ LMCT at 310 nm, while the $\text{S} \rightarrow \text{M(II)}_{\sigma}$ transition at higher energy is not observed. For Ni(II), the observed transition at 280 nm must be due to $\text{S} \rightarrow \text{M(II)}_{\sigma^*}$ since no other transition is

(38) Frotin, F.; Martinez, A.; Peynot, P.; Mitra, S.; Holz, R. C.; Gigliome, C.; Meinel, T. *Mol. Cell Proteomics* **2006**, *5* (12), 2336–49.

(39) Brown, D. C.; Collins, K. D. *J. Biol. Chem.* **1991**, *266* (3), 1597–604.

(40) May, S. W.; Kuo, J. Y. *Biochemistry* **1978**, *17* (16), 3333–8.

(41) Ragunathan, K. G.; Bharadwaj, P. K. *J. Chem. Soc., Dalton Trans.* **1992**, 2417–22.

(42) Rosenfield, S. G.; Berends, H. P.; Gelmini, L.; Stephan, D. W.; Mascharak, P. K. *Inorg. Chem.* **1987**, *26* (17), 2792–7.




Figure 3. Electronic spectroscopy of Ni or Co-substituted RcnR. (A) UV-vis difference spectra of 1:1 Ni(II)-RcnR (60 μ M) (orange line) and 1:1 Co(II)-RcnR (85 μ M) (blue line) were obtained in buffer M. A metal-free sample was subtracted to obtain the difference spectrum. Data were collected every 0.5 nm. (B) Stoichiometric addition of NiCl₂ to RcnR (60 μ M). (C) Stoichiometric addition of CoCl₂ to RcnR (88 μ M). Absorbance signal saturated at 1.0 Co/monomer and 1.2 Ni/monomer under the same solution conditions as in panel A.

possible, and the energy relative to the Co(II) case reflects the higher energy of the transition moderated by the stabilization of the metal 3d-orbitals with respect to the sulfur 2p-orbitals that occurs with increasing the effective nuclear charge on the metal.

Metal-Binding Stoichiometry of RcnR. Independent experimental methods were used to determine the metal-binding stoichiometry. Titration of 60 μ M RcnR with NiCl₂ and 88 μ M with CoCl₂ revealed a saturation of the absorbance signals that corresponded to M(II)-RcnR complexes with stoichiometries of 1.2 Ni(II)/monomer (Figure 3B) and 1.0 Co/monomer (Figure 3C), or about four metals per tetramer. Because the absorbance change upon metal binding was linear at these concentrations, no accurate determination of binding affinities could be made. The number of metal-binding sites per RcnR monomer also was determined using the colorimetric divalent metal indicator, PAR (described in the Experimental Procedures). The stoichiometries were found to be 0.7 ± 0.1 Co(II)/RcnR monomer and 1.2 ± 0.1 Ni(II)/RcnR monomer. The nickel content also was confirmed using AAS, which yielded 1.03 ± 0.01 Ni(II)/RcnR




Figure 4. Ni(II)- and Co(II)-binding affinities determined by competition assay. (A) Mag-fura 2 (0.5 μ M) fluorescence ($\lambda_{\text{ex}} = 380$ nm and $\lambda_{\text{em}} = 505$ nm) in absence or presence of RcnR monomer (500 nM) and indicated concentrations of NiCl₂. The solid line represents the fit to a model of four independent binding sites per RcnR tetramer ($K_d < 25 \pm 4$ nM). (B) Fura 2 (0.3 μ M) fluorescence ($\lambda_{\text{ex}} = 360$ nm and $\lambda_{\text{em}} = 500$ nm) in the absence or presence of RcnR monomer (360 nM) and the indicated concentrations of CoCl₂. The solid line is a fit to a four independent binding site model ($K_d < 5 \pm 2$ nM).

monomer. These data are all consistent with one metal-binding site per RcnR monomer or four binding sites per tetramer.

Metal Affinities. To more accurately define metal-binding affinities, chelator competition assays were performed. It has been previously determined that mag-fura 2 fluorescence is quenched upon 1:1 complex formation with Ni(II).^{43,44} To apply this approach to RcnR, the affinities for the dye itself were first determined by direct titration of 0.5 μ M mag-fura 2 with NiCl₂ in buffer M. Data were fit to a single site model, giving an affinity of 330 nM for Ni(II). The value is slightly higher than the published value of 130 nM, which was carried out under different solution conditions.⁴³

Titration were then performed with 0.5 μ M mag-fura 2 with 500 nM RcnR monomer. The presence of RcnR resulted in a shift in the Ni-dependent fluorescence quenching curve (Figure 4A), indicating competition for Ni binding. However, the shape of the nickel-binding curve indicated that the protein was nearly fully saturated before quenching of mag-fura 2 began. Therefore, only a lower limit of the affinity could be determined from this experiment ($K_d < 25 \pm 4$ nM; four noninteracting sites model).

Similarly, to examine Co(II) interactions, CoCl₂ titrations were performed with 600 nM RcnR and 300 nM fura 2. Titration with fura 2 alone resulted in an affinity of 201 nM for the Co

(43) Golynskiy, M. V.; Gunderson, W. A.; Hendrich, M. P.; Cohen, S. M. *Biochemistry* **2006**, *45* (51), 15359–72.

(44) Kwan, C. Y.; Putney, J. W., Jr *J. Biol. Chem.* **1990**, *265* (2), 678–84.




Figure 5. *K*-edge XAS spectra of Ni(II)–RcnR (left): WT (orange) and Cys35Ala (brown) and Co(II)–RcnR (right): WT (light blue), and Cys35Ala (dark blue) in 300 mM NaBr buffer. (Top) XANES spectra: Insets are enlargements of the peaks associated with the $1s \rightarrow 3d$ and $1s \rightarrow 4p_z$ transitions. (Bottom) Unfiltered EXAFS data and fits (black). Fits shown are from Table 1 and correspond to fits 5 and 10 for WT and Cys35Ala Ni(II)–RcnR and fits 15 and 20 for WT and Cys35Ala Co(II)–RcnR, respectively.

fura 2 complex (1:1 complex expected).⁴⁵ When RcnR was added, a noticeable shift in the cobalt titration curve was observed (Figure 4B), but again, only a lower limit of the affinity of RcnR for Co(II) (5 ± 2 nM) could be determined by this method.

Metal Site Structure. XANES. XAS spectroscopy of Ni(II)– and Co(II)–RcnR was used to obtain structural information about the metal coordination environment. Ni(II)- and Co(II)-substituted samples of a Cys35Ala RcnR mutant also were studied to explore S coordination at the metal site. Nickel and cobalt *K*-edge XANES analysis provides information about the coordination number and geometry at each site.⁴⁶ In the WT Ni(II)–RcnR complex, the pre-edge XANES image exhibits only one small pre-edge transition at ~ 8330 eV (peak area = 0.036(7) eV), associated with a $1s \rightarrow 3d$ electronic transition (Figure 5). In the absence of any other pre-edge features, this spectrum is diagnostic for a six-coordinate, pseudo-octahedral Ni(II) center.⁴⁷ The spectrum of Cys35Ala Ni(II)–RcnR shows a clear change in the intensity of the $1s \rightarrow 3d$ transition (peak area = 0.070(16) eV) and a new feature, a shoulder at ~ 8338 eV, associated with a $1s \rightarrow 4p_z$ transition (Figure 5). The increased intensity of the $1s \rightarrow 3d$ transition and the observation of a second pre-edge feature are both consistent with a five-coordinate, square pyramidal nickel center.⁴⁷

The XAS spectrum of Cys35Ala Ni(II)–RcnR (and no other sample) exhibited a slight change with each sequential scan, indicating some exposure time-dependent radiation damage. The

shift in the Ni *K*-edge energy and changes in the XANES and EXAFS spectra during scans 1–5 (used for the summed file) are too slight to interpret as a specific change in oxidation state or ligand environment (Supporting Information Figure S4). The peak area of the $1s \rightarrow 3d$ transition increases from 0.056(25) eV for scan 1 to 0.067(21) eV for scan 5, which are indistinguishable and lie well within the range observed for five-coordinate Ni(II) centers.⁴⁷

In the WT Co(II)–RcnR complex, the pre-edge XANES spectrum features a single small pre-edge transition at ~ 7710 eV (peak area = 0.097(4) eV) that is associated with a $1s \rightarrow 3d$ electronic transition (Figure 5). The small peak area and the absence of any other pre-edge features are consistent with a six-coordinate, distorted octahedral geometry. While the peak area is slightly larger than usual for octahedral Co(II) complexes, it is significantly smaller than the peaks observed for five-coordinate or tetrahedral Co(II) complexes.⁴⁸ A slight increase in the $1s \rightarrow 3d$ peak area also was observed in other mixed-ligand, distorted octahedral Co(II) centers.⁸ The XANES spectrum obtained for Cys35Ala Co(II)–RcnR is very similar to that obtained for the WT complex, with only one notable pre-edge feature at ~ 7710 eV (Figure 5). The peak area (0.075(6) eV) is slightly smaller than the corresponding feature in the spectrum of the WT cobalt center and is also consistent with a six-coordinate site.

EXAFS. EXAFS analysis provides information about the types of scattering atoms and metric details about the complex.⁴⁶ The results of EXAFS analyses on the RcnR samples are summarized in Figure 5 and Table 1, which contains five example fits for the EXAFS spectrum of each sample. Three of these fits represent the best fits of scattering arising from the

(45) Gryniewicz, G.; Poenie, M.; Tsien, R. Y. *J. Biol. Chem.* **1985**, *260* (6), 3440–50.

(46) Scott, R. A. Measurement of metal–ligand distances by EXAFS. In *Enzyme Structure, Part J*; Hirs, C. H. W., Timasheff, S. N., Eds.; Academic Press: San Diego, 1985; Vol. 117, pp 414–459.

(47) Colpas, G. J.; Maroney, M. J.; Bagyinka, C.; Kumar, M.; Willis, W. S.; Suib, S. L.; Baidya, N.; Mascharak, P. K. *Inorg. Chem.* **1991**, *30* (5), 920–8.

(48) Padden, K. M.; Krebs, J. F.; MacBeth, C. E.; Scarrow, R. C.; Borovik, A. S. *J. Am. Chem. Soc.* **2001**, *123* (6), 1072–9.

Table 1. EXAFS Fits for RcnR Complexes

sample/fit	N^a	r (Å) ^b	σ^2 ($\times 10^3$ Å ⁻²) ^c	ΔE_0 (eV)	GOF ^d
Ni(II)–RcnR	Fourier-filtered fits with a backtransform window of 1–2.3 Å				
1	6 Ni–N/O	2.05(2)	10.4(17)	7.6(21)	0.66
2	5 Ni–N/O	2.07(12)	7.7(12)	10.4(15)	0.48
	1 Ni–N/O	2.34(350)	[7.7] ^e	[10.4]	
3	5 Ni–N/O	2.06(1)	8.1(7)	9.5(9)	0.31
	1 Ni–S	2.57(2)	8.2(27)	[9.5]	
	Fourier-filtered fits with a backtransform window of 1–4 Å				
4	2 Ni–N _{imd}	1.99(2)	3.4(19)	10.6(9)	0.55
	3 Ni–N/O	2.12(1)	[3.4]	[10.6]	
	1 Ni–S	2.55(2)	7.5(22)	[10.6]	
	unfiltered EXAFS fit from 2 to 12.5 Å ⁻¹				
5	2 Ni–N _{imd}	1.986(5)	3.3(6)	9.8(3)	0.70
	3 Ni–N/O	2.112(3)	[3.3]	[9.8]	
	1 Ni–S	2.543(7)	7.4(8)	[9.8]	
C35A Ni–RcnR	Fourier-filtered fits with a backtransform window of 1–2.3 Å				
6	5 Ni–N	1.87(1)	5.7(10)	2.5(27)	0.76
7	4 Ni–N/O	1.88(1)	4.1(5)	6.6(17)	0.43
	1 Ni–N/O	2.15(2)	[4.1]	[6.6]	
8	4 Ni/N/O				no fit
	1 Ni–S				
	Fourier-filtered fits with a backtransform window of 1–4 Å				
9	1 Ni–N _{imd}	2.31(3)	4.1(5)	3.0(17)	0.65
	4 Ni–N/O	1.87(1)	[4.1]	[3.0]	
	unfiltered EXAFS fit from 2 to 14.0 Å ⁻¹				
10	1 Ni–N _{imd}	2.316(8)	4.1(2)	4.4(4)	0.72
	4 Ni–N/O	1.876(2)	[4.1]	[4.4]	
Co(II)–RcnR	Fourier-filtered fits with a backtransform window of 1–2.3 Å				
11	6 Co–N/O	2.05(3)	14.8(19)	–0.9(21)	0.44
12	5 Co–N/O	2.05(200)	14.8(32)	–0.9(27)	0.48
	1 Co–N/O	2.06(1000)	[14.8]	[–0.9]	
13	5 Co–N/O	1.97(2)	16.4(15)	–8.0(13)	0.16
	1 Co–S	2.25(1)	6.6(10)	[8.0]	
	Fourier-filtered fits with a backtransform window of 1–4 Å				
14	2 Co–N _{imd}	1.94(3)	6.6(27)	–2.8(16)	0.51
	3 Co–N/O	2.11(3)	[6.6]	[–2.8]	
	1 Co–S	2.24(4)	13.1(60)	[–2.8]	
	4 Co–C _{imd}	3.01(4)	13.4(60)	[–2.8]	
	unfiltered EXAFS fit from 2 to 12.5 Å ⁻¹				
15	2 Co–N _{imd}	1.945(12)	6.9(13)	–3.0(7)	0.76
	3 Co–N/O	2.113(14)	[6.9]	[–3.0]	
	1 Co–S	2.238(18)	12.6(28)	[–3.0]	
	4 Co–C _{imd}	3.009(19)	13.5(28)	[–3.0]	
C35A Co–RcnR	Fourier-filtered fits with a backtransform window of 1–2.3 Å				
16	6 Co–N/O	1.92(3)	15.6(29)	–5.2(48)	0.62
17	4 Co–N/O	1.92(2)	6.7(11)	–2.8(23)	0.43
	2 Co–N/O	2.12(3)	[6.7]		
18	5 Co–N/O				no fit
	1 Co–S				
	Fourier-filtered fits with a backtransform window of 1–4.0 Å				
19	2 Co–N _{imd}	1.87(7)	14.6(66)	–1.6(21)	0.66
	4 Co–N/O	1.97(4)	[14.6]	[–1.6]	
	4 Co–C _{imd}	2.89(5)	12.6(66)	[–1.6]	
	unfiltered EXAFS fit from 2 to 14.0 Å ⁻¹				
20	2 Co–N _{imd}	1.867(23)	14.3(22)	–1.6(6)	0.74
	4 Co–N/O	1.975(11)	[14.3]	[–1.6]	
	4 Co–C _{imd}	2.886(18)	13.2(24)	[–1.6]	

^a N is the number of scattering atoms at a specific distance. ^b r (Å) is the radial distance. ^c σ^2 is the root-mean-squared disorder in the Ni–X distance, where X is the scattering atom. ^d GOF (goodness-of-fit) was calculated as described previously.⁴⁸ The accuracy of distance determined by EXAFS for atoms in the first coordination sphere of the metal was limited to ± 0.02 Å by theoretical phase parameters; numbers in parentheses represent SD from the least-squares fits. ^e Values of parameters in brackets were constrained to be equal to those of other scattering atoms in the same shell.

first coordination sphere atoms (Fourier-filtered data with a backtransform window of 1.0–2.3 Å, not corrected for phase shifts) and include the best single shell fit and best two shell fits with and without an S-donor ligand. In the fourth fit, data are included to allow the determination of the presence of histidine ligands

and an estimate of the number of histidine imidazoles bound to be made (Fourier-filtered data with a back-transform window of 1.0–4.0 Å, not corrected for phase shifts). The fifth fit is a refinement against unfiltered EXAFS data. These data and best fits to unfiltered data are shown in Figure 5. Additional fits of

Fourier-filtered data are included in the Supporting Information (Figure S5 and associated Tables).

The EXAFS spectrum arising from the first coordination sphere of the metal in WT Ni(II)–RcnR is best modeled as a six-coordinate site composed of two shells of ligands, including one S scattering atom (Figure 5 and Table 1). The number of scattering atoms in the first coordination sphere is in agreement with the XANES analysis of the Ni(II) site (*vide supra*). Addition of a S-donor at a long distance (2.57(2) Å) improves the GOF by 35% over the best fit composed of two shells of N/O scattering atoms. Data obtained from samples prepared in buffers containing 300 mM NaBr or 300 mM NaCl were essentially identical (Supporting Information Figure S6), confirming the assignment of a S scattering atom and not a Cl[−] ion from the buffer. Analysis of the EXAFS data including scattering atoms in the second and third coordination sphere confirms the presence of histidine ligation (the best fit with histidine ligands has a GOF that is 15% better than the best fit obtained without imidazole parameters using the same back-transform limits) and an estimate of two to three histidine ligands. Refinement of the models against unfiltered data led to similar conclusions, with the fit obtained for two histidine ligands having the lowest GOF value (Supporting Information Tables S1 and S3). Fits obtained using either Fourier-filtered or unfiltered data clearly show a substantial improvement upon inclusion of a long Ni–S vector (Table 1 and Supporting Information Tables S1 and S3). The EXAFS analysis is therefore consistent with a Ni(His)₂(Cys)(N/O)₃ site in WT Ni(II)–RcnR.

The EXAFS spectrum obtained for Cys35Ala Ni(II)–RcnR further supports the inclusion of an S-donor in the coordination sphere of WT Ni(II)–RcnR and identifies the ligand involved as Cys35. The first coordination sphere EXAFS data are best modeled by a single type of scatterer, N/O-donors (Table 1). Splitting the coordination sphere into two shells improves the GOF by 43%. The number of N/O ligands determined for the first coordination sphere is five, which is also in agreement with the XANES analysis. Multiple scattering analysis of data including second and third coordination sphere atoms is consistent with one histidine ligand, although the improvement over the fit obtained without imidazole parameters is modest (7%). There is no evidence in the data for a Br[−] ion from the buffer coordinating to the Ni(II) center. The EXAFS analysis of Cys35Ala Ni(II)–RcnR is therefore consistent with a Ni(N/O)₅ site, where the number of histidine ligands is not clear. This might be due to the large distribution in Ni–NHis distances (static disorder) that occurs as a result of the loss of C35, a possibility that is consistent with the range of Ni–N/O distances observed (1.87–2.31 Å).

The analysis of first coordination sphere EXAFS data for Co(II)–RcnR also reveals the presence of a S-donor ligand. The addition of a single S-donor to a shell of five N/O-donors greatly improves the GOF as compared to a fit containing six N/O-donor ligands (67%, Table 1). Again, data obtained in 300 mM NaCl or NaBr containing buffers gave indistinguishable results, confirming the assignment of an S-donor ligand. The total number of ligands (six) agrees well with the XANES analysis (*vide supra*). Analysis of the spectrum using multiple scattering parameters to account for histidine ligation is consistent with the presence of *ca.* two histidine ligands (15% improvement in GOF). The structure that emerges from the EXAFS analysis for WT Co(II)–RcnR is Co(His)₂(Cys)(N/O)₃, a result that is similar to the Ni site structure in WT Ni(II)–RcnR. Since there is only one Cys ligand present in *E.*

coli RcnR, the two metals both bind Cys35 in six-coordinate sites that also contain at least two imidazole ligands. The largest difference apparent in the EXAFS spectra of the two metal sites is the M–S distance, which is short for Co(II) (2.24(2) Å) and long for Ni(II) (2.54(2) Å).

The EXAFS arising from first coordination sphere scattering atoms in the Cys35Ala Co(II)–RcnR center is best modeled by six N/O-donors at two different distances (Table 1). Inclusion of a S-donor ligand causes the fit to diverge. Multiple scattering analysis of the Co(II) site is consistent with the presence of 2 ± 1 imidazole ligands. Thus, the perturbation in the metal site for the Cys35Ala mutants is distinct. In the case of Ni(II), the site is five-coordinate, reflecting the loss of the S-donor. For Co(II), the site binds an unknown sixth N/O-donor ligand. Both Cys35Ala sites have poorly refined data for histidine ligation.

EPR. Analysis of the XAS data from both Ni(II)– and Co(II)–RcnR is consistent with six-coordinate, M(Cys)(His)₂(N/O)₃ complexes but differs strikingly in the M(II)–S distance found for the Cys ligand (2.24(2) Å for Co(II) and 2.54(2) Å for Ni(II)). Six-coordinate Ni(II) complexes are invariably high-spin, *S* = 1 complexes, and the metal–ligand distances found for the Ni(II)–RcnR complex are consistent with this electronic configuration.⁴⁹ Both high-spin (*S* = 3/2) and low-spin (*S* = 1/2) Co(II) complexes are common but are easily distinguished by EPR spectroscopy.⁵⁰ The EPR spectrum of Co(II)–RcnR collected at 12 K shows features that are typical of an *S* = 3/2 Co(II) center (Supporting Information Figure S7). Integration of the EPR spectrum shows an absorption curve that lies on a flat baseline, indicating that no saturation effects are seen. The experimental spectrum can be simulated by a high-spin, *S* = 3/2 Co center with *M_s* = 1/2, *D* ≫ *hν* (50 cm^{−1} was used), *g_{x,y}* = 2.210, *g_z* = 2.195, and *E/D* = 0.07. These values are typical for high-spin five- or six-coordinate Co(II) in protein–ligand environments.⁵¹

Identification of Metal-Sensing Determinants by Amino Acid Substitutions. Site-directed mutagenesis of the *rcnR* gene (pJII14) was carried out to gain functional insight into the amino acids that form the metal-binding site, as suggested by spectroscopic measurements and sequence conservation. The regulation of RcnA expression by the RcnR mutants was tested *in vivo* using the previously described LacZ assay.⁹ A mutation that reduced metal binding should generate a constitutively active repressor and result in low LacZ activity, even in the presence of Ni(II) or Co(II). Importantly, any mutation that caused the protein to be poorly expressed, unstable, or unable to bind DNA would lead to high LacZ activity in both the presence and the absence of these metal ions. Mutated RcnR residues included all five histidine residues (three are highly conserved), a subset of conserved Glu and Asp residues, and the highly conserved Cys35 (see Supporting Information). An alanine insertion mutation (A2*) results in an expressed protein with N-terminal sequence NH₂–Ala₂*Ser₃His₄ after processing, instead of NH₂–Ser₂His₃, thereby disrupting the spacing between His₃ and NH₂ terminus. This mutation was constructed to test the role of amino-terminal coordination in metal

(49) Melnik, M.; Sramko, T.; Dunajjurco, M.; Sirota, A.; Jona, E.; Holloway, C. E. *Rev. Inorg. Chem.* **1995**, *15* (1–2), 139–44.

(50) Bennett, B. EPR of cobalt-substituted zinc enzymes. In *Metals in Biology: Applications of High Resolution EPR to Metalloenzymes*; Hanson, G. R., Berliner, L. J., Eds.; Springer: New York, 2007; p 29.

(51) Crawford, P. A.; Yang, K. W.; Sharma, N.; Bennett, B.; Crowder, M. W. *Biochemistry* **2005**, *44* (13), 5168–76.

responsiveness based on the precedent of N-terminal Ni-coordination seen with Ni-SOD.^{52,53}

Ni(II) Responsiveness. Two mutations (Ala2* and His3Leu) completely abolished Ni-responsive LacZ expression. A third mutation (His64Leu) showed only partial induction (44%) as compared to WT. Mutants at two positions (Cys35Ala and His60Ala) that lost Co(II) responsiveness maintained sensitivity to Ni(II), suggesting a key structural difference in the metal-mediated inhibition of DNA binding by Ni(II) versus Co(II). The response of Cys35Ala to Ni(II) was unexpected based on the XAS analysis for this protein, which showed the loss of S-coordination. A Cys35Leu mutation was nonresponsive to Ni(II), which could result from steric perturbation of the Ni(II)-binding site. Additional mutations of His60 to Leu and Asn also maintained Ni sensitivity, suggesting that this His residue is not a Ni ligand, in contrast to results for Co(II). Only the mutation of His60 to Arg completely abolished nickel responsiveness, most likely due to steric effects from the large, positively charged Arg side chain. Mutation of Asp7, Asp30, His33, Glu34, Glu63, His67, Asp77, Asp79, Asp86, or Tyr88 had no effect on Ni(II) responsiveness, eliminating these residues as possible Ni ligands.

Co(II) Responsiveness. Mutant RcnR constructs were tested for LacZ activity in the presence of 200 μM CoCl_2 . LacZ activity was increased ~ 20 -fold under these experimental conditions when WT RcnR is present (Figure 1). The Ala2*, His3Leu, Cys35Ala, His60Ala/Asn/Leu/Arg, and His64Leu mutants were nonresponsive ($<10\%$ WT) to added Co(II) and showed constitutive, low LacZ activity. These mutations potentially account for five of the six cobalt ligands determined by XAS. As with Ni(II), mutation of Asp7, Asp30, His33, Glu34, Glu63, His67, Asp77, Asp79, Asp86, or Tyr88 had little or no effect on Co(II)-dependent LacZ expression as compared to WT RcnR.

UV-vis Absorption Spectroscopy of RcnR Mutants. UV-vis spectroscopy of mutant RcnR proteins was used to explore the relationship between M(II) coordination and LacZ assay data, in particular, the differences between Cys35, His60, and His64 in Ni(II) and Co(II) coordination. Individual mutations were constructed in the RcnR expression vector. Mutant proteins were overexpressed and purified the same as for WT RcnR, with similar yields. UV-vis spectra for M(II)-substituted RcnRs (1:1 metal/RcnR) are shown in the Supporting Information (Figure S8). The extinction coefficients of the absorbance maxima for each M(II)-RcnR variant are listed in Table 2. The fully Ni(II)- and Co(II)-responsive D30N RcnR mutant (Figure 6) served as a control protein that exhibited little perturbation of the UV-vis spectra of the M(II)-substituted proteins.

Ni(II)-Substituted Mutants. WT Ni-RcnR exhibited minimal absorbance features between 400 and 700 nm, as expected of a six-coordinate metal site that lacks d-d electronic transitions. Decreases in the Ni(II)-coordination number are expected to increase absorbance in the visible range. An alteration in the interaction with Cys35 is expected to shift the energy or magnitude of the 280 nm $\text{S} \rightarrow \text{Ni}$ LMCT peak. All RcnR mutants, except H3L, bound ~ 1 equiv of Ni/RcnR monomer (Supporting Information Figure S9). H3L RcnR showed no features in the difference UV-vis spectrum indicative of bound Ni(II) atoms. For both C35A and C35L RcnR, the 280 nm $\text{S} \rightarrow \text{Ni}$ LMCT feature was absent, while the 235 nm feature

Table 2. UV-vis Spectral Features for M(II)-Substituted RcnRs

RcnR variant	Ni(II)		Co(II)	
	λ (nm)	ϵ ($\text{M}^{-1} \text{cm}^{-1}$)	λ (nm)	ϵ ($\text{M}^{-1} \text{cm}^{-1}$)
WT	235	3937	238	4675
			240	4609
			314	2664
A2*	286	1778	549	shoulder
	240	10555	239	12200
	428	136	286	shoulder
			322	4681
			557	shoulder
H3L	none		238	2733
			296	2270
			597	194
D30N			619	214
	234	4420	242	4100
	280	1709	312	2500
C35A	240	8308	238	4598
	456	76	272	1678
			325	850
C35L			491	86
	240	6052	239	3626
	452	48	346	shoulder
H60L	237	3462	238	9539
	281	2407	307	5134
	287	2429	450	shoulder
H60R			586	shoulder
	238	6840	240	14280
	290	shoulder	284	14145
H64L	450	45	550	shoulder
	237	3666	239	10148
	278	2546	289	6922
	380	40	550	shoulder
	600	19		

remained. Additionally, both of these mutants had a feature at ~ 450 nm, which can be attributed to a Ni(II) d-d transition. These data indicate that both mutants have four- or five-coordinate sites, the latter being consistent with XAS analysis of C35A Ni(II)-RcnR. The Ni(II)-A2*-RcnR spectrum contained a feature at 420 nm ($\epsilon_{420} = 136 \text{ M}^{-1} \text{cm}^{-1}$) that is of similar intensity to one observed for the square-planar Ni(II) site of *E. coli* NikR,⁵⁴ and a similar geometry may be adopted for A2* Ni(II)-RcnR. This mutant also lacked the 280 nm $\text{S} \rightarrow \text{M}$ LMCT feature, suggesting a loss of Cys coordination. The H64L Ni(II)-RcnR spectrum contained two weak features at 380 and 600 nm not seen for WT. These features are consistent with a six-coordinate Ni(II) site⁵⁵ that is clearly distinct from WT Ni(II)-RcnR. Except for A2* and H3L, each mutant shows a significant shift in the 280 nm $\text{S} \rightarrow \text{M}$ LMCT feature. Together, these data strongly indicate that the NH_2 group of the amino-terminus and the side chains of His3, Cys35, and His64 coordinate nickel. When these residues were disrupted, the proteins adopted altered coordination geometries (4, A2*; 5, C35A; and 6, H64L) or did not bind Ni at all (H3L). In contrast, the H60L Ni(II)-RcnR spectrum showed neither an increase in the intensity of d-d features nor a significant perturbation of the 280 nm peak, suggesting that this residue is not involved in Ni coordination. These data are consistent with the LacZ data for the H60L/N mutants. Only when His60 was substituted with the bulky, charged Arg side chain was the UV-vis Ni(II) spectrum affected, with features suggesting a five-coordinate metal site. The effect of H60R mutation on Ni(II) responsiveness indicates that this residue is near the metal site but does not contribute to Ni(II) coordination.

(52) Wuerges, J.; Lee, J. W.; Yim, Y. I.; Yim, H. S.; Kang, S. O.; Carugo, K. D. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101* (23), 8569–74.

(53) Barondeau, D. P.; Kassmann, C. J.; Bruns, C. K.; Tainer, J. A.; Getzoff, E. D. *Biochemistry* **2004**, *43* (25), 8038–47.

(54) Chivers, P. T.; Sauer, R. T. *J. Biol. Chem.* **2000**, *275* (26), 19735–41.

(55) Glennon, J. D.; Sarkar, B. *Biochem. J.* **1982**, *203* (1), 15–23.




Figure 6. Effect of side-chain substitution of conserved residues on RcnR function *in vivo*. (Top) Mutants that affect either Co(II) or Ni(II) responsiveness. (Bottom) Mutants that do not affect Ni(II) and Co(II) responsiveness. β -Galactosidase activity was measured for the P_{rcnA} -lacZ reporter construct encoding WT or mutant *rcnR* genes (Supporting Information Table S8) in *E. coli* strain PC563 grown anaerobically in LB media in the absence (black bars) or presence of 1 mM NiCl₂ (orange bars) or 200 μ M CoCl₂ (blue bars). Each mutant is normalized to WT activity with either NiCl₂ or CoCl₂. A relative activity of 1 corresponds to ~168 Miller units with Ni(II) and 209 Miller units for Co(II). All data are normalized to WT plus Ni(II) for ease of comparison. Error bars show the SD of the LacZ activity of three separate samples. Mutations were constructed in pJ114 as described in the Experimental Procedures.

Co(II)-Substituted Mutants. Unlike Ni(II)-RcnR UV-vis spectra, the d-d transitions of WT and mutant Co(II)-RcnR proteins were difficult to resolve due to the extended tail of the S \rightarrow Co LMCT (Figure 3; ~315 nm). However, other differences can be observed for Co(II)-substituted RcnR mutants (see Supporting Information). These changes, combined with alterations to the S \rightarrow Co LMCT feature, are diagnostic of changes in Co(II) coordination. As with WT RcnR, all mutant proteins tested bound ~1 equiv of Co(II) per RcnR monomer (see Supporting Information). The C35A and C35L Co(II)-RcnR spectra lacked a strong absorbance feature at 315 nm. These data confirm the assignment of this peak as S \rightarrow Co LMCT. They also provide further support for C35 as a Co ligand, in agreement with XAS analysis of C35A Co(II)-RcnR. H3L RcnR still bound Co(II), although the Co(II) affinity appeared to be significantly weaker than for WT based on the sigmoidal shape of the titration curve. This mutant also showed distinct UV-vis spectral features in the d-d transition region with peaks at 597 nm ($\epsilon = 194 \text{ M}^{-1} \text{ cm}^{-1}$) and 619 nm ($\epsilon = 214 \text{ M}^{-1}$

cm^{-1}). These values are most consistent with a five-coordinate Co(II) complex.⁵⁶

Spectra for A2*, H60L, and H64L Co(II)-RcnR were not as distinctly different from WT, but each showed a large increase in absorbance for the S \rightarrow Co transition and a shift in energy for this transition (A2*, longer wavelength and H60L and H64L, shorter wavelengths). Changes also were observed between 500 and 700 nm, but these differences are difficult to resolve due to overlap with other features. It is likely that the mutant sites remain six-coordinate but with a variable sixth ligand as seen by XAS for C35A Co(II)-RcnR. While precise characterization of these mutant sites cannot be made, it is clear that the mutations altered the Co(II) site and further support the model of Co(II) coordination by the NH₂-terminus, H3, C35, H60, and H64 suggested by the LacZ assay data.

Discussion

RcnR plays a pivotal role in intracellular sensing of Ni(II) and Co(II) ions in *E. coli*, where it must specifically interact with these metal ions when they are deemed to be in excess. The identification of the metal-binding ligands by a combination of XAS and mutagenesis experiments provides a starting point for understanding the allosteric regulation of RcnR activity by Ni(II) and Co(II). The characterization of the Ni(II)- and Co(II)-binding sites also permits a comparison of RcnR with NikR, the other nickel-responsive regulator in *E. coli*, as well as Ni(II)-/Co(II)-responsive regulators in other organisms. The results of this work, in combination with the prior study of Cu(I)-responsive CsoR,¹² established a framework for understanding the ligand selectivity of other members of the RcnR/CsoR family of transcription factors.

Ni(II) and Co(II) bind to RcnR with a 1:1 stoichiometry and upper limits on the dissociation constants of 25 and 5 nM, respectively. These affinities make RcnR sensitive to low levels of free nickel and cobalt ions (lower limit ~1 nM) in *E. coli* that, in the absence of other effects, could lead to relatively rapid induction of *rcnA* expression in response to excess metal. Both Ni(II) and Co(II) increase the stability of the RcnR tetramer without altering its oligomeric state or producing large-scale changes in secondary structure. The identity of the subunit interface required for RcnR tetramerization is unknown. The structure of *M. tuberculosis* CsoR revealed a dimeric species,¹² from which a reasonable structural homology model of RcnR can be constructed. The solution and crystal structure data presented for CsoR emphasized the dimeric form of the protein, but there was also evidence for a tetrameric species. In particular, two distinct dimer-dimer interfaces are present in the unit cell, either of which may be relevant to tetramer formation by RcnR or may simply result from packing in the crystal lattice. CsoR also has a 30-amino acid C-terminal extension not visible in the crystal structure that may influence the multimeric state of the protein. RcnR dimers may occur in the cell, where its concentration is likely an order of magnitude less than those used in the analytical ultracentrifugation experiments. Differences in the oligomeric state between the two proteins may impact the mode of DNA binding and recognition, which is currently unknown. Nonetheless, the metal-binding properties of both CsoR and RcnR can be fully explained by examining only the dimeric assembly of the protein monomers.

The XAS analyses for Co(II)- and Ni(II)-RcnR reveal similar, but not identical, six-coordinate binding sites for each metal. XANES analyses indicate a pseudo-octahedral geometry for both metal ions, and EXAFS analyses are consistent with a

(56) Outten, C.; Tobin, D.; Penner-Hahn, J.; O'Halloran, T. V. *Biochemistry* **2001**, *40*, 10417-23.

similar set of six ligands: Cys(His)₂(N/O)₃ (Figure 5 and Table 1). The major difference seen between Co(II) and Ni(II) centers is the M–S distance, which is long for Ni(II)–RcnR (2.54 Å) and short for Co(II)–RcnR (2.24 Å). The 2.54 Å Ni–S distance is higher than the mean distance for unidentate S-donor ligands of all types (2.47 Å)⁴⁹ and at the high end of the distances found in the small number of structurally characterized high-spin, six-coordinate Ni(II) thiolate complexes that are known. Six-coordinate Ni(II) complexes are invariably high-spin ($S = 1$), which accounts for longer nickel–ligand distances in these complexes. The lability of the high-spin $S = 1$ configuration also likely accounts for the fact that only a few high-spin nickel thiolates have been characterized, and these feature chelating thiolate ligands.^{41,42} For example, diaqua bisdimethylaminothioquadronickel(II) (NiS₂N₂O₅C₁₂H₂₀) contains an average Ni–S distance of 2.469(1) Å,⁴¹ tetraethylammonium tris-2-thiopyridiniumnickelate(II) (Ni(SC₅H₄N)₃) contains Ni–S distances ranging from 2.518(1) to 2.541(1) Å,⁴² and tetraphenylphosphonium tris-2-thiopyrrolidiniumnickelate(II) (Ni(SC₄H₃N₂)₃ · CH₃CN) contains Ni–S bond distances ranging from 2.460(2) to 2.545(2) Å.⁴² The Ni–S distance in Ni(II)–RcnR is clearly distinct from the short Ni–S distances found in planar, $S = 0$, NiS₄ complexes (mean Ni–S = 2.18 Å).⁴⁹ The Ni–N average distances observed for Ni(II)–RcnR (1.98 and 2.11 Å) also are consistent with the high-spin configuration and fall in the range observed for structurally characterized high-spin Ni(II)N₆ complexes (mean Ni–N = 2.07 Å).⁴⁹ Thus, the metal–ligand distances found for the WT Ni(II)–RcnR complex are consistent with a high-spin ($S = 1$) complex. Co(II)–RcnR has an EPR spectrum that is typical of a high-spin, $S = 3/2$ center,⁵¹ and clearly different from the low-spin, $S = 1/2$, complexes that have been characterized.⁵⁷ The metal–ligand distances found for six-coordinate Co(II) thiolates are variable [2.338(1)–2.585(1) Å],^{58,59} a reflection of the $S = 3/2$, d^7 electronic configuration, which lacks an unfilled π -symmetry orbital in contrast to d^8 Ni(II). An example of a structurally characterized six-coordinate Co(II) thiolate complex with a confirmed $S = 3/2$ ground state ($\mu_{\text{eff}} = 4.83 \mu_{\text{B}}$) is dimethanol bisdimethylaminothioquadratecobalt(II) (CoS₂N₂O₅C₁₂H₂₀), which features an average Co(II)–S distance of 2.500(1) Å.⁶⁰ The Co–S distance found for Co(II)–RcnR lies at the lower end of the range of Co(II)–S distances observed. The difference between Co(II)–S and Ni(II)–S distances in RcnR suggests that either the Co(II)–S bond is supported by a π -interaction that is absent for Ni(II) or that the protein conformation allows stronger Co(II)–S binding. A likely consequence of the shorter Co(II)–S interaction is a different protein conformation resulting from Co(II) versus Ni(II) binding.

The structural characterization of Ni(II)- and Co(II)-binding sites of RcnR allows a comparison with Ni(II) and Co(II) sites in other metalloregulators from *E. coli* and other bacteria. This study also completes structural characterization of the three known Ni(II)-binding sites involved in transcriptional regulation in *E. coli*. The two nickel-responsive transcriptional regulators from *E. coli*, RcnR and NikR, operate in a sequential fashion

in response to increasing intracellular nickel.⁹ NikR binds to DNA in the presence of Ni(II) and represses the transcription of the genes encoding the NikABCDE uptake permease. In contrast, RcnR-dependent repression of *rcnA* is maintained until *nikABCDE* expression is fully repressed, then increasing Ni(II) releases RcnR from DNA. NikR contains two Ni(II)-binding sites, termed high- and low-affinity. The high-affinity Ni(II)-binding site ($K_{\text{d}} = \sim 1 \text{ pM}$)^{7,35} is a low-spin ($S = 0$), four-coordinate planar complex with His₃Cys coordination^{61,62} that is essential for biological function.⁶³ Changes in high-affinity Ni(II) site occupancy are thought to tune *nikABCDE* expression according to nickel–enzyme expression levels.^{9,63} The low-affinity Ni sites ($K_{\text{d}} = 5 \text{ nM}$)⁶⁴ are six-coordinate but do not employ thiolate coordination.⁸ Low-affinity site occupancy increases DNA-binding affinity ($K_{\text{d}} = \sim 5 \text{ pM}$)^{35,64} and results in the full repression of *nikABCDE* expression.⁶³ The Ni(II) site in RcnR is high-spin and six-coordinate with an affinity similar to the six-coordinate site of NikR, but their relative numbers in the cell are probably quite different—the NikR low-affinity sites occur only when bound to DNA, so there are two per cell (considering that there is only one known NikR DNA-binding site in the chromosome), while the number of RcnR sites is likely 1–2 orders of magnitude greater based on the range of intracellular concentrations observed for different repressor proteins in *E. coli*.^{35,65–67}

The differences in the electronic structures of the three sites have several ramifications for the physical and functional properties of the Ni(II) sites. First, the metal–ligand bond distances are quite different, seen most dramatically in the Ni–SCys distances for NikR⁶¹ (2.13 Å) and RcnR (2.54 Å). These differences reflect the increased ionic radius of the high-spin ion that occurs when σ^* levels are populated. The mechanism for ligand exchange reactions for the two types of Ni(II) centers are also expected to differ. High-spin Ni(II) centers typically undergo ligand exchange via ligand dissociation, which is favored by the high-spin electronic configuration. In contrast, four-coordinate planar complexes undergo ligand exchange via a mechanism involving association of the incoming ligand and therefore depend strongly on the nature of the nucleophile involved.⁶⁸ Thus, the NikR high-affinity site, which is not easily accessible in the protein, is not prone to easy nickel release once the metal is bound. A specific mechanism, such as a conformational change, might be required to release nickel. The occupancy of the high-affinity site of NikR is inversely correlated with nickel–enzyme expression,^{9,63} and its function is consistent with the conservation of biosynthetic resources rather than stress. In contrast, the six-coordinate Ni(II) sites of NikR and RcnR appear to be designed for spontaneous metal release in response to decreasing metal ion concentration, as both of these sites function to respond to fluctuations in

(57) Brennan, B. A.; Alms, G.; Nelson, M. J.; Durney, L. T.; Scarrow, R. C. *J. Am. Chem. Soc.* **1996**, *118* (38), 9194–5.

(58) Fun, H.-K.; Raj, S. S.; Zhu, X.-H.; Chen, X.-F.; Zuo, J.-L.; You, X.-Z. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1999**, *55*, 896–9.

(59) Yu, P.; Zhuang, B.; Huang, L.; Pan, B. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1996**, *52*, 630–2.

(60) Hehl, U.; Hinse, P.; Frolich, R.; Mattes, R. *Z. Anorg. Allg. Chem.* **2002**, *628*, 770–8.

(61) Carrington, P. E.; Chivers, P. T.; Al-Mjeni, F.; Sauer, R. T.; Maroney, M. J. *Nat. Struct. Biol.* **2003**, *10* (2), 126–30.

(62) Schreiter, E. R.; Sintchak, M. D.; Guo, Y.; Chivers, P. T.; Sauer, R. T.; Drennan, C. L. *Nat. Struct. Biol.* **2003**, *10* (10), 794–9.

(63) Rowe, J. L.; Starnes, G. L.; Chivers, P. T. *J. Bacteriol.* **2005**, *187* (18), 6317–23.

(64) Bloom, S. L.; Zamble, D. B. *Biochemistry* **2004**, *43* (31), 10029–38.

(65) Gilbert, W.; Muller-Hill, B. *Proc. Natl. Acad. Sci. U.S.A.* **1966**, *56* (6), 1891–8.

(66) Gunsalus, R. P.; Miguel, A. G.; Gunsalus, G. L. *J. Bacteriol.* **1986**, *167* (1), 272–8.

(67) Kolodrubetz, D.; Schleif, R. *J. Mol. Biol.* **1981**, *149* (1), 133–9.

(68) Wilkins, R. G. *Kinetics and Mechanism of Reactions of Transition Metal Complexes*; VCH Publishers, Inc.: New York, 1991.

intracellular nickel concentrations that occur under stress conditions.^{9,63}

The use of a single cysteine thiolate ligation in NikR and RcnR would seem to be an ideal choice for a metal ion electronic structure sensor. The thiolate sulfur is both a σ - and a π -donor and is therefore able to discriminate between metal ions by sensing the occupancy of both sets of metal d-orbitals. Considering the Co(II)- and Ni(II)-substituted forms of RcnR and NikR, low-spin Ni(II) has filled π -symmetry d-orbitals and the highest energy σ -orbital ($d_{x^2-y^2}$) is directed at the ligands and is unoccupied. This electronic structure leads to a very short in-plane Ni–S distance (2.13 Å). In contrast, for high-spin Ni(II), there is still no vacancy in the π -symmetry orbitals, but the σ -symmetry orbitals are both singly occupied, resulting in a much longer Ni(II)–S_{Cys} distance. For Co(II), the π -symmetry orbitals have one vacancy, opening up a possible π -interaction, resulting in an intermediate M–S distance (2.26 Å). Thus, the use of a single cysteine ligand in metallosensors is a useful mechanism for interrogating the electronic structure of the bound metal ion, a critical aspect of generating the appropriate metal-specific allosteric response.

Mutation of RcnR identified specific residues involved in metal binding and support the ligand set determined from EXAFS analysis. The mutant data reveal that Ni(II) and Co(II) use distinct, but overlapping, ligand spheres to regulate RcnR DNA-binding activity. The mutation of His3 showed the most severe effects on metal binding, virtually eliminating UV–vis spectral features for Ni(II) and substantially decreasing Co(II)-binding affinity. The involvement of the N-terminus in the metal-binding site is supported by the CsoR structure, where the amino terminus is near the Cu(I)-binding site (Figure 7).¹² The extension of the N-terminus by one amino acid (A2* mutant) also impaired Ni(II) and Co(II) binding and responsiveness. The involvement of the N-terminal amino group and the side chain of the second residues is similar to that observed for Ni–SOD.^{52,53} In this case, the deprotonated main chain amide of residue 2 also coordinates the Ni atom, suggesting that RcnR also uses main chain coordination by the amide of His3 for Ni(II) and Co(II) binding. Both Ni(II) and Co(II) are known to bind Gly–His and deprotonate the main chain amide to form a stable coordination complex,^{69,70} providing a chemical precedent for our model. The A2* mutation likely impairs Ni(II) responsiveness by forming a square-planar ATCUN motif (Xaa–Xaa–His).^{70,71} The tripartite coordination of Ni(II) and Co(II) by the two N-terminal residues of RcnR is a key aspect of its selectivity toward metals with preferences for higher coordination numbers.

His64 also contributes to the responsiveness of Co(II) and Ni(II). The slightly different effect of the His64Leu mutation on the metal response for Ni(II) and Co(II) is consistent with expectations arising from differences in electronic structures of metal coordination complexes. This residue at this position in CsoR is Cys, suggesting that changes in this residue position will be important for metal selectivity and responsiveness regardless of coordination number.

The differential contributions of Cys35 and His60 to Co(II) and Ni(II) responsiveness suggest that these residues play different roles in the structural response of RcnR to metal binding. The spectroscopic studies of RcnR demonstrate that Cys35 is a ligand for both metals. The basis for the absence of




Figure 7. Fingerprint of the RcnR/CsoR family. (A) Image of Cu(I)–CsoR (PDB ID 2HH7) showing the residues (RcnR numbering) that coordinate the cuprous ion (orange sphere). The two different subunits of the CsoR dimer are shown in light blue and orange. Text in parentheses refers to the identity of RcnR Ni/Co coordinating residues. (B) Scheme of CsoR primary structure showing the positions of the fingerprint amino acids identified by bioinformatics analysis. The orange and blue bars correspond to the regions of structure shown in panel A. (C) Summary of the sequence fingerprint, gene neighborhood, and ligand predictions of different classes within the CsoR/RcnR family. See Supporting Information for more details.

an effect of the Cys35Ala mutation on Ni(II) responsiveness is currently unclear. Cys35 may play a more prominent role in metal selectivity rather than an allosteric response to metal binding. In contrast to Cys35, the mutation of His60 has little effect on the spectroscopic properties of the Ni(II) site. This position is invariant in RcnR and CsoR proteins, and so its dispensability with regard to Ni(II) response is unexpected but might reflect another difference between the ligands used to bind Ni(II) and Co(II).

The minimal structure that emerges from the combination of XAS and mutagenesis analysis, M(Cys)(His)₂(NH₂–Ser)(N/O)₂, clearly identifies five metal–ligands (the NH₂-terminus, the His3 amide, and the side chains of His3, Cys35, and His64) for both Ni(II) and Co(II), and a sixth, His60, for Co(II). The difference between EXAFS and mutagenesis, combined with structural homology, in assigning the number of His ligands is not unreasonable owing to the effect of His conformation on fitting these residues to EXAFS data.⁷² A sequence alignment of the predicted RcnR proteins (see Supporting Information) does not reveal a highly conserved sixth residue with a side chain likely to function in metal binding (Asp, Glu, Met, His, and Cys). Mutation of several candidate residues, including those closest to the M(II)-binding site, did not cause any changes in Co(II) or Ni(II) responsiveness, leaving the identity of the sixth Ni(II) unresolved. The mutagenesis approach will only identify side chain ligands, and so a Ni(II) coordination sphere completed by a second main chain amide will require additional experi-

(69) Farkas, E.; Sovago, I.; Gergely, A. *J. Chem. Soc., Dalton Trans.* **1983**, 1545–51.

(70) Sovago, I.; Osz, K. *Dalton Trans.* **2006**, (32), 3841–54.

(71) Harford, C.; Sarkar, B. *Acc. Chem. Res.* **1997**, *30* (3), 123–30.

(72) Ferreira, G. C.; Franco, R.; Mangravita, A.; George, G. N. *Biochemistry* **2002**, *41* (15), 4809–18.

mental approaches for identification. Alternately, the coordination sphere could be completed by a ligand from the solvent (aqua or hydroxo) but not a halide ion, based on XAS data.

The regulation of DNA binding through allosteric changes communicated between metal- and DNA-binding sites is commonly observed in metalloregulator proteins.¹ Giedroc and co-workers showed for Zn-responsive CzrA and SmtB proteins that two residues of the tetrahedral metal-binding site are critical for transmitting the metal-binding state of the site to the rest of the protein.^{73,74} The other two residues are primarily important for maintaining the tetrahedral coordination geometry. These observations suggest that within a particular structural class of metalloregulator, one or more master residues are critical for allostery. RcnR and CsoR share three conserved ligand positions (35, 60, and 64), suggesting that one or more of these residues may be essential for the allosteric response to metal binding. The regulation of RcnR by Co(II) and Ni(II) binding is coupled with a change in stability. For Co(II) and Ni(II), the ligand-binding sites utilize three distinct regions of the protein primary sequence and link these elements together (Figure 7). The stability change could include a rearrangement of secondary structure elements relative to the metal-free protein or altering the orientation of monomers/domains relative to one another as seen for the iron-responsive regulator DtxR.⁷⁵ Either of these mechanisms is consistent with the absence of a change in RcnR secondary structure upon metal binding. The unaltered sensitivity of the Cys35Ala and His60 mutants to Ni(II) suggests either that these residues are not important for the allosteric response or that more than one allosteric network exists in this protein family. The Cys35 mutant data suggest that Ni(II)-dependent allostery originates from intramonomer metal coordination, in contrast to Co(II)-RcnR and Cu(I)-CsoR, where this residue is essential for activity. There is no functional requirement to preserve a single allosteric mechanism for the metal-dependent inhibition of DNA binding for a particular metalloregulator. The primary requirement of this class of metalloregulator is that metal binding reduces DNA-binding affinity. Thus, Ni(II)- and Co(II)-RcnR are likely distinct structural entities whose common feature is an inability to bind DNA with appreciable affinity.

The requirement for thiolate coordination in Co(II) discrimination by RcnR is currently a singular occurrence. RcnR is functionally analogous to *M. tuberculosis* NmtR and KmtR, which also utilize six-coordinate geometry to bind Ni(II) and Co(II) between two monomer chains.^{6,76,77} However, KmtR and NmtR use only N and O ligands for metal coordination. Neither protein appears to utilize NH₂-terminal, main-chain amide, and thiolate coordination environment proposed here for RcnR. Additionally, mutational studies of putative metal-binding ligands in both proteins have shown identical effects, or lack thereof, on Ni and Co responsiveness.⁷⁷ KmtR only recently was identified,⁷⁷ so the physiological requirement for two functionally similar Ni-/Co-responsive regulators in the same

organism is not fully appreciated. However, the mutagenesis studies of NmtR and KmtR demonstrate that, despite their structural homology, the structural location of the metal-binding sites in each protein is distinct. This difference also manifests itself in the metal-binding affinities of the two proteins, wherein KmtR has a higher relative affinity for Ni(II) and Co(II) than NmtR.⁷⁷

NmtR and KmtR, like RcnR, are members of a large family of metalloregulators (ArsR/SmtB) that show distinct metal-binding specificities that correlate with differences in coordination number. Interestingly, the Zn-/Co-responsive *Staphylococcus aureus*⁶ CzrA protein uses a four-coordinate tetrahedral site. The higher coordination number observed in NmtR is thought to be achieved by the recruitment of two His ligands from a short C-terminal extension in the protein that is not present in CzrA,⁷⁶ similar to the recruitment of N-terminal residues for Ni/Co coordination by RcnR. The MerR family uses a variable loop region between two helices to increase the coordination number and alter metal specificity.⁷⁸ These observations suggest that increases in metal site coordination number within a structural family generally require recruitment of ligands from distinct regions of the protein that are more tolerant to amino acid insertions and deletions, such as loops and the amino and carboxy termini.

The RcnR/CsoR family contains several distinct functional classes that likely respond to very different types of allosteric effectors. Ensuring ligand-binding specificity would require adaptation of the general binding site that has been identified from these studies of RcnR and those of CsoR. CsoR coordinates Cu(I) with three ligands (Cys35, His60, and Cys64, using RcnR numbering),^{12,79} while RcnR responds biologically to metals that require higher coordination spheres by recruiting ligands from the N-terminal region, which is not used by CsoR for metal coordination, as well as a residue changes at position 64. The identification of four RcnR residues that when mutated drastically affect metal binding and metal responsiveness suggests that RcnR-/CsoR-like sequences can be analyzed to make predictions about ligand-binding specificities. For example, sequences containing His/Cys/His/His residues at positions 3, 35, 60, and 64 (RcnR numbering), respectively, are likely Ni(II)- and Co(II)-responsive regulators (Figure 7). Proteins with Xaa/Cys/His/Cys residues (Xaa = nonconserved residue) are likely Cu(I)-responsive regulators. Thus, the residues at these four positions provide a fingerprint that likely correlates with the ligand-binding specificity of a particular protein class in the largely uncharacterized RcnR/CsoR superfamily.

Using these minimal sequence criteria, ~450 unique sequences of the RcnR/CsoR family (COG 1937) were categorized into six different groups (Figure 7). CsoR-like regulators (Xaa/Cys/His/Cys) are the most abundant class, with 242 sequences. Of the remaining sequences, 72 (47 sequences with <95% identity; see Supporting Information) are predicted to be Ni-/Co-responsive transcriptional regulators. All sequences with this pattern are expected to express a protein with only one residue preceding His3,³⁸ suggesting a strong bias toward the N-terminal coordination motif proposed here. All of the proposed *rcnR* genes have a neighboring gene that encodes a putative metal efflux protein (Figure 7), likely responsible for Ni/Co export. Position 3 in other groups varies widely and may not play a

(73) Eicken, C.; Pennella, M. A.; Chen, X.; Koshlap, K. M.; VanZile, M. L.; Sacchetti, J. C.; Giedroc, D. P. *J. Mol. Biol.* **2003**, *333* (4), 683–95.

(74) Pennella, M. A.; Arunkumar, A. I.; Giedroc, D. P. *J. Mol. Biol.* **2006**, *356* (5), 1124–36.

(75) Pohl, E.; Holmes, R. K.; Hol, W. G. *J. Biol. Chem.* **1998**, *273* (35), 22420–7.

(76) Cavet, J. S.; Meng, W.; Pennella, M. A.; Appelhoff, R. J.; Giedroc, D. P.; Robinson, N. J. *J. Biol. Chem.* **2002**, *277* (41), 38441–8.

(77) Campbell, D. R.; Chapman, K. E.; Waldron, K. J.; Tottey, S.; Kendall, S.; Cavallaro, G.; Andreini, C.; Hinds, J.; Stoker, N. G.; Robinson, N. J.; Cavet, J. S. *J. Biol. Chem.* **2007**, *282* (44), 32298–310.

(78) Changela, A.; Chen, K.; Xue, Y.; Holschen, J.; Outten, C.; O'Halloran, T. V.; Mondragon, A. *Science* **2003**, *301*, 1383–7.

(79) Smaldone, G. T.; Helmann, J. D. *Microbiology* **2007**, *153* (12), 4123–8.

role in ligand binding in these other proteins. Thus, the RcnR class may be unique because it has recruited residues from the N-terminal region to aid in metal recognition.

Formaldehyde-responsive FrmRs⁸⁰ (12 sequences) can be singled out based on their fingerprint (Xaa/Cys/His/Xaa fingerprint) and proximity to formaldehyde dehydrogenase genes. The structural basis for formaldehyde responsiveness by FrmR is not known, and the involvement of a metal ligand has not been explored. Additional sequence fingerprints for uncharacterized proteins also were detected, and predictions can be made about ligand specificities based on gene neighborhood analysis. For example, 92 genes encoding the Xaa/Cys/(Tyr, Asn, or Gly)/Cys proteins are preferentially near genes with functions related to antibiotic or oxidative stress responses, suggesting that not all RcnR/CsoR regulators are metal responsive. Some organisms, such as *Deinococcus geothermalis*,⁸¹ have multiple RcnR/CsoR family members, emphasizing the need for precise structural control of ligand-binding specificity. Further biochemical and biological studies will be required to identify the signals for each type of regulator. Once these signals are identified, the structural basis for ligand recognition and allosteric regulation can be characterized in detail.

(80) Herring, C. D.; Blattner, F. R. *J. Bacteriol.* **2004**, *186* (20), 6714–20.

(81) Makarova, K. S.; Omelchenko, M. V.; Gaidamakova, E. K.; Matrosova, V. Y.; Vasilenko, A.; Zhai, M.; Lapidus, A.; Copeland, A.; Kim, E.; Land, M.; Mavrommatis, K.; Pitluck, S.; Richardson, P. M.; Detter, C.; Brettin, T.; Saunders, E.; Lai, B.; Ravel, B.; Kemner, K. M.; Wolf, Y. I.; Sorokin, A.; Gerasimova, A. V.; Gelfand, M. S.; Fredrickson, J. K.; Koonin, E. V.; Daly, M. J. *PLoS ONE* **2007**, *2* (9), e955.

Acknowledgment. This work was supported by NSF Grant MCB0520877 to P.T.C., NIH Grant GM-69696 to M.J.M., and NIH National Research Service Award T32 GM08515 (S.L.). We thank the Chivers lab for helpful discussions, Haoran Li for technical assistance with LacZ assays, Johnna Roose for her AAS expertise, Mike Bradley for help with sequence analysis, and Profs. Brian Bennett and Paul Lahti for help with simulation of the Co(II)–RcnR spectrum and helpful discussions. XAS data collection was carried out at the Stanford Synchrotron Radiation Laboratory (SSRL), a national user facility operated by Stanford University on behalf of the U.S. Department of Energy, Office of Basic Energy Sciences. The SSRL Structural and Molecular Biology Program is supported by the Department of Energy, Office of Biological and Environmental Research and by the National Institutes of Health, National Center for Research Resources, Biomedical Technology Program.

Supporting Information Available: Concentration dependence of RcnR stability, plots of analytical ultracentrifugation data, UV–vis spectra IAM-modified RcnR and RcnR mutants, metal titrations of RcnR mutants, Fourier-filtered and Fourier transform spectra for metal-substituted RcnR, EPR spectrum of Co(II)-substituted RcnR, sequence alignment of RcnR family, sequence alignment of representative RcnR/CsoR family groups, tables of EXAFS fits for M(II) complexes, and tables of strains, plasmids, and oligonucleotides. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA710067D