



Addition/Correction

Isotopically Discriminated NMR Spectroscopy: A Tool for Investigating Complex Protein Interactions in Vitro

Alexander P. Golovanov, Richard T. Blankley, Johanna M. Avis, and Wolfgang Bermel J. Am. Chem. Soc., 2008, 130 (27), 8870-8870 • DOI: 10.1021/ja8036004 • Publication Date (Web): 05 June 2008 Downloaded from http://pubs.acs.org on February 8, 2009

More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

View the Full Text HTML





Isotopically Discriminated NMR Spectroscopy: A Tool for Investigating Complex Protein Interactions in Vitro [*J. Am. Chem. Soc.* 2007, *129*, 6528–6535]. Alexander P. Golovanov,* Richard T. Blankley, Johanna M. Avis, and Wolfgang Bermel

Supporting Information. In Figure S1a,b, the delay shown on the pulse sequence diagrams was labeled incorrectly as "k $t_1/4$ ". The labels should read " $(1-k)t_1/4$ ".

Supporting Information Available: Complete refs 3 and 4, detailed description of IDIS-HSQC and -TROSY pulse sequences with diagrams (corrected), illustration of the effect of incomplete incorporation of ¹³C label in doubly labeled protein on IDIS spectra, and reference spectra of unbound [¹⁵N,¹³C]N3 and [¹⁵N]- WW3-4. This material is available free of charge via the Internet at http://pubs.acs.org.This material is available free of charge via the Internet at http://pubs.acs.org.

JA8036004

10.1021/ja8036004 Published on Web 06/05/2008