

Additions and Corrections

**Contact Model for the Prediction of NMR
N#H Order Parameters in Globular Proteins**

Fengli Zhang, and Rafael Bru#schweiler

J. Am. Chem. Soc., **2008**, 130 (28), 9178-9178 • DOI: 10.1021/ja8020445 • Publication Date (Web): 24 June 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009

More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Contact Model for the Prediction of NMR N–H Order Parameters in Globular Proteins [*J. Am. Chem. Soc.* **2002**, *124*, 12654–12655]. Fengli Zhang and Rafael Brüschweiler*2406

Page 12654. In eq 1, it was inadvertently omitted that the interatomic distances $r_{i-1,k}^O$ and $r_{i,k}^H$ are defined by the corresponding internuclear distances *shortened by 1.2 Å*.

Supporting Information Available: Additional information about eq 1. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA8020445

10.1021/ja8020445

Published on Web 06/24/2008