


Communication

A Targetable Fluorescent Probe for Imaging Hydrogen Peroxide in the Mitochondria of Living Cells

Bryan C. Dickinson, and Christopher J. Chang

J. Am. Chem. Soc., **2008**, 130 (30), 9638-9639 • DOI: 10.1021/ja802355u • Publication Date (Web): 08 July 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

A Targetable Fluorescent Probe for Imaging Hydrogen Peroxide in the Mitochondria of Living Cells

Bryan C. Dickinson and Christopher J. Chang*

Department of Chemistry, University of California, Berkeley, California 94720

Received April 2, 2008; E-mail: chrischang@berkeley.edu


Hydrogen peroxide (H_2O_2) is an increasingly recognized small-molecule mediator of physiology, aging, and disease in living organisms.^{1–6} In this regard, aberrant production or accumulation of H_2O_2 within cellular mitochondria over time due to environmental stress(es) and/or genetic mutation(s) is connected to serious diseases where age is a risk factor, including cancer⁷ and neurodegenerative Alzheimer's, Parkinson's, and Huntington's diseases.^{8,9} Indeed, overexpression and mitochondrial targeting of catalase, a peroxide-detoxifying enzyme, can increase life span in mouse models.¹⁰ On the other hand, newer data suggest that controlled bursts of mitochondrial H_2O_2 can also serve beneficial roles for cell survival, growth, differentiation, and maintenance.^{3–6}

New imaging methods that allow visualization of localized production and accumulation of mitochondrial H_2O_2 in living samples are potentially useful for disentangling the complex contributions of this reactive oxygen species (ROS) to both healthy and diseased states. Synthetic fluorescent H_2O_2 indicators that can be targeted to precise subcellular locations offer one approach to this goal and do not require transfection like their protein counterparts,^{11,12} but traditional ROS indicators such as dihydro-rhodamine (DHR) are uncharged and hence not preferentially localized in cells before oxidation.¹³ In addition, DHR and related dyes are not specific for H_2O_2 over other ROS. Accordingly, mitochondrial-targeted small molecules for detection of specific ROS remain rare,^{13,14} and none of the probes reported to date are selective for H_2O_2 . We now report the synthesis and applications of mitochondria peroxy yellow 1 (MitoPY1), a new type of fluorophore for imaging mitochondrial H_2O_2 in living cells with ROS and spatial specificity.

Our overall strategy for fluorescence imaging of mitochondrial H_2O_2 in living systems is to create bifunctional dyes that contain both a peroxide-responsive element and a mitochondrial-targeting moiety. For the latter purpose, we were inspired by the use of phosphonium head groups by Murphy and others to deliver antioxidants, electrophiles, and EPR and optical probes to mitochondria, as these and related lipophilic cations selectively accumulate in this organelle due to proton gradient considerations.^{14–16} In addition, we sought a modular synthetic route that would allow facile introduction of a phosphonium or any other desired targeting group after installation of the boronate switch, which circumvents potential complications arising from sensitive functionalities that are incompatible with palladium-catalyzed Miyaura–Suzuki reactions typically used to introduce the H_2O_2 -cleavable boronate cage. Both of these design criteria can be met by the approach outlined in Scheme 1 for the synthesis of MitoPY1. The ability to append additional groups postboronation offers a host of opportunities for generating new multifunctional H_2O_2 imaging probes.

MitoPY1 features two major visible region absorptions ($\lambda_{\text{abs}} = 489 \text{ nm}$, $\epsilon = 14\,300 \text{ M}^{-1} \text{ cm}^{-1}$; 510 nm , $\epsilon = 14\,200 \text{ M}^{-1} \text{ cm}^{-1}$) and a weak emission ($\lambda_{\text{em}} = 540 \text{ nm}$, $\Phi = 0.019$, Figure S1) in 20 mM HEPES, pH 7. Reaction of MitoPY1 with H_2O_2 triggers a

Scheme 1. Synthesis and Activation of MitoPY1



fluorescence increase by its conversion to MitoPY1ox, which possesses one major absorption band at 510 nm ($\epsilon = 22\,300 \text{ M}^{-1} \text{ cm}^{-1}$) and enhanced emission ($\lambda_{\text{em}} = 528 \text{ nm}$, $\Phi = 0.405$). Kinetics measurements of the H_2O_2 -mediated boronate deprotection were performed under pseudo-first-order conditions ($5 \mu\text{M}$ dye, 10 mM H_2O_2), giving an observed rate constant of $k = 2.0(1) \times 10^{-3} \text{ s}^{-1}$. Figure 1 shows the relative turn-on fluorescence responses of MitoPY1 to a panel of biologically relevant ROS. Owing to its chemospecific boronate switch,^{17,18} the probe is selective for H_2O_2 over ROS like superoxide, nitric oxide, and hydroxyl radical.

MitoPY1 was then tested for its ability to both target the mitochondria and respond to H_2O_2 in living biological systems. Cervical cancer HeLa cells loaded with $5 \mu\text{M}$ MitoPY1 for 1 h at $37 \text{ }^\circ\text{C}$ show faint but measurable levels of fluorescence in discrete




Figure 1. Fluorescence responses of $5 \mu\text{M}$ MitoPY1 to various reactive oxygen species (ROS). Bars represent relative responses at 0, 5, 15, 30, 45, and 60 min after addition of each ROS. Data shown are for 10 mM $\text{O}_2\cdot^-$, $200 \mu\text{M}$ NO, and $100 \mu\text{M}$ for all other ROS. Data were acquired at $25 \text{ }^\circ\text{C}$ in 20 mM HEPES, pH 7, with excitation $\lambda = 503 \text{ nm}$ and emission collected between 510 and 750 nm .




Figure 2. Confocal fluorescence images of live HeLa cells with increases in mitochondrial H_2O_2 levels visualized using MitoPY1. Images displayed represent emission intensities collected in optical windows between 527 and 601 nm upon excitation at 510 nm for MitoPY1. HeLa cells incubated with $5 \mu\text{M}$ MitoPY1 for 60 min at 37°C and imaged with MitoPY1 (a), MitoTracker Red and Hoechst (overlay, b), and MitoPY1 with MitoTracker Red (overlay, c). HeLa cells incubated with $5 \mu\text{M}$ MitoPY1 for 60 min at 37°C with $100 \mu\text{M}$ H_2O_2 added for the final 40 min and imaged with MitoPY1 (d), MitoTracker Red and Hoechst (overlay, e), MitoPY1 and MitoTracker Red (overlay, f), and brightfield (g) with $20 \mu\text{m}$ scale bar. HeLa cells incubated with $5 \mu\text{M}$ MitoPY1 for 60 min at 37°C and imaged with MitoPY1 (h), MitoTracker Red and Hoechst (overlay, i), and MitoPY1 with MitoTracker Red (overlay, j). HeLa cells incubated for 24 h with 1 mM paraquat, then washed and incubated with $5 \mu\text{M}$ MitoPY1 for 60 min at 37°C and imaged with MitoPY1 (k), MitoTracker Red and Hoechst (overlay, l), MitoPY1 and MitoTracker Red (overlay, m), and brightfield (n) with $20 \mu\text{m}$ scale bar.

subcellular locations as determined by confocal microscopy (Figure 2a). Co-staining experiments with 50 nM MitoTracker Deep Red, a commercially available mitochondrial indicator (Figure 2b,c), or 500 nM LysoTracker Red, a lysosomal indicator (Figures S4–S7), establish that the observed fluorescence from MitoPY1 is localized to the mitochondria of these live cells. Addition of $100 \mu\text{M}$ H_2O_2 to HeLa cells loaded with MitoPY1 display a marked localized increase in fluorescence compared to control cells (Figure 2d–f). Again, co-staining with MitoTracker confirms that the dye is retained in the mitochondria and detects localized rises in H_2O_2 concentrations. Brightfield measurements and nuclear staining with Hoechst 33342 indicate that the cells are viable throughout the imaging experiments (Figures 2b,e,g). In addition, control experiments using a probe lacking the phosphonium targeting moiety (ContPY1, Figures S9–S12) or the oxidized probe (MitoPY1ox, Figures S13–S18) confirm that only MitoPY1 targets the mitochondria, and complementary flow cytometry experiments (Figure S8) provide supporting data over a larger population of cells. Finally, analogous experiments in Cos-7, HEK293, and CHO.K1 cell lines give similar results and expand the scope of the probe (Figures S5–S7). Taken together, these data establish that MitoPY1 is targeted to cellular mitochondria, where it can respond to localized changes in H_2O_2 levels in living samples.

Finally, we sought to utilize MitoPY1 to visualize endogenous production of H_2O_2 in the mitochondria of living cells. To this end, we treated HeLa cells with paraquat, a small-molecule inducer of oxidative stress that produces Parkinson's-like phenotypes.¹⁹ The images in Figure 2h–n show clear increases in mitochondrial-localized H_2O_2 levels detected with MitoPY1 within cells that had been exposed to 1 mM paraquat compared to control cells (IC_{50} of paraquat in HeLa cells is 1.02 mM).²⁰ These data indicate that MitoPY1 is sensitive enough to detect local mitochondrial H_2O_2 elevations associated with oxidative stress in this Parkinson's model.

To close, we have presented the synthesis, properties, and biological applications of MitoPY1, a new targeted fluorescent probe that can selectively detect H_2O_2 in the mitochondria of living cells. Our data show that MitoPY1 is capable of imaging changes in the levels of H_2O_2 within the mitochondria of a variety of mammalian cell lines, as well as H_2O_2 elevations caused by an oxidative stress model of Parkinson's disease. In addition to applying MitoPY1 and related chemical tools for studies of mitochondrial redox biology, we anticipate that this modular probe scaffold should prove useful

for creating new multifunctional probes for targeting, activation, and detection in living systems and are actively pursuing these possibilities.

Acknowledgment. We thank the Beckman, Packard, and Sloan Foundations, and the NIH (GM 79465) for providing funding for this work. B.C.D. thanks the NIH Chemical Biology Graduate Program (T32 GM066698) for support. We thank Holly Aaron (UCB Molecular Imaging Center) and Ann Fischer (UCB Tissue Culture Facility) for expert technical assistance.

Supporting Information Available: Synthetic and experimental details (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Rhee, S. G. *Science* **2006**, *312*, 1882–1883.
- (2) Stone, J. R.; Yang, S. *Antioxid. Redox Signal.* **2006**, *8*, 243–270.
- (3) Veal, E. A.; Day, A. M.; Morgan, B. A. *Mol. Cell* **2007**, *26*, 1–14.
- (4) D'Autr aux, B.; Toledano, M. B. *Nat. Rev. Mol. Cell Biol.* **2007**, *8*, 813–824.
- (5) Giorgio, M.; Trinei, M.; Migliaccio, E.; Pelicci, P. G. *Nat. Rev. Mol. Cell Biol.* **2007**, *8*, 722–728.
- (6) Poole, L. B.; Nelson, K. J. *Curr. Opin. Chem. Biol.* **2008**, *12*, 18–24.
- (7) Finkel, T.; Serrano, M.; Blasco, M. A. *Nature* **2007**, *448*, 767–774.
- (8) Barnham, K. J.; Masters, C. L.; Bush, A. I. *Nat. Rev. Drug Discovery* **2004**, *3*, 205–214.
- (9) Lin, M. T.; Beal, M. F. *Nature* **2006**, *443*, 787–795.
- (10) Schriener, S. E.; Jinfond, N. J.; Martin, G. M.; Treuting, P.; Ogburn, C. E.; Emond, M.; Coskun, P. E.; Ladiges, W.; Wolf, N.; Van Remmen, H.; Wallace, D. C.; Rabinovitch, P. S. *Science* **2005**, *308*, 1909–1911.
- (11) Belousov, V. V.; Fradkov, A. F.; Lukyanov, K. A.; Staroverov, D. B.; Shakhbazov, K. S.; Tersikh, A. V.; Lukyanov, S. *Nat. Methods* **2006**, *3*, 281–286.
- (12) Hanson, G. T.; Aggeler, R.; Oglesbee, D.; Cannon, M.; Capaldi, R. A.; Tsien, R. Y.; Remington, S. *J. Biol. Chem.* **2004**, *279*, 13044–13053.
- (13) Koide, Y.; Urano, Y.; Kenmoku, S.; Kojima, H.; Nagano, T. *J. Am. Chem. Soc.* **2007**, *129*, 10324–10325.
- (14) Robinson, K. M.; Janes, M. S.; Pehar, M.; Monette, J. S.; Ross, M. F.; Hagen, T. M.; Murphy, M. P.; Beckman, J. S. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 15038–15043.
- (15) Murphy, M. P.; Smith, R. A. *Annu. Rev. Pharmacol. Toxicol.* **2007**, *47*, 629–656.
- (16) Hardy, M.; Chalier, F.; Ouari, O.; Finet, J.; Rockenbauer, A.; Kalyanaraman, B.; Tordo, P. *Chem. Commun.* **2007**, 1083–1085.
- (17) Chang, M. C. Y.; Pralle, A.; Isacoff, E. Y.; Chang, C. J. *J. Am. Chem. Soc.* **2004**, *126*, 15392–15393.
- (18) Miller, E. W.; Tulyathan, O.; Isacoff, E. Y.; Chang, C. J. *Nat. Chem. Biol.* **2007**, *3*, 263–267.
- (19) McCormack, A. L.; Thiruchelvam, M.; Manning-Bog, A. B.; Thiffault, C.; Langston, J. W.; Cory-Clechta, D. A.; Di Monte, D. A. *Neurobiol. Dis.* **2002**, *10*, 119–127.
- (20) Park, S. Y.; Choi, J. *Environ. Int.* **2007**, *33*, 817–822.

JA802355U