


Palladium-Catalyzed Amination of Aryl and Heteroaryl Tosylates at Room Temperature

Tokutaro Ogata, and John F. Hartwig

J. Am. Chem. Soc., **2008**, 130 (42), 13848-13849 • DOI: 10.1021/ja805810p • Publication Date (Web): 24 September 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



Palladium-Catalyzed Amination of Aryl and Heteroaryl Tosylates at Room Temperature

Tokutaro Ogata and John F. Hartwig*

Department of Chemistry, University of Illinois, 600 South Mathews Avenue, Urbana, Illinois 61801


Received July 24, 2008; E-mail: jhartwig@uiuc.edu

Palladium-catalyzed aminations of aryl halides occur under mild conditions with many catalysts,^{1–3} but aminations of aryl sulfonates have been more challenging. The coupling of amines with aryl triflates has been known for many years,¹ but aryl triflates are expensive to prepare and less stable to hydrolysis than other aryl sulfonates. Aryl tosylates are less expensive to prepare and are stable, crystalline solids. However, they are also less reactive. Although Suzuki and Kumada couplings of aryl tosylates at room temperature have been reported,^{4,5a} only one coupling of amines with aryl tosylates at room temperature or with low catalyst loadings has been reported.⁵ Moreover, the coupling of amines with heteroaryl tosylates has not been reported, and the couplings of primary amines with aryl tosylates are rare.

We previously reported the oxidative addition of aryl tosylates at room temperature,^{5a} but this fast addition did not lead to a general coupling with amines under similar conditions. Here, we report a new method to create a catalyst that couples aryl tosylates with primary alkylamines, arylamines, and N–H imines with fast rates and high turnover numbers under mild conditions. This process was achieved by combining the hindered Josiphos ligand CyPF-*t*-Bu (Table 1) with the unconventional Pd(0) precursor Pd[P(*o*-tol)₃]₂, which we show can be made in a practical manner.

The difference between the rates of oxidative addition and the catalytic coupling stems from the difficulty in accessing the (CyPF-*t*-Bu)Pd intermediate under mild conditions. Typical catalyst precursors include Pd(II) acetate or chloride complexes, which require reduction to the true Pd(0) catalyst. To avoid the need for reduction to Pd(0), Pd_n(dba)_m is often used, but Pd(CyPF-*t*-Bu)(dba) generated from CyPF-*t*-Bu and Pd₂(dba)₃ dissociates dba particularly slowly, presumably because the strongly donating bisphosphine leads to strong backbonding into the electron-poor dba ligand. Thus, an alternative precatalyst was needed to couple aryl tosylates with amines under mild conditions.

Instead of using the combination of CyPF-*t*-Bu and typical catalyst precursors, we sought a practical synthesis of Pd(CyPF-*t*-Bu)[P(*o*-tol)₃]₂ (1), which oxidatively adds aryl tosylates at room temperature.^{5a} Pd(0) catalyst 1 is generated in solution at room temperature within minutes from Pd[P(*o*-tol)₃]₂ (2) and CyPF-*t*-Bu (3),⁶ but 2 has rarely been used in catalytic chemistry because of its lack of commercial availability and difficult synthesis.⁷ Thus, we sought a more convenient method to prepare 1 independently or a convenient method to prepare precursor 2.



Our efforts to isolate 1 in high yield were hampered by the high solubility of this species in organic solvents. Thus, we sought to simplify the synthesis of Pd(0) complex 2 and to use 1 generated in situ. On the basis of the most efficient synthesis of Pd[P(*t*-Bu)₃]₂,⁸ we prepared 2 in 95% isolated yield from the reaction of P(*o*-tol)₃ and Pd(dba)₂ in DMF.⁹ Complex 2 precipitated after 1 h at room temperature without the need for excess P(*o*-tol)₃. Complex 2 is stable for > 1 week in air as a

Table 1. Coupling of Aryl and Heteroaryl Tosylates with Primary Alkylamines and Arylamines Catalyzed by Pd[P(*o*-tol)₃]₂ and CyPF-*t*-Bu (1:1)^a

entry	substrate	R	cat. (%)	time (h)	yield (%) ^b		
1		X = OTs	octyl	0.1	24	94	
2 ^c		OTs	octyl	0.1	10 min	97	
3		OTf	octyl	0.1	48	29	
4		OMs	octyl	0.1	24	– ^d	
5		OSO ₂ Ph-4-F	octyl	0.1	48	76	
6		R ₁ = H	^t Bu	0.1	24	87	
7		H	³ Bu	0.2	24	83	
8		H	⁹ Hex	0.1	24	72	
9		H	Bn	0.1	24	90	
10		H	1-methylbenzyl ^e	1.0	24	95 ^f	
11		H	PhNHCH ₂ H ₄ NH ₂ ^g	0.5	24	96 ^h	
12		H	Ph ₂ C=N ^h	0.1	48	91	
13		Me	⁹ Bu	0.2	24	99	
14			octyl	0.05	24	97	
15 ^c			octyl	0.01	48	84	
16			R ₁ = CN	octyl	0.5	24	99
17 ⁱ			CO ₂ Et	octyl	1.0	24	99
18			CO ₂ ^t Bu	octyl	1.0	24	88
19	OMe		Bn	0.5	24	87	
20		4-Me-C ₆ H ₄	0.1	24	87		
21		4-OMe-C ₆ H ₄	0.1	24	77		
22		2-Me-C ₆ H ₄	0.2	24	75		
23 ^j		2-pyridyl	1.0	24	71		
24		3-OMe-5-CF ₃ -C ₆ H ₄	1.0	48	62(88 ^k)		
25		4-Me-C ₆ H ₄	0.1	24	99		
26		2-Me-C ₆ H ₄	0.2	24	70		
27		4-Me-C ₆ H ₄	0.2	24	96		
28		octyl	0.5	24	90		
29		4-OMe-C ₆ H ₄	1.0	48	63		
30		octyl	1.0	24	80		
31		octyl	1.0	24	82		
32		octyl	0.2	24	91		
33		^t Bu	0.2	24	97		
34		Bn	0.2	24	87		
35		4-Me-C ₆ H ₄	1.0	24	78		

^a Reactions conducted with a 1:1 ratio of metal to ligand, 1 mmol ArOTs, 1.2 equiv amine and 1.4 equiv NaOt-Bu in 1 mL toluene. ^b Isolated yields are an average of two runs. ^c At 80 °C. ^d Phenol was formed in 23% yield. ^e From phenethylamine that is stated to be 99% ee. ^f Produced in 99% ee. ^g Run using 1.5 equiv amine. ^h *N,N*-diphenylethylenediamine was obtained as a single product. ⁱ Reaction conducted with 1.2 equiv K₃PO₄ at 100 °C. ^j At 110 °C. ^k Run using 2.0 mol % catalyst.

solid at room temperature.¹⁰ Because CyPF-*t*-Bu is also stable in air, both catalyst components can be stored outside of a

drybox, and a solid mixture of the two materials can mimic a single-component catalyst (vide infra).¹¹

Examples of the reactions of aryl tosylates and heteroaryl tosylates with various alkylamines are summarized in Table 1. Reactions with 0.05–1.0 mol % of the combination of **2** and **3** in toluene at room temperature produced the corresponding monoarylamines in excellent yields. Most reactions were complete at room temperature within 24 h. However, similar yields could also be obtained after short (<10 min) times at 80 °C.

Reactions of linear primary amines were fast and occurred in high yield with 0.05–0.1 mol % catalyst (entries 1, 6, 9, and 14). The reaction of octylamine with 2-naphthyl tosylate was particularly efficient and occurred at 80 °C in 84% isolated yield with only 0.01 mol % of catalyst (entry 15). This turnover number of 8400 is the highest for any type of coupling of an aryl tosylate of which we are aware.^{12–14} The reactions of hindered α -branched primary amines, such as cyclohexylamine and *sec*-butylamine, occurred in high yields at room temperature with just 0.1 to 0.2 mol % of catalyst (entries 7–8, and 13). Both electron-rich and electron-poor aryl tosylates reacted in high yield with primary amines without formation of any diarylamine. These reactions occurred with aryl tosylates containing ortho substituents, as well as aryl tosylates containing cyano and carboalkoxy groups.

Because this new catalyst does not require reduction to a palladium(0) species or dissociation of dba, primary arylamines also couple with aryl tosylates at room temperature. Although these reactions were slightly slower than those of alkylamines, reactions of electron-rich arylamines (entries 20–21, 25, and 27), including an ortho-substituted arylamine (entries 22 and 26), occurred in good-to-excellent yield with complete selectivity for monoarylation. Reactions of electron-deficient primary aryl- and heteroarylamines also occurred. Although these reactions required higher catalyst loadings than the reactions of electron-rich primary arylamines (entries 23–24), they still occurred at elevated temperatures or with extended reaction times at room temperature.

The scope of the couplings of this catalyst also includes the first aminations of heteroaryl tosylates (entries 28–35). These reactions occurred under the same conditions we developed for the amination of aryl tosylates. Pyridyl and quinolyl tosylates underwent reaction with primary alkyl- and arylamines in good-to-excellent yields. The reactions of 2- and 3-pyridyl and 6- and 8-quinolyl tosylate also occurred, although 0.5 and 1.0 mol % catalyst was needed in some cases.


Consistent with the high selectivity for reactions of primary amines, the reactions of secondary amines were slower. The reactions of aryl tosylates with morpholine, dibutylamine, and *N*-methylaniline gave no coupled products or low yield of products at 110 °C with 1 mol % catalysts.

A comparison of these reactions to those of aryl tosylates conducted with other precatalysts showed the value of initiating reactions with P(*o*-tol)₃-ligated **2**. No reaction of octylamine with phenyl tosylate catalyzed by 0.1 mol % Pd(dba)₂ and **3** occurred at room temperature. The same reaction catalyzed by 0.1 mol % of Pd(OAc)₂ and **3** at room temperature for 48 h gave the product in 6% yield, as determined by GC, and this reaction catalyzed by 0.1 mol % PdCl₂(CyPF-*t*-Bu) occurred in only 65% isolated yield. Indicating the importance of Josiphos ligand **3**, the reaction of phenyl tosylate with octylamine catalyzed by the combination of 0.1 mol % Pd(OAc)₂, Pd(dba)₂, or Pd[P(*o*-tol)₃]₂ as precursor and 0.1–0.25 mol % Q-phos,¹⁵ X-phos,^{16,17} SIPr,³ DPPF, or BINAP as ligand gave no coupled products or very low yield of products, even in toluene at 110 °C (see Supporting Information for details).

Finally, a solid mixture of the two catalyst components catalyzes the reaction with efficiency equal to that of the catalyst generated from the two separate solids. Although expected, this procedure does allow the use of the combination of **2** and **3** as if it were an air-stable single-component catalyst.

A comparison of the reactions of different types of aryl sulfonates revealed some unusual trends (Table 1, entries 1–5). Most striking,

reactions of aryl tosylates were faster than those of aryl triflates when conducted as separate reactions (c.f. Table 1, entry 1 vs entry 3). Reactions of octylamine with a 1:1 ratio of these two aryl sulfonates formed mostly phenol (14%) with only 4% yield of the coupled product in the presence of 0.1 mol % catalyst. Reactions with Cs₂CO₃¹⁸ did not improve the yield from reactions of triflates. In addition, aryl tosylates reacted faster than aryl chlorides. The reaction of 3- or 4-chlorophenyl tosylate led to coupling at the tosylate group (eq 2). These relative rates contrast those catalyzed by complexes of dialkyl-*o*-biarylphosphines,¹⁹ but parallel those catalyzed by complexes of secondary phosphine oxides.¹³



In summary, we have developed a highly efficient catalyst system for the amination of aryl tosylates at room temperature, as well as the first examples of the Pd-catalyzed amination of heteroaryl tosylates. The use of an unusual precatalyst now prepared in a practical fashion enabled us to achieve this high activity.

Acknowledgment. We thank the NIH (GM-55382) for support of this work, Johnson-Matthey for a gift of PdCl₂, and Solvias for a gift of CyPF-*t*-Bu. T.O. thanks JSPS for a fellowship.

Supporting Information Available: All experimental procedures and spectroscopic data of new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Hartwig, J. F. *Acc. Chem. Res.* published online, Aug. 6, 2008, <http://dx.doi.org/10.1021/ar800098p>. (b) Hartwig, J. F. In *Modern Arene Chemistry*; Astruc, D., Ed.; Wiley-VCH: Weinheim, Germany, 2002; pp 107. (c) Hartwig, J. F. In *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; Wiley-Interscience: New York, 2002; pp 1051. (d) Muci, A. R.; Buchwald, S. L. *Top. Curr. Chem.* **2002**, *219*, 131.
- (2) For a recent review of aminations conducted with *o*-biaryl dialkylphosphine ligands, see: Surry, D. S.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2008**, *47*, 6338.
- (3) For a recent report of aminations conducted with *N*-heterocyclic carbenes, see: Marion, N.; Navarro, O.; Mei, J.; Stevens, E. D.; Scott, N. M.; Nolan, S. P. *J. Am. Chem. Soc.* **2006**, *128*, 4101.
- (4) (a) Tang, Z.-Y.; Hu, Q.-S. *J. Am. Chem. Soc.* **2004**, *126*, 3058. (b) Tang, Z.-Y.; Spinella, S.; Hu, Q.-S. *Tetrahedron Lett.* **2006**, *47*, 2427. (c) Kobayashi, Y.; Mizojiri, R. *Tetrahedron Lett.* **1996**, *37*, 8531. (d) Limmert, M. E.; Roy, A. H.; Hartwig, J. F. *J. Org. Chem.* **2005**, *70*, 9364. (e) Fürstner, A.; Leitner, A.; Mendez, M.; Krause, H. *J. Am. Chem. Soc.* **2002**, *124*, 13856. (f) Fürstner, A.; Leitner, A. *Angew. Chem., Int. Ed.* **2002**, *41*, 609.
- (5) (a) Roy, A. H.; Hartwig, J. F. *J. Am. Chem. Soc.* **2003**, *125*, 8704. For two examples of the amination of aryl tosylates at elevated temperatures with a related catalyst, see: (b) Hamann, B. C.; Hartwig, J. F. *J. Am. Chem. Soc.* **1998**, *120*, 7369. For recent work on the amination of aryl mesylates, at elevated temperatures, see: (c) So, C. M.; Zhou, Z.; Lau, C. P.; Kwong, F. Y. *Angew. Chem., Int. Ed.* **2008**, *47*, 6402.
- (6) Ligand **3** is commercially available from Solvias and Strem.
- (7) (a) Paul, F.; Patt, J.; Hartwig, J. F. *Organometallics* **1995**, *14*, 3030. (b) Böhm, V. P. W.; Herrmann, W. A. *Chem.—Eur. J.* **2001**, *7*, 4191.
- (8) Dai, C.; Fu, G. C. *J. Am. Chem. Soc.* **2001**, *123*, 2719.
- (9) Complex **2** is commercially available from Johnson-Matthey and Strem Chemicals (#46-0262).
- (10) Samples left in air as a solid for one week were unchanged, as judged by ¹H NMR and ³¹P NMR spectroscopy.
- (11) Naber, J. R.; Buchwald, S. L. *Adv. Synth. Catal.* **2008**, *350*, 957.
- (12) For reactions of activated aryl tosylates with 0.5 mol % palladium, see refs 13 and 14.
- (13) Ackermann, L.; Althammer, A. *Org. Lett.* **2006**, *8*, 3457.
- (14) Zhang, L.; Meng, T.; Wu, J. *J. Org. Chem.* **2007**, *72*, 9346.
- (15) Shelby, Q.; Kataoka, N.; Mann, G.; Hartwig, J. F. *J. Am. Chem. Soc.* **2000**, *122*, 10718.
- (16) Huang, X.; Anderson, K. W.; Zim, D.; Jiang, L.; Klapars, A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2003**, *125*, 6653.
- (17) Reaction with 2 mol % Pd(OAc)₂ and 5 mol % X-phos at 110 °C for 18 h occurred to 84% conversion.
- (18) Åhman, J.; Buchwald, S. L. *Tetrahedron Lett.* **1997**, *38*, 6363.
- (19) Nguyen, H. N.; Huang, X.; Buchwald, S. L. *J. Am. Chem. Soc.* **2003**, *125*, 11818.

JA805810P