


Design of Quantum Dot-Conjugated Lipids for Long-Term, High-Speed Tracking Experiments on Cell Surfaces

Michael J. Murcia, Daniel. E. Minner, Gina-Mirela Mustata, Kenneth Ritchie, and Christoph A. Naumann

J. Am. Chem. Soc., **2008**, 130 (45), 15054-15062 • DOI: 10.1021/ja803325b • Publication Date (Web): 21 October 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Design of Quantum Dot-Conjugated Lipids for Long-Term, High-Speed Tracking Experiments on Cell Surfaces

Michael J. Murcia,[†] Daniel. E. Minner,[†] Gina-Mirela Mustata,[‡] Kenneth Ritchie,[‡] and
Christoph A. Naumann^{*†}

*Department of Chemistry and Chemical Biology, Indiana University—Purdue University
Indianapolis, Indianapolis, Indiana 46202-3274, and Department of Physics,
Purdue University—West Lafayette, West Lafayette, Indiana 47907-2036*

Received May 5, 2008; E-mail: Naumann@chem.iupui.edu

Abstract: The current study reports the facile design of quantum dot (QD)-conjugated lipids and their application to high-speed tracking experiments on cell surfaces. CdSe/ZnS core/shell QDs with two types of hydrophilic coatings, 2-(2-aminoethoxy)ethanol (AEE) and a 60:40 molar mixture of 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine and 1,2-dipalmitoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[methoxy(polyethylene glycol-2000)], are conjugated to sulfhydryl lipids via maleimide reactive groups on the QD surface. Prior to lipid conjugation, the colloidal stability of both types of coated QDs in aqueous solution is confirmed using fluorescence correlation spectroscopy. A sensitive assay based on single lipid tracking experiments on a planar solid-supported phospholipid bilayer is presented that establishes conditions of monovalent conjugation of QDs to lipids. The QD-lipids are then employed as single-molecule tracking probes in plasma membranes of several cell types. Initial tracking experiments at a frame rate of 30 frames/s corroborate that QD-lipids diffuse like dye-labeled lipids in the plasma membrane of COS-7, HEK-293, 3T3, and NRK cells, thus confirming monovalent labeling. Finally, QD-lipids are applied for the first time to high-speed single-molecule imaging by tracking their lateral mobility in the plasma membrane of NRK fibroblasts with up to 1000 frames/s. Our high-speed tracking data, which are in excellent agreement with previous tracking experiments that used larger (40 nm) Au labels, not only push the time resolution in long-time, continuous fluorescence-based single-molecule tracking but also show that highly photostable, photoluminescent nanoprobes of 10 nm size can be employed (AEE-coated QDs). These probes are also attractive because, unlike Au nanoparticles, they facilitate complex multicolor experiments.

Introduction

The present surge in life science is linked intimately to the development of new experimental tools that allow the study of biological processes at the molecular level. In particular, optical single-molecule imaging techniques have emerged as powerful tools to detect heterogeneities below the diffraction limit of optical microscopy and to study dynamic processes in cellular systems at the single-molecule level because these techniques enable individual molecules to be tracked with a spatial resolution of 20–30 nm and a time resolution in the micro-second range. For example, single-molecule imaging techniques have been instrumental in changing our view of plasma membranes from a featureless lipid bilayer with embedded membrane proteins¹ to a complex, compartmentalized system exhibiting a wide variety of length scale-dependent dynamic processes.² Interestingly, not only labeled membrane proteins but also lipid-based single-molecule tracking probes have emerged as powerful tools in the in-depth characterization of

membrane heterogeneities.^{3,4} Traditionally, the detection of single biomolecules has been accomplished via conjugated colloidal gold or fluorescent latex spheres using nanovided microscopy and single-particle tracking.^{5–11} The main limitations of these traditional probes for single-molecule imaging are their relatively large size (gold, 40–50 nm; latex, 0.1–1 μm), which considerably exceeds that of the biomolecules labeled, and the difficulty to avoid cross-linking of biomolecules. These limitations motivated the development of single-molecule imaging using fluorescent dye labels. Single fluorophores were first successfully imaged at very low temperature using an

[†] Indiana University—Purdue University Indianapolis.

[‡] Purdue University—West Lafayette.

(1) Singer, S. J.; Nicolson, G. L. *Science* **1972**, *175*, 720–731.

(2) Kusumi, A.; Nakada, C.; Ritchie, K.; Murase, K.; Suzuki, K.; Murakoshi, H.; Kasai, R. S.; Kondo, J.; Fujiwara, T. *Annu. Rev. Biophys. Biomol. Struct.* **2005**, *34*, 351–378.

(3) Fujiwara, T.; Ritchie, K.; Murakoshi, H.; Jacobson, K.; Kusumi, A. *J. Cell Biol.* **2002**, *157*, 1071–1082.

(4) Murase, K.; Fujiwara, T.; Umemura, Y.; Suzuki, K.; Iino, R.; Yamashita, H.; Saito, M.; Murakoshi, H.; Ritchie, K.; Kusumi, A. *Biophys. J.* **2004**, *86*, 4075–4093.

(5) Geerts, H.; De Brabander, M.; Nuydens, R.; Geuens, S.; Moeremans, M.; De Mey, J.; Hollenbeck, P. *Biophys. J.* **1987**, *52*, 775–782.

(6) Sheetz, M. P.; Turney, S.; Qian, H.; Elson, E. L. *Nature* **1989**, *340*, 284–288.

(7) Lee, G. M.; Ishihara, A.; Jacobson, K. A. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 6274–6278.

(8) Ghosh, R. N.; Webb, W. W. *Biophys. J.* **1994**, *66*, 1301–1318.

(9) Qian, H.; Sheetz, M. P.; Elson, E. L. *Biophys. J.* **1991**, *60*, 910–921.

(10) Fein, M.; Unkeless, J.; Chuang, F. Y. S.; Sassaroli, M.; da Costa, R.; Väänänen, H.; Eisinger, J. *J. Membr. Biol.* **1993**, *135*, 83–92.

(11) Kusumi, A.; Sako, Y.; Yamamoto, M. *Biophys. J.* **1993**, *65*, 2021–2140.

intensified CCD camera.¹² Diffusion studies of membrane constituents became possible after the time resolution of individual dye tracking was increased into the millisecond range.¹³ Although dyes overcome limitations of colloidal Au probes in terms of size and monovalent labeling of biomolecules, their application to single-molecule tracking remains limited due to poor photostability, thus limiting the length of individual trajectories.


More recently, photoluminescent quantum dots (QDs) have emerged as an attractive alternative for fluorescence-based imaging because they combine small size, brightness, high photostability, and broad absorption and narrow, size-tunable emission bands.^{14–16} In particular, the superior brightness and photostability make QDs powerful probes for the single-molecule tracking with very high time/spatial resolution. Due to their unique absorbance and emission properties, QDs are also particularly beneficial for multicolor experiments because just one laser excitation wavelength is needed to excite QDs of different emission properties. To allow biological imaging applications, QDs need to be made water-soluble using hydrophilic surface coatings.^{17,18} Initially, this was achieved by replacing the coordinating solvent trioctylphosphine oxide with monothiols containing hydrophilic terminal moieties, such as 3-mercaptopropionic acid and mercaptoethanol.^{19–23} Although these coatings are very thin, they show limited stability.²⁴ Therefore, several alternative coatings strategies have been developed, which are based on silanes forming a stable shell via cross-linking,^{16,25,26} peptides,²⁷ polyelectrolytes,²⁸ polysac-

charides,²⁹ avidin,³⁰ and QD-encapsulating amphiphilic molecules, such as hydrophilic polymers with hydrophobic side chains or mixtures of phospholipids and lipopolymers.³¹ The encapsulation with amphiphiles is interesting because the protecting coordinating solvent layer does not need to be replaced, thus preventing oxidative damage. Since their introduction as biological imaging probes in 1998,^{32,33} QDs have found widespread application in biological imaging at the *in vitro* and *in vivo* levels.^{14–16} Dahan and co-workers first successfully demonstrated that QDs are excellent imaging probes for single-molecule tracking experiments.³⁴ In their initial work, QDs were employed to study the lateral diffusion of glycine receptors and their entry into neuronal synapses. Meanwhile, QDs have been employed as single-molecule tracking probes to analyze the lateral mobility of several membrane receptors, including EGF,³⁵ GABA,³⁶ and Kv2.1.³⁷ In these applications, streptavidin-coated QDs were conjugated to either biotinylated membrane receptors or receptor ligands. Streptavidin or avidin coatings are attractive because they allow for the strong, specific binding of QDs to biotinylated biomolecules. Their disadvantages, however, are the relatively large coating thickness, leading to probe sizes of more than 15 nm, and the difficulty to monovalently label biomolecules (each streptavidin or avidin molecule binds up to four biotins).

Here we report for the first time the successful monovalent conjugation of 2-(2-aminoethoxy)ethanol (AEE)- and phospholipid/lipopolymer (LIPO)-capped CdSe/ZnS quantum dots to phospholipids and their application in lipid tracking experiments on the cell surface using high-speed single-molecule fluorescence microscopy. The thicker LIPO coating was chosen because it is very stable and was already successfully applied in long-term cellular imaging studies.³⁸ The thinner AEE coating was pursued because the overall probe size can be reduced and because a similar coating, which is based on short poly(ethylene oxide) chains of just two ethylene oxide segments, was shown to prevent the nonspecific adsorption of proteins quite effectively.³⁹ As reported recently, confocal fluorescence correlation spectroscopy (FCS) is employed to verify the colloidal stability of AEE- and LIPO-coated QDs and to determine their hydrodynamic radii.⁴⁰ QD conjugation to lipids is achieved by incorporating a well-adjusted amount of maleimide-containing

- (12) (a) Güttler, F.; Irmgartinger, T.; Plakhotnik, T.; Renn, A.; Wild, U. P. *Chem. Phys. Lett.* **1994**, *217*, 393–397. (b) Moerner, W. E.; Plakhotnik, T.; Irmgartinger, T.; Croci, M.; Palm, V.; Wild, U. W. *J. Phys. Chem.* **1994**, *98*, 7382–7389.
- (13) Schmidt, T.; Schuetz, G. J.; Baumgartner, W.; Gruber, H. J.; Schindler, H. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 2926–2929.
- (14) Alivisatos, A. P.; Gu, W. W.; Larabell, C. *Annu. Rev. Biomed.* **2005**, *7*, 55–76.
- (15) Chan, W. C. W. *Biol. Blood Marrow Transplant* **2006**, *12* (1, Suppl. 1), 87–91.
- (16) Michalet, X.; Pinaud, F. F.; Bentolila, L. A.; Tsay, J. M.; Doose, S.; Li, J. J.; Sundaresan, G.; Wu, A. M.; Gambhir, S. S.; Weiss, S. *Science* **2005**, *307*, 538–544.
- (17) Parak, W. J.; Gerion, D.; Pellegrino, T.; Zanchet, D.; Micheel, C.; Williams, S. C.; Boudreau, R.; Le Gros, M.; Larabell, C. A.; Alivisatos, A. P. *Nanotechnology* **2003**, *14*, 15–27.
- (18) Murcia, M. J.; Naumann, C. A. Biofunctionalization of fluorescent nanoparticles. In *Biofunctionalization of Nanoparticles*; Kumar, C., Ed.; Wiley-VCH: Weinheim, 2005; pp 1–40.
- (19) Peng, X.; Schlamp, M. C.; Kadavanich, A. V.; Alivisatos, A. P. *J. Am. Chem. Soc.* **1997**, *119*, 7019–7029.
- (20) Passow, T.; Leonardi, K.; Hommel, D. *Phys. Status Solidi* **2001**, *224*, 143–146.
- (21) Pathak, S.; Choi, S. K.; Arnheim, N.; Thompson, M. E. *J. Am. Chem. Soc.* **2001**, *123*, 4103–4104.
- (22) Mattoussi, H.; Mauro, J. M.; Goldman, E. R.; Anderson, G. P.; Sundar, V. C.; Mikulec, F. V.; Bawendi, M. G. *J. Am. Chem. Soc.* **2000**, *122*, 12142–12150.
- (23) Mattoussi, H.; Mauro, J. M.; Goldman, E. R.; Green, T. M.; Anderson, G. P.; Sundar, V. C.; Bawendi, M. G. *Phys. Status Solidi* **2001**, *224*, 277–283.
- (24) Aldana, J.; Wang, Y. A.; Peng, X. G. *J. Am. Chem. Soc.* **2001**, *123*, 8844–8850.
- (25) Gerion, D.; Pinaud, F.; Williams, S. C.; Parak, W. J.; Zanchet, D.; Weiss, S.; Alivisatos, A. P. *J. Phys. Chem.* **2001**, *105*, 8861–8871.
- (26) Parak, W. J.; Gerion, D.; Zanchet, D.; Woerz, A. S.; Pellegrino, T.; Micheel, C.; Williams, S. C.; Seitz, M.; Bruehl, R. E.; Bryant, Z.; Bustamante, C.; Bertozzi, C. R.; Alivisatos, A. P. *Chem. Mater.* **2002**, *14*, 2113–2119.
- (27) Lagerholm, B. C.; Wang, M.; Ernst, L. A.; Ly, D. H.; Liu, H.; Bruchez, M. P.; Waggoner, A. S. *Nano Lett.* **2004**, *4*, 2019–2022.
- (28) Jaffar, S.; Nam, K. T.; Khademhosseini, A.; Xing, J.; Langer, R. S.; Belcher, A. M. *Nano Lett.* **2004**, *4*, 1421–1425.

- (29) Xie, M.; Liu, H. H.; Chen, P.; Zhang, Z. L.; Wang, X. H.; Xie, Z. X.; Du, Y. M.; Pan, B. Q.; Pang, D. W. *Chem. Commun.* **2005**, *44*, 5518–5520.
- (30) Goldmann, E. R.; Balighian, E. D.; Mattoussi, H.; Kuno, M. K.; Mauro, J. M.; Tran, P. T.; Erson, G. P. *J. Am. Chem. Soc.* **2002**, *124*, 6378–6382.
- (31) Dubertret, B.; Skourides, P.; Norris, D. J.; Noireaux, V.; Brivanlou, A. H.; Libchaber, A. *Science* **2002**, *298*, 1759–1762.
- (32) Brunchez, M. J.; Moronne, M.; Gin, P.; Weiss, S.; Alivisatos, A. P. *Science* **1998**, *281*, 2013–2016.
- (33) Chan, W. C. W.; Nie, S. *Science* **1998**, *281*, 2016–2018.
- (34) Dahan, M.; Levi, S.; Luccardini, C.; Rostaing, P.; Riveau, B.; Triller, A. *Science* **2003**, *302*, 442–445.
- (35) Lidke, D. S.; Nagy, P.; Heintzmann, R.; Arndt-Jovin, D. J.; Post, J. N.; Grecco, H. E.; Jares-Erijman, E. A.; Jovin, T. M. *Nat. Biotechnol.* **2004**, *22*, 198–203.
- (36) Bouzigues, C.; Morel, M.; Triller, A.; Dahan, M. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104*, 11251–11256.
- (37) Tamkun, M. M.; O'Connell, K. M. S.; Roliq, A. S. *J. Cell Sci.* **2007**, *120*, 2413–2423.
- (38) Chapman, R. G.; Ostuni, E.; Liang, M. N.; Meluleni, G.; Kim, E.; Yan, L.; Pier, G.; Warren, H. S.; Whitesides, G. M. *Langmuir* **2001**, *17*, 1225–1233.
- (39) Murcia, M. J.; Shaw, D. L.; Long, E. L.; Naumann, C. A. *Opt. Commun.* **2008**, *281*, 1771–1780.
- (40) Murcia, M. J.; Shaw, D. L.; Woodruff, H.; Naumann, C. A.; Young, B. Y.; Long, E. L. *Chem. Mater.* **2006**, *18*, 2219–2225.

Scheme 1. Reaction Scheme Illustrating the Synthesis Steps for the Design of AEE-Coated QDs

cross-linker molecules into the QD coatings and by reacting QDs with sulfhydryl lipids. To evaluate the inertness of the QD probes in the presence of a membrane and to confirm the monovalent labeling of QDs to lipids, the lateral diffusion of QD-conjugated lipids is first examined in a solid-supported phospholipid bilayer and compared to diffusion data obtained using dye-labeled lipids. QD-lipids are then incorporated into plasma membranes of several cell types, including COS-7, HEK-293, 3T3, and NRK fibroblasts, and their lateral diffusion behavior is examined using a frame rate of 30 frames per second (fps). Finally, high-speed imaging experiments on NRK fibroblasts are presented, where QD-lipids are tracked with frame rates of up to 1000 fps. The results from these experiments are in very good agreement with previous single particle tracking data using notably larger colloidal Au probes. By pushing the time resolution in fluorescence-based wide-field single-molecule imaging, we show that QD-lipids are promising imaging probes in state-of-the-art high-speed single-molecule tracking experiments on cell surfaces.

Experimental Section

Materials. Cadmium acetate and elemental selenium powder were purchased from Strem Chemicals (Newburyport, MA). Trioctylphosphine (TOP), trioctylphosphine oxide (TOPO), hexadecylamine (HDA), 3-mercaptopropionic acid (MPA), 2-(2-aminoethoxy)ethanol (AEE), di-*tert*-butyl dicarbonate (DiBOC), trifluoroacetic acid (TFA), zinc nitrate hexahydrate, potassium ethylxanthate, and HPLC-grade solvents were purchased from Aldrich Chemical (Milwaukee, WI). The heterobifunctional cross-linker reagent *N*-[*p*-maleimidophenyl]isocyanate (PMPI) and amide bond catalyst 1-ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC) were obtained from Pierce Biotechnology (Rockford, IL). The phospholipids 1-stearoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (SOPC), 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC), 1,2-dioleoyl-*sn*-glycero-3-phosphoethanolamine (DOPE), and 1,2-dipalmitoyl-*sn*-glycero-3-phosphothioethanol (DHPTE), as well as the lipopolymers 1,2-dipalmitoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[methoxy(polyethylene glycol)-2000] (DPPE-PEG2000) and 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[maleimide(polyethylene glycol)-2000] (ammonium salt) (DSPE-PEG2000-MAL), were purchased from Avanti Polar Lipids (Alabaster, AL). The

fluorescent lipid probe *N*-(6-tetramethylrhodaminethiocarbonyl)-1,2-dihexadecanoyl-*sn*-glycero-3-phosphoethanolamine (TRITC-DHPE) was acquired from Invitrogen/Molecular Probes (Eugene, OR). Water utilized was purified using a Milli-Q water purification system (Millipore, Milford, MA).

Methodology. (a). Synthesis of CdSe/ZnS QDs. CdSe/ZnS core/shell QDs were synthesized using a facile sonochemical approach, as reported previously.⁴⁰ In short, TOPO-coated CdSe QDs were synthesized in a one-pot synthesis using HDA, Cd(OAc)₂, and TOPO. The ZnS shell was added using zinc ethylxanthate. CdSe/ZnS core/shell QDs synthesized using this procedure were found to show high crystallinity, quantum yields of 50–60%, narrow emission spectra (fwhm ~25 nm), and size distributions of ~10%.⁴⁰

(b). Formation of 2-(2-Aminoethoxy)ethanol (AEE)-Coated CdSe/Zn QDs. Reaction Scheme 1 illustrates the synthesis steps for the formation of AEE-coated CdSe/ZnS QDs. The weakly bound TOPO coating of TOPO-CdSe/ZnS QDs was first replaced with the more tightly binding MPA. In a typical preparation, 0.5 g of the crude, shelled sample was washed and precipitated three times with a solution of 50:50 methanol/acetone and dissolved in 12 mL of chloroform with 0.25 g of MPA. The solution was stirred under argon for 24 h and centrifuged to separate out the hydrophilic nanocrystals. The nanocrystals were washed in hexane (three times) and dried under vacuum at room temperature. The dry product was then dispersed in water. The pH of the aqueous nanoparticle solution was then adjusted to 6.5, and 30 mg of AEE was added as the solution was stirred in an ice bath. The amide bond formation between MPA and AEE was catalyzed by the addition of 10 mg of EDC. To facilitate the linkage to phospholipids, AEE was mixed with the heterobifunctional cross-linker 2-(2-aminoethoxy)ethyl 4-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)phenylcarbamate (AEE to cross-linker molar mixing ratio, 100:1). Some control experiments were conducted using lower and higher molar cross-linker concentrations. To complete this reaction, the solution was stirred for 1 h at 5 °C. The solution was then centrifuged at 15 000 rpm and decanted to remove any excess AEE/cross-linker as well as any nanoparticle aggregates. The synthesis of the heterobifunctional cross-linker is described below.

(c). Synthesis of Heterobifunctional Amine–Maleimide Cross-Linker. First, the primary amine of AEE was protected with DiBOC. Here, 2.5 g of AEE was dissolved in 30 mL of water with 2 equiv (4.04 g) of sodium bicarbonate. A second solution

containing 5.19 g of DiBOC dissolved in 10 mL of dioxane was prepared and added to the AEE solution. The mixture was stirred for 24 h under argon, after which the protected AEE was extracted three times with 25 mL of ethyl acetate. The ethyl acetate solution was condensed to product via rotary evaporation. The resulting protected AEE product was dissolved in dry chloroform and reacted with the heterobifunctional cross-linker PMPI in a 1:2 ratio over 24 h. After completion of the reaction, the chloroform was vacuumed off. To remove the DiBOC protecting group, the orange, oily product was dissolved in 10 mL of a 5:1 dichloromethane (DCM)/TFA solution and stirred for 2 h. The DCM/TFA mixture was vacuumed off, giving a dry product. As verified by NMR, this final product is the carboxyl- and sulfhydryl-reactive heterobifunctional cross-linker 2-(2-aminoethoxy)ethyl 4-(2,5-dioxo-2,5-dihydro-1*H*-pyrrol-1-yl)phenylcarbamate.

(d). Formation of Lipopolymer (LIPO)-Encapsulated CdSe/Zn QDs. By adapting the coating procedure reported by Dubertret et al.,³¹ crude, shelled QDs were first washed and precipitated three times with a solution of 50:50 methanol/acetone to remove any excess coordinating solvent. The hydrophobic TOPO-capped nanoparticles were then dispersed in chloroform containing 5.5×10^{-6} mol of 40 mol % DPPE-PEG2000, 59 mol % DPPC, and 0.04 mol % DSPE-PEG2000-MAL. Some control experiments were also conducted using lower and higher DSPE-PEG2000-MAL molar concentrations. The chloroform was removed via evaporation, and the residue was heated to 80 °C before 1 mL of distilled water was added, resulting in an optically transparent solution of LIPO-coated QDs.

(e). Formation of QD-Conjugated Lipids in Solid-Supported Phospholipid Bilayer. Solid-supported phospholipid bilayers were formed via successive Langmuir–Blodgett (LB) and Langmuir–Schaefer (LS) film-transfer steps following standard procedures.^{41,42} Specifically, microscopy coverslips were cleaned by baking at 515 °C for 1 h, followed by sonication for 30 min each in the following solutions: 1% SDS, methanol saturated with NaOH, and 0.1% HCl, using a bath sonicator.⁴² After each washing step, the microscopy coverslips were rinsed in Milli-Q water. To form the first (LB) monolayer, a chloroform solution of SOPC was spread at the air–water interface of a film balance with a dipper (Labcon, Darlington, UK). Prior to film transfer, the lateral pressure of the SOPC monolayer was adjusted to 30 mN/m. To form the second (LS) monolayer, a chloroform solution containing SOPC and DHPTE (10⁻⁸ mol %) was spread on the air–water interface and compressed to 30 mN/m. In the case of dye experiments, DHPTE was replaced by the fluorescent TRITC-DHPE. After completion of the LB/LS transfer, the bilayer-containing cover glass was assembled as the bottom part of a cuvette with a glass cylinder perched atop it. In this open geometry, QD conjugation was achieved by adding maleimide-functionalized QDs to the cuvette and rinsing off the excess (unbound) QDs. To achieve monovalent labeling of QDs to lipids, the maleimide surface concentration on AEE- and LIPO-coated QDs was systematically increased, and their binding to a solid-supported SOPC bilayer containing 10⁻⁸ mol % DHPTE was studied using wide-field single-molecule fluorescence microscopy. For both the AEE and LIPO QDs, the required ratio of maleimide to inert surface molecules was determined to be 1:100. In addition, SOPC membranes containing no DHPTE were incubated with both maleimide-functionalized QD systems, and no specific binding was observed. Figure 1 illustrates schematics of AEE- (top, left) and LIPO-coated (top, right) QD-conjugated lipids and the design of QD-conjugated lipids in a solid-supported bilayer (bottom).

(f). Reconstitution of QD-Lipids into Plasma Membranes. A vesicle fusion technique was used to incorporate QD-labeled lipids

into living cell membranes. Small unilamellar vesicles (SUVs) were created in phosphate-buffered saline (PBS) buffer by a commonly used rod sonication technique, with the composition being a 3:1 ratio of SOPC/DOPE, as this mixture is known to be fusogenic with the plasma membrane of cells.^{43,44} In addition, the vesicles also contained a small quantity (10⁻³%) of the thiol-functionalized lipid, DHPTE. Based upon the number of DHPTE molecules in the vesicle solution, an order of magnitude fewer maleimide-functionalized QDs were added to the solution to bind the vesicles. As verified by FCS, this procedure ensured that there were no excess functionalized QDs in the vesicle solution which could potentially bind to proteins or other cysteine-containing biomolecules in the cell membrane. Upon binding the maleimide-functionalized QDs to the SUVs, they were allowed to fuse with the cells for 15–20 min before the excess vesicles were rinsed off and the cells were imaged. Quantum dots were tracked only if they remained in focus during lateral diffusion, thus excluding tracking artifacts due to either unfused or internalized imaging probes.

(g). Cell Culture. NRK fibroblasts were cultured in Dubelco's Minimum Essential Medium (DMEM) with 10% FetalPlex Animal Serum Complex (Gemini Bioproducts). Cells were plated on 18 mm diameter coverslips for single fluorophore imaging and used the next day. COS-7, HEK-293, and Swiss-3T3 cells were cultured by standard procedures, plated on 18 mm diameter coverslips for single fluorophore imaging, and used the next day.

Detailed information about the experimental methods (confocal fluorescence correlation spectroscopy, wide-field single-molecule fluorescence microscopy, and oblique angle fluorescence microscopy) is provided in the Supporting Information.

Results and Discussion

Prior to conjugation of QDs to lipids, the colloidal stability and the photophysical properties of capped QDs were determined in aqueous solution. Because FCS is a powerful method to obtain the necessary information,^{40,45,46} Figure 2 illustrates corresponding FCS autocorrelation curves of rhodamine dyes and of CdSe/ZnS QDs with MPA, AEE, and LIPO coatings obtained after $t = 10$ min (left) and $t = 1$ week (right).⁴¹ Using eq 2.3 (in Supporting Information), the autocorrelation curves in Figure 2A provide the following hydrodynamic radii after 10 min: $r(\text{rhodamine}) = 0.6$ nm, $r(\text{MPA}) = 3.3$ nm, $r(\text{AEE}) = 4.5$ nm, and $r(\text{LIPO}) = 14.6$ nm. As expected, the obtained radii indicate that AEE-coated QDs are smaller than LIPO-coated systems. The obtained radii also suggest that AEE- and LIPO-QDs exist in solution as individual, colloidally stable nanocrystals. The colloidal stability is evaluated more rigorously by comparing hydrodynamic radii determined from QD samples after $t = 10$ min (Figure 2, left) and $t = 1$ week (Figure 2, right). While the hydrodynamic radius of the MPA-QD sample, which was employed as a negative control, increased significantly to $r(\text{MPA}) = 12.1$ nm, those of the AEE- and LIPO-QDs samples remain largely unchanged, with $r(\text{AEE}) = 5.3$ nm and $r(\text{LIPO}) = 16.3$ nm. These findings not only verify the excellent colloidal stability of LIPO QDs, which already have been applied successfully in long-term cellular studies,³¹ but also highlight the promising properties of the smaller AEE QDs. The very good colloidal stability of AEE-coated QDs observed

(43) Farhood, H.; Serbina, N.; Huang, L. *Biochim. Biophys. Acta* **1995**, *1235*, 289–295.

(44) Miller, C. R.; Bondurant, B.; McLean, S. D.; McGovern, K. A.; O'Brien, D. F. *Biochemistry* **1998**, *37*, 12875–12883.

(45) Pellegrino, T.; Manna, L.; Kudera, S.; Liedl, T.; Koktysh, D.; Rogach, A. L.; Keller, S.; Raedler, J.; Natile, G.; Parak, W. J. *Nano Lett.* **2004**, *4*, 703–707.

(46) Zhang, P.; Li, L.; Dong, C.; Qian, H.; Ren, J. *Anal. Chim. Acta* **2005**, *546*, 46–51.

(41) Deverall, M. A.; Gindl, E.; Sinner, E.-K.; Besir, H.; Ruehe, J.; Saxton, M. J.; Naumann, C. A. *Biophys. J.* **2005**, *88*, 1875–1886.

(42) Garg, S.; Ruehe, J.; Luedtke, K.; Jordan, R.; Naumann, C. A. *Biophys. J.* **2007**, *92*, 1263–1270.




Figure 1. Schematic illustration of AEE- (top, left) and LIPO-coated QDs (top, right) and QD-conjugated lipids in a solid-supported phospholipid bilayer (bottom). The LIPO-coated QDs also contain phospholipids, which are omitted in the corresponding schematic.




Figure 2. FCS autocorrelation curves of MPA-, AEE-, and LIPO-QDs in 0.1 M PBS buffer, pH 7.0, after sample preparation at $t = 10$ min (left) and $t = 1$ week (right). The rhodamine data are added as a reference.

is reasonable because similar ultrathin poly(ethylene oxide) coatings consisting of not more than three ethylene oxide segments were effective in preventing the nonspecific adsorption of proteins.³⁸

The quality of the surface coatings was further evaluated by monitoring the photoluminescence properties of capped QDs in aqueous solution. Specifically, FCS was applied to determine the photon counts per particle of MPA-, AEE-, and LIPO-coated QDs over a time period of 1 week, which provided 550–690 for MPA-coated QDs, 530–640 for AEE-coated QDs, and 850–1000 for LIPO-coated QDs. Most

importantly, for all three coatings, no drop in the counts per particle was observed over the time period of 1 week. The lower counts per particle of MPA- and AEE-QDs relative to LIPO-QDs are caused partially by the fluorescence-reducing effect of the thiol exchange reaction employed in both cases. In addition, the reduced emission observed could result from the thinner barrier between the nanocrystal surface and water (relative to LIPO-QDs). If water does reach the nanoparticle surface, it can oxidize the nanocrystal, which decreases the emission efficiency. We have demonstrated this phenomenon by intentionally stripping off the dynamic surface-exchanged

nanoparticle coatings with repeated washings, or dialysis, resulting in quenched fluorescence (data not shown).

After AEE- and LIPO-coated QDs were verified to show good colloidal stability and photophysical properties in aqueous solution, their properties in a solid-supported phospholipid bilayer were studied. Initially, the inertness of maleimide-functionalized AEE- and LIPO-QDs was explored qualitatively by monitoring their nonspecific adsorption to a bilayer without sulfhydryl lipids using wide-field single-molecule fluorescence microscopy. These control experiments verified no nonspecific binding of QDs on the bilayer for both types of coated QDs (data not shown). In the next step, QDs were conjugated to sulfhydryl lipids in the lipid bilayer. To achieve monovalent labeling, which is essential for single-molecule imaging, the surface of the QD must exhibit the proper concentration of maleimide. In particular, too elevated maleimide concentrations may cause individual QDs to bind to multiple sulfhydryl lipids. To determine conditions where QDs are monovalently labeled to lipids, QDs with systematically varied maleimide concentrations were added to a solid-supported SOPC bilayer containing 10^{-8} mol % of maleimide-reactive DHPTE, and their binding behavior was analyzed using single-molecule fluorescence microscopy. Specifically, four samples of maleimide-functionalized QDs were created for both the heterobifunctional linker and the maleimide lipopolymer. The ratio of maleimide linker to AEE, or maleimide lipopolymer to lipopolymer, was increased for each subsequent QD sample. The linker/passive molecule ratios used were 1:10000, 1:1000, 1:100, and 1:10. Starting with the QD sample containing the lowest amount of maleimide groups, the samples were screened in order of increasing concentration for binding to DHPTE-containing membranes via single-molecule fluorescence microscopy. Monovalent labeling is expected at the lowest ratio of maleimide to inert surface molecules leading to QD binding, which was found to be 1:100 for AEE and LIPO coatings.

After verification of the inertness of AEE- and LIPO-QDs in the presence of a phospholipid bilayer, and after identification of the appropriate maleimide concentration on the QD surface for lipid tracking experiments, maleimide-functionalized QDs were tested as lipid tracking probes for single-molecule imaging on model membranes and were compared to dye tracking probes (TRITC-DHPE). Figure 3 compares square-displacement (r^2) histograms of TRITC-DHPE and LIPO-QD-conjugated lipids in a solid-supported SOPC bilayer for $t_{\text{lag}} = 50$ ms. Notably, statistically identical mean-square displacement values of $\langle r^2 \rangle_{\text{dye}} = 0.19 \pm 0.05 \mu\text{m}^2$ and $\langle r^2 \rangle_{\text{QD}} = 0.20 \pm 0.05 \mu\text{m}^2$ can be derived from both histograms. This excellent agreement between QD and dye probes is important because it shows that QDs did not cross-link multiple lipids and that monovalent labeling of QDs to sulfhydryl lipids was achieved. It should be emphasized that single and cross-linked lipids show different diffusion behavior.⁷

The great similarity of QD- and dye-based lipid tracking results in Figure 3 also corroborates the lack of nonspecific binding between lipid-conjugated QD probes and the SOPC bilayer. Of course, to investigate the inertness of lipid-conjugated QDs more rigorously, the diffusion properties need to be analyzed over a larger range of different time lags. Unlike dyes, individual QDs can be tracked over longer time periods due to their superior photostability. Figure 4 illustrates $\langle r^2 \rangle$ -time plots of AEE- and LIPO-QDs conjugated to a solid-supported SOPC bilayer which span a time period of 2 s. Both plots are well-described by linear fits, thus verifying Brownian diffusion of




Figure 3. Representative square-displacement (r^2) histograms obtained from single lipid tracking experiments on dye-labeled (TRITC-DHPE) and LIPO-QD-conjugated lipids in a solid-supported phospholipid bilayer. The resulting mean-square displacements of $\langle r^2 \rangle_{\text{dye}} = 0.19 \pm 0.01 \mu\text{m}^2$ and $\langle r^2 \rangle_{\text{QD}} = 0.20 \pm 0.01 \mu\text{m}^2$ are statistically identical and suggest that QD-lipids diffuse like their dye-labeled counterparts. The $\langle r^2 \rangle$ data are based on a time lag of $t_{\text{lag}} = 50$ ms.




Figure 4. $\langle r^2 \rangle$ -time plots of LIPO- and AEE-coated QDs conjugated to thiolipids in the top monolayer of a solid-supported SOPC bilayer over a time period of 2 s. The dashed lines show the best linear fits associated with the Brownian diffusion model. The identical slopes of both plots verify that LIPO and AEE coatings result in identical lipid diffusion properties.

QD-conjugated lipids in the SOPC bilayer. Furthermore, the almost identical slopes of the two plots also show that AEE- and LIPO-QD-conjugated lipids are characterized by the same lateral mobility. Overall, the conducted model membrane experiments successfully identified conditions of monovalent QD labeling of lipids, thus facilitating single-molecule tracking studies of these probes on cell surfaces.

To incorporate QD-lipids into plasma membranes, maleimide-functionalized AEE-QDs were bound to DHPTE in fusogenic SUVs composed of a mixture of SOPC and DOPE (molar ratio, 3:1) containing 10^{-3} mol % DHPTE. Again, to ensure mono-




Figure 5. FCS autocorrelation data and corresponding best fits of free AEE-QDs (top, left), vesicles labeled with the dye lipid TRITC-DHPE (top, right), maleimide-functionalized AEE-QDs bound to DHPTE-containing vesicles (bottom, left), and DHPTE-containing vesicles with bound and unbound maleimide-functionalized AEE-QDs (bottom, right). The data illustrate that QD-labeled vesicles can be prepared without free, excess QDs.




Figure 6. Diffusion coefficient histograms for QD-labeled lipids in three cell lines, COS 7, 3T3, and HEK293.

valent binding of QDs to DHPTE, the linker-to-passive molecule ratio on the QD surface was set to 1:100. Furthermore, to avoid an excess of unbound QDs in the vesicle solution, the amount of QDs added was kept 1 order of magnitude lower than the amount of DHPTE. Figure 5 shows FCS autocorrelation data and best-fitting curves of free QDs (top, left), vesicles labeled with dye-lipid TRITC-DHPE (top, right), QDs bound to vesicles (bottom, left), and QDs bound to vesicles + free excess QDs (bottom, right). Not surprisingly, free QDs and TRITC-labeled vesicles are best described by one-component fits with notably different diffusion times of $\tau_{D1} = 1533 \mu\text{s}$ and $\tau_{D1} = 3118 \mu\text{s}$, respectively. When all maleimide-functionalized QDs are bound to DHPTE-containing vesicles, the FCS data match those obtained using TRITC-containing vesicles. Again, the FCS curve is best described by a one-component fit. Furthermore, the diffusion time of $\tau_{D1} = 3256.5 \mu\text{s}$ is indistinguishable (within the experimental error of 5%) from the value determined using TRITC-containing vesicles. A two-component fit did not

Table 1. Diffusion Coefficients of Dye-Labeled and QD-Lipids in COS-7, 3T3, and HEK293

| probe | cell line | $D (\mu\text{m}^2 \text{s}^{-1})$ | reference |
|------------------------|-----------|-----------------------------------|-----------|
| diIC ₁₈ | COS-7 | 1.1 | 49 |
| diIC ₁₆ | COS-7 | 0.9 | 49 |
| QD-DHPTE | COS-7 | 1.07 | this work |
| Cy3-DOPE | HEK293 | 0.41 | 4 |
| QD-DHPTE | HEK293 | 0.70 | this work |
| diIC ₁₈ lcc | 3T3 | 0.68 | 50 |
| QD-DHPTE | 3T3 | 0.63 | this work |

improve the fit quality and resulted in almost identical diffusion times of $\tau_{D1} = 3247 \mu\text{s}$ and $\tau_{D2} = 3249 \mu\text{s}$, respectively. In contrast, with an excess of QDs, the FCS data are best described using a two-component model. Here the diffusion times of $\tau_{D1} = 1743 \mu\text{s}$ and $\tau_{D2} = 3428 \mu\text{s}$ are in good agreement with corresponding results obtained using free QDs and QDs bound to vesicles, respectively. Overall, these experiments provide




Figure 7. Representative trajectories of AEE-coated QDs bound to DHPTE in the plasma membrane of NRK cells determined at different frame rates of 30, 120, 400, and 1000 fps.

experimental evidence that samples of QD-labeled vesicles can be prepared, where free, excess QDs are statistically insignificant.

Upon verifying the proper conjugation of QDs to vesicles, the QD-containing SUVs were fused to the plasma membrane of cells. As already mentioned in the Experimental Section, QD-lipids were only tracked as long as they remained in the focus of the microscope, thus excluding tracking artifacts due to either unfused or internalized imaging probes. Figure 6 illustrates diffusion coefficient histograms of QD-conjugated lipids obtained from tracking experiments on COS7, 3T3, and HEK293 cells using a frame rate of 30 fps. Table 1 summarizes the average diffusion coefficients from these studies, together with previously reported data obtained using dye-labeled lipids. The good agreement between QD and dye data, particularly for COS-7 and 3T3 cells, indicates that individual QD-conjugated lipids were tracked on the cell surface. Only the HEK293 data show a moderate discrepancy. However, Murase and co-workers reported a large variation of diffusion coefficients among individual dye-labeled lipids tracked, resulting in an SD of $\pm 0.18 \mu\text{m}^2/\text{s}$.⁴ Overall, Figure 6 and Table 1 demonstrate that QDs conjugated to lipids maintain their inertness as tracking probes on the cell surface.

The main advantage of QD probes over their dye counterparts is expected to be seen in the case of fluorescence high-speed imaging. The higher brightness of the QD-lipids in comparison to dye-labeled lipids makes them particularly suited for faster than video rate fluorescence microscopy. Further, the use of organic dyes in high-speed imaging requires increased excitation laser powers, which decreases the lifetime of the organic dyes. In comparison, the photostability of the QD probes allows for long-time, high-speed single-particle imaging. Figure 7 presents typical trajectories from high-speed tracking experiments on the plasma membrane of NRK fibroblasts using QD-lipids with AEE coatings. These trajectories demonstrate the high photostability of QDs and the ability to conduct long-term tracking experi-

ments with these fluorescent probes. Figure 8 illustrates lateral diffusion and compartment size data from high-speed tracking experiments on AEE- (left) and LIPO-coated (right) QD-conjugated lipids in the plasma membrane of NRK fibroblasts. Here tracking was conducted using four different frame rates of 30, 120, 400, and 1000 fps. Regardless of the type of QD coating, the analysis shows that the membrane consists of two sizes of compartments, consistent with those observed previously with 40 nm diameter gold-labeled lipids.⁴⁸ As the frame rate increases, the measured diffusion coefficient of QD-lipids increases as expected, and the smaller compartments dominate in the analysis. For comparison, the diffusion coefficient determined with 40 nm diameter gold-labeled lipids viewed at 500 fps was $1.6 \mu\text{m}^2/\text{s}$,⁴⁷ which agrees well with this study of AEE and LIPO QD-lipids at 400 fps, where $0.85 \pm 0.15 \mu\text{m}^2/\text{s}$ was obtained. Overall, the tracking data in Figure 8 illustrate that AEE and LIPO QD-lipids show very similar diffusion properties on the cell surface. The current studies also demonstrated that fluorescence-based single-molecule tracking experiments can be conducted with frame rates of up to 1000 fps, which currently is the maximum frame rate of our imaging system. At 1000 fps, the signal-to-noise ratio, measured as the ratio of the background subtracted signal to the standard deviation in the fluctuations of the background subtracted signal, ranged from 2 to 2.5 on live cells. What is perhaps most important is that the photostability of the QD probes allows imaging at these high frame rates for over a second long. Such long trajectories will allow for quantitative analysis of structure

(47) Fujiwara, T.; Ritchie, K.; Murakoshi, H.; Jacobson, K.; Kusumi, A. *J. Cell Biol.* **2002**, *157*, 1071–1081.

(48) Ritchie, K.; Shan, X.-Y.; Kondo, J.; Iwasawa, K.; Fujawara, T. *Biophys. J.* **2005**, *88*, 2266–2277.

(49) Goodwin, J. S.; Drake, K. R.; Rimmert, C. L.; Kenworthy, A. K. *Biophys. J.* **2005**, *89*, 1398–1410.

(50) Metcalf, T. N.; Wang, J. L.; Schindler, M. *Proc. Natl. Acad. Sci. U.S.A.* **1986**, *83*, 95–99.




Figure 8. Frame-rate-dependent change in histograms of the diffusion coefficient and compartment size obtained using AEE- (left) and LIPO-coated QD-lipids (right) in the plasma membrane of NRK cells. The data illustrate the features of hop diffusion at high frame rates.

and interactions in the membrane. The access to higher frame rates is intriguing because it improves the spatial resolution of the imaging approach.⁴⁸ The improved spatial resolution is particularly advantageous to probe sub-diffraction-limit-size heterogeneities in cellular systems, which could previously be studied only using much larger Au labels.

Conclusion

Previous single-molecule tracking experiments on lipids have been limited by either a probe size, notably exceeding that of the labeled biomolecule (colloidal Au), or poor photostability (fluorescent dyes). In the current study, these limitations are overcome by conjugating CdSe/ZnS QDs with bioinert surface properties to lipids and tracking their lateral mobility in planar model membranes and in plasma membranes. Our studies show that highly photoluminescent, capped CdSe/ZnS with a size of less than 10 nm can be applied as powerful tracking probes in single-lipid tracking applications. The results presented herein highlight their particular strength in state-of-the-art high-speed single-molecule imaging. This experimental approach is very attractive for obtaining information about heterogeneities of biomolecules with a spatial resolution of less than 100 nm, which, so far, have only been detectable using 40–50 nm colloidal Au probes. Since molecular crowding is a common feature of many cellular regions, the smaller size QDs are

expected to outperform their colloidal Au counterparts in many imaging applications. QDs are also advantageous because, unlike Au probes, they can be employed in multicolor experiments where different types of biomolecules can be monitored at the same time. Although the current work has focused on QD-conjugated lipids, the presented technology is easily applicable to the long-term, high-speed tracking of individual proteins. For example, as will be described in a future paper, conjugation of QDs to proteins was easily accomplished by replacing maleimide functional groups on the QD surface with succinimidyl ester, which is reactive to amine groups of proteins (data not shown).

Acknowledgment. This research was supported in part by the National Science Foundation (grant MCB-0416779) and the National Institutes of Health (grant R21 DK77051-01).

Supporting Information Available: Detailed information about the methodology, including confocal fluorescence correlation spectroscopy, wide-field single-molecule fluorescence microscopy, oblique angle fluorescence microscopy, and single-molecule tracking analysis. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA803325B