


Conversion of Allylic Alcohols to Stereodefined Trisubstituted Alkenes: A Complementary Process to the Claisen Rearrangement

Justin K. Belardi, and Glenn C. Micalizio

J. Am. Chem. Soc., **2008**, 130 (50), 16870-16872 • DOI: 10.1021/ja8074242 • Publication Date (Web): 20 November 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



ACS Publications
 High quality. High impact.

Conversion of Allylic Alcohols to Stereodefined Trisubstituted Alkenes: A Complementary Process to the Claisen Rearrangement

Justin K. Belardi and Glenn C. Micalizio*

Department of Chemistry, Yale University, New Haven, Connecticut 06520–8107

Received September 19, 2008; E-mail: glenn.micalizio@yale.edu

Methods for the synthesis of geometrically defined trisubstituted olefins define a pillar of modern synthetic organic chemistry. From a target-based perspective, these stereodefined structural motifs are ubiquitous in natural products and molecules of biomedical and physical relevance (Figure 1). From a reactivity-based perspective,




Figure 1. Natural products possessing stereodefined (*Z*)-trisubstituted alkenes.

geometrically defined olefins serve as a foundation for stereoselective synthesis. These factors have driven the invention of a large variety of chemical methods for the convergent synthesis of stereodefined olefins. While many of these methods proceed from carbonyl addition chemistry or alkyne functionalization, the use of allylic alcohol derivatives in sigmatropic processes defines a powerful means to access a subset of stereodefined polysubstituted olefins.¹ Of these, Claisen-based methods have been particularly effective at establishing stereodefined (*E*)-trisubstituted olefins. Here, we describe a metal-mediated reductive cross-coupling reaction that defines a stereochemically complementary means of converting allylic alcohols to products related to those derived from Claisen rearrangement (Figure 2). While describing a unique stereoselective transformation for complex molecule synthesis, this study also defines a novel reductive cross-coupling reaction between alkenes and allylic alcohols.^{2,3}




Figure 2. A stereochemically unique method for the synthesis of stereodefined trisubstituted alkenes from allylic alcohols.

Recently, we demonstrated that allylic alcohols are useful substrates in titanium-mediated reductive cross-coupling reactions with internal alkynes.⁴ In these reactions, 1,4-dienes result from C–C bond formation between preformed titanium–alkyne complexes and allylic alkoxides. While quite useful for the stereoselective synthesis of substituted 1,4-dienes, we wondered whether a

related reductive cross-coupling process could define a stereoselective convergent pathway to isolated di- and trisubstituted olefins. To accomplish such a transformation, we targeted a reductive cross-coupling reaction between allylic alcohols and vinylsilanes.⁵




Figure 3. Preliminary study of stereoselection in reductive cross-coupling of allylic alcohols with vinylsilanes.

Our initial studies, depicted in Figure 3, provided some hope that the desired stereoselective transformation would be possible. In general, the preformed lithium alkoxide of an allylic alcohol was combined with vinyltrimethylsilane in Et₂O, cooled, and treated with the combination of ClTi(Oi-Pr)₃ and C₅H₉MgCl (–78 to 0 °C).⁶ While cross-coupling of allylic alkoxides **1** and **4** with vinyltrimethylsilane (**2**) provided cross-coupled products **3** and **5** in 58–66% yield, these reactions proceeded without stereoselection (*E/Z* = 1:1). In contrast, reductive cross-coupling of the (*Z*)-disubstituted alkene **6** with **2** provided the (*E*)-alkene **7** in 64% yield (*E/Z* = 9:1). Highest levels of (*E*)-selectivity were observed in the reaction of **8** with **2**. This process provided **9** in 69% yield with ≥20:1 selectivity; defining a stereoselective transformation that also establishes a quaternary center.⁷

While the cross-coupling reaction of terminally substituted allylic alcohols (i.e., **6** and **8**) delivers stereodefined (*E*)-disubstituted alkenes, the reaction of allylic alcohols bearing a 1,1-disubstituted olefin proceeds in a stereochemically unique manner. Reductive cross-coupling of **10** with **2** delivers **11** in 50% yield, with ≥20:1 selectivity, favoring the formation of the central stereodefined (*Z*)-trisubstituted alkene. Similarly, the coupling of the trisubstituted allylic alcohol **12** with **2** provides **13** in 65% yield (*Z:E* ≥ 20:1).

While preliminary studies investigating the coupling of simple acyclic- and cyclic alkenes with vinyltrimethylsilane indicate that this reaction is flexible and stereoselective (Figure 3 and Table 1, entries 1–3), we searched to identify a coupling partner that would allow for facile oxidation of the C–Si bond resident in the products. The combination of these two reactions, cross-coupling and oxidation, would then define a means to access stereodefined products related to those derived from Claisen rearrangement.¹

Table 1

entry	allylic alcohol	vinylsilane	yield (%)	(Z:E) dr	product
1 ^a			62	≥ 20:1	
2 ^a			71	≥ 20:1	
3 ^b			64	5:1	

Reaction conditions: (a) *n*-BuLi (1 equiv), vinylsilane (3 equiv), CITi(O*i*-Pr)₃ (3 equiv), C₅H₉MgCl (6 equiv) (−78 to 0 °C), then HCl (1 N); (b) *n*-BuLi (2 equiv), vinylsilane (3 equiv), CITi(O*i*-Pr)₃ (3 equiv), C₅H₉MgCl (6 equiv) (−78 to 0 °C), then HCl (1 N).

As illustrated in Figure 4, reductive cross-coupling of allylic alcohols **6** and **12** with vinyltrimethylchlorosilane (**20**) proceeds in a stereoselective manner, and delivers the corresponding silyl ethers **21** and **22** in 53% and 75% yield (*E/Z* = 10:1 to ≥ 20:1). Oxidation of the C–Si bond under standard conditions⁸ then delivers the stereodefined unsaturated primary carbinol (i.e., **22** → **23**). While products like **23** could be derived from **12** by the application of well-known Claisen rearrangement-based procedures, the cross-coupling reaction described here has the potential to deliver stereodefined products not readily accessible with these robust [3,3]-sigmatropic rearrangement processes. For example, Claisen rearrangement of **10**, followed by carbonyl reduction, provides the (*E*)-trisubstituted olefin **24** with high levels of stereoselection (*E/Z* ≥ 20:1).⁹ In this complementary process, reductive cross-coupling of **10** with vinyltrimethylchlorosilane (**20**), followed by oxidation, provides the isomeric (*Z*)-trisubstituted olefin **25** in 58% yield (*Z/E* ≥ 20:1).¹⁰

As illustrated in Table 2, this (*Z*)-selective reductive cross-coupling reaction is useful for the stereoselective functionalization of a variety of allylic alcohols (entries 1–5). Additionally, stereochemically defined products can be prepared from the coupling of mixtures of isomeric allylic alcohols (i.e., entries 6 and 7). Interestingly, coupling of **36** with **20** does not proceed in a




Figure 4. Cross-coupling reactions with vinyltrimethylchlorosilane.




Figure 5. A stereochemically complementary process with respect to the Claisen rearrangement. Reaction conditions: (a) Johnson *o*-ester Claisen rearrangement; (b) reduction;⁹ (c) **20**, CITi(O*i*-Pr)₃, *c*-C₅H₉MgCl, Et₂O (−78 to −50 °C), then cool to −78 °C and add lithium alkoxide of **10** (−78 to 0 °C) then, HCl (1 N) (75%, *Z/E* ≥ 20:1); (d) *t*-BuOOH, CsOH·H₂O, TBAF, DMF, 70 °C.

similarly stereoconvergent manner, indicating a potential role of the PMB ether in the stereochemical course of this reaction (entry 8).¹²

The regio- and stereochemical control observed in this allylic alcohol functionalization process is consistent with the empirical model depicted in Figure 6. In short, preassociation of the allylic alkoxide with a preformed titanacyclopropane (derived from the vinylsilane) produces an intermediate mixed titanate ester capable of rearrangement via formal metallo-[3,3]-rearrangement.⁴ While the C–C bond formation proceeds with allylic transposition, stereochemical control is thought to derive from minimization of

Table 2.

entry	allylic alcohol	vinylsilane	yield (%) ^a	(Z:E)	product
1			51	≥ 20:1	
2			58	≥ 20:1	
3			69	≥ 20:1	
4			56	≥ 20:1	
5			64	≥ 20:1	
6	34 ; R = TBS ¹¹	20	66	≥ 20:1	37 ; R = H
7	35 ; R = TBDPS ¹¹	20	61	≥ 20:1	37 ; R = H
8	36 ; R = PMB ¹¹	20	61	1.2:1 ¹²	38 ; R = PMB

^a Yield reported is over the two-step process: (1) Reductive cross-coupling (**20**, CITi(O*i*-Pr)₃, *c*-C₅H₉MgCl, Et₂O (−78 to −50 °C), then cool to −78 °C and add lithium alkoxide of the allylic alcohol (−78 to 0 °C) then, HCl (1 N)); (2) oxidation (*t*-BuOOH, CsOH·H₂O, TBAF, DMF, 70 °C).

nonbonded steric interactions in a boatlike conformation (i.e., **A'** and **B'**) where the σ_{C-Ti} bond is aligned with the $\pi_{C=C}$ bond.¹³




Figure 6. Model of stereoselection.

In sum, we have described a new regio- and stereoselective reductive cross-coupling reaction between allylic alcohols and vinylsilanes. This reaction proceeds with allylic transposition, delivers products with stereodefined di- and trisubstituted olefins, and provides a means to establish allylic tertiary and quaternary carbon centers. In addition to defining a novel olefin functionalization reaction and metal-mediated reductive cross-coupling process,¹⁴ this reaction provides a stereochemically unique pathway to functionalized acyclic products not readily accessible with modern [3,3]-sigmatropic rearrangement reactions.¹ Future study will explore both the utility of this process in target-oriented synthesis and the interplay between allylic alcohol substitution and selectivity.

Acknowledgment. We gratefully acknowledge financial support of this work by the American Cancer Society (Grant RSG-06-117-01), the Arnold and Mabel Beckman Foundation, Boehringer Ingelheim, Eli Lilly & Co., and the National Institutes of Health—NIGMS (Grant GM80266).

Note Added in Proof. During the course of our studies, a related process was published by Professor Jin K. Cha: Lysenko, I. L.; Kim, K.; Lee, H. G.; Cha, J. K. *J. Am. Chem. Soc.*, published online November 6, 2008 <http://dx.doi.org/10.1021/ja806440m>.

Supporting Information Available: Experimental procedures and tabulated spectroscopic data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For recent reviews, see: (a) Ziegler, F. E. *Chem. Rev.* **1988**, *88*, 1423–1452. (b) *The Claisen Rearrangement*: Hiersemann, M., Nubbemeyer, U., Eds.; Wiley-VCH: Weinheim, Germany, 2007, p 571. See also: (c) Johnson, W. S.; Werthem, A. L.; Bartlett, W. R.; Brocksom, T. J.; Li, T. T.; Faulkner, D. J.; Petersen, M. J. *J. Am. Chem. Soc.* **1970**, *92*, 741–743. (d) Perrin, C. L.; Faulkner, D. J. *Tetrahedron Lett.* **1969**, *10*, 2783–2786. (e) Funk, R. L.; Stallman, J. B.; Wos, J. A. *J. Am. Chem. Soc.* **1993**, *115*, 8847–8848 For Claisen rearrangement processes leading to (*Z*)-disubstituted olefins, see: (f) Nonoshita, K.; Banno, H.; Maruoka, K.; Yamamoto, H. *J. Am. Chem. Soc.* **1990**, *112*, 316–322. (g) Fernández de la Pradilla, R.; Montero, C.; Tortosa, M. *Org. Lett.* **2002**, *237*, 2–2376. (h) Takanami, T.; Hayashi, M.; Suda, K. *Tetrahedron Lett.* **2005**, *46*, 2893–2896. For a rare case of a [3,3]-rearrangement for the stereoselective synthesis of (*Z*)-trisubstituted olefins, see: (i) Krafft, M. E.; Dasse, O. A.; Jarrett, S.; Fievre, A. *J. Org. Chem.* **1995**, *60*, 5093–5101 For an example of a (*Z*)-selective [2,3]-Wittig rearrangement to access related products, see: (j) Still, W. C.; Mitra, A. *J. Am. Chem. Soc.* **1978**, *100*, 1927–1928.
- (2) For metal-mediated dimerization of terminal alkenes, see: (a) Isakov, V. E.; Kulinkovich, O. G. *Synlett* **2003**, 967–970. (b) Lee, J. C.; Sung, M. J.; Cha, J. K. *Tetrahedron Lett.* **2001**, *42*, 2059–2061. (c) Reference 5. For dimerization of, or coupling with, ethylene in route to metallacyclopentanes, see: (d) McDermott, J. X.; Wilson, M. E.; Whitesides, G. M. *J. Am. Chem. Soc.* **1976**, *98*, 6529–6536. (e) Grubbs, R. H.; Miyashita, A. *J. Chem. Soc., Chem. Commun.* **1977**, 864–865. (f) Cohen, S. A.; Auburn, P. R.; Bercaw, J. E. *J. Am. Chem. Soc.* **1983**, *105*, 1136–1143. (g) Mashima, K.; Takaya, H. *Organometallics* **1985**, *4*, 1464–1466. (h) Mashima, K.; Sakai, N.; Takaya, H. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 2475–2483. (i) Thorn, M. G.; Hill, J. E.; Waratuke, S. A.; Johnson, E. S.; Fanwick, P. E.; Rothwell, I. P. *J. Am. Chem. Soc.* **1997**, *119*, 8630–8641 For the dimerization of chalcones, see: (j) Schobert, R.; Maaref, R.; Dürr, S. *Synlett* **1995**, 83–84 For cross-coupling of 1,3-dienes with α -olefins, see: (k) Waratuke, S. A.; Thorn, M. G.; Fanwick, P. E.; Rothwell, A. P.; Rothwell, I. P. *J. Am. Chem. Soc.* **1999**, *121*, 9111–9119.
- (3) For titanium alkoxide-mediated coupling reactions of allylic alcohols and ethers with Grignard reagents, see: (a) Kulinkovich, O. G.; Epstein, O. L.; Isakov, V. E.; Khamel'nitskaya, I. A. *Synlett* **2001**, 49–52. (b) Matyushenkov, E. A.; Churikov, D. G.; Sokolov, N. A.; Kulinkovich, O. G. *Russian J. Org. Chem.* **2003**, *39*, 478–485 For Cp₂ZrCl₂-catalyzed carbomagnesiation of allylic alcohols and ethers, see: (c) Hoveyda, A. H.; Xu, Z. *J. Am. Chem. Soc.* **1991**, *113*, 5079–5080. (d) Hourri, A. F.; Didiuk, M. T.; Xu, Z.; Horan, N. R.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1993**, *115*, 6614–6624. (e) Hoveyda, A. H.; Morken, J. P. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1262–1284.
- (4) (a) Kolundzic, F.; Micalizio, G. C. *J. Am. Chem. Soc.* **2007**, *129*, 15112–15113 For reductive cross-coupling of allenic alcohols with alkynes via formal metallo-[3,3] rearrangement, see: (b) Shimp, H. L.; Hare, A.; McLaughlin, M.; Micalizio, G. C. *Tetrahedron* **2008**, *64*, 3437–3445 For reductive cross-coupling of allenic alcohols with aromatic imines via formal metallo-[3,3] rearrangement, see: (c) McLaughlin, M.; Shimp, H. L.; Navarro, R.; Micalizio, G. C. *Synlett* **2008**, 735–738.
- (5) For the first report documenting the ability of vinylsilanes to participate in the ligand exchange reaction with presumed alkoxytitanacyclopropane intermediates, see: Mizojiri, R.; Urabe, H.; Sato, F. *J. Org. Chem.* **2000**, *65*, 6217–6222.
- (6) For the use of cycloalkylmagnesium halide reagents to generate presumed alkoxytitanacyclopropane intermediates for intermolecular Kulinkovich cyclopropanation, see: Lee, J.; Kim, H.; Cha, J. K. *J. Am. Chem. Soc.* **1996**, *118*, 4198–4199.
- (7) For reviews on the construction of quaternary carbon centers, see: (a) Martin, S. F. *Tetrahedron* **1980**, *36*, 419–460. (b) Corey, E. J.; Buzman-Perez, A. *Angew. Chem., Int. Ed.* **1998**, *37*, 388–401. (c) Denissova, I.; Barriault, L. *Tetrahedron* **2003**, *59*, 10105–10146.
- (8) Smitrovich, J. H.; Woerpel, K. A. *J. Org. Chem.* **1996**, *61*, 6044–6046.
- (9) Abouabdellah, A.; Bonnet-Delpon, D. *Tetrahedron* **1994**, *50*, 11921–11932.
- (10) Titanium-mediated reductive cross-coupling reactions of allylic alcohols with internal alkynes proceed with a similar sense of stereoselection. See ref 4a for details.
- (11) Derived from the addition of 2-propenylmagnesium bromide to the corresponding chiral aldehyde. See Supporting Information for details.
- (12) The stereochemistry of the major/minor isomer was not determined. Future studies will examine the relationship of the relative stereochemistry of **36** on the stereochemical course of the reductive cross-coupling reaction.
- (13) This empirical model does not yet address the number of ligands present on the metal center in the transition state. Others have suggested–ate complexes as reactive intermediates in the Kulinkovich reaction: (a) Kulinkovich, O. G.; Kananovich, D. G. *Eur. J. Org. Chem.* **2007**, 212, 1–2132. (b) Kananovich, D. G.; Kulinkovich, O. G. *Tetrahedron* **2008**, *64*, 1536–1547 For the proposal of related intermediates in the reductive ethylation of allylic ethers, see: (c) Matyushenkov, E. A.; Churikov, D. G.; Sokolov, N. A.; Kulinkovich, O. G. *Russ. J. Org. Chem.* **2003**, *39*, 478–485.
- (14) While this reaction has not yet been rendered catalytic in the metal (Ti), the process provides a stereochemically unique transformation of great potential utility in organic synthesis. Like the Claisen rearrangement, these reactions proceed by substrate control, thereby eliminating the necessity to control stereochemistry by reagent- or catalyst-based methods. Finally, due to the low cost of the metal-containing reagents, and benign nature of the byproducts (TiO₂ and magnesium(II) salts), the reaction in its current form should be of great utility in organic chemistry.

JA8074242