

## Demonstration of a Peroxide Shunt in the Tetrahydropterin-Dependent Aromatic Amino Acid Monooxygenases


Jorge Alex Pavon and Paul F. Fitzpatrick\*

Departments of Biochemistry and Biophysics and of Chemistry, Texas A&M University, College Station, Texas 77843

Received January 7, 2009; E-mail: fitzpatrick@biochem.uthscsa.edu

Phenylalanine hydroxylase (PheH), tyrosine hydroxylase (TyrH), and tryptophan hydroxylase (TrpH) are nonheme iron monooxygenases that catalyze the insertion of an oxygen atom from O<sub>2</sub> into the aromatic side chain of their corresponding substrates using a tetrahydropterin (PH<sub>4</sub>) substrate as the reductant.<sup>1–3</sup> Their active sites each contain a mononuclear iron coordinated by two histidines and a glutamate,<sup>4–6</sup> an arrangement that has been termed a 2-his-1-carboxylate facial triad.<sup>7,8</sup> Scheme 1 shows the chemical mechanism proposed for PheH, TyrH, and TrpH.<sup>3</sup> The hydroxylating intermediate is an Fe(IV)O capable of aromatic, benzylic, and aliphatic hydroxylation.<sup>9,10</sup> This species has recently been detected in TyrH by freeze-quench Mössbauer spectroscopy.<sup>11</sup> The spectra and reactivity of the Fe(IV)O intermediate resemble those in members of the  $\alpha$ -ketoglutarate-dependent hydroxylase family, which also contain a mononuclear iron coordinated by a 2-his-1-carboxylate facial triad.<sup>12,13</sup>


### Scheme 1



In Scheme 1 the PH<sub>4</sub> supplies two electrons to reduce one atom of O<sub>2</sub> to the level of water, but it plays no role in the actual oxygen transfer to the amino acid substrate. This suggests that it could be possible to bypass the PH<sub>4</sub> and generate the Fe(IV)O intermediate directly with an alternative oxygen donor. Such a shunt has been possible in the cases of the heme-based cytochrome P450,<sup>14</sup> the binuclear nonheme methane monooxygenase,<sup>15</sup> and mononuclear nonheme dioxygenases,<sup>16,17</sup> but not with a mononuclear nonheme monooxygenase.

We now report that H<sub>2</sub>O<sub>2</sub> can replace PH<sub>4</sub> and O<sub>2</sub> to support amino acid hydroxylation by the aromatic amino acid hydroxylases. Incubation of PheH,<sup>18,19</sup> TyrH,<sup>20</sup> or TrpH<sup>19,21</sup> with phenylalanine and H<sub>2</sub>O<sub>2</sub> results in the formation of tyrosine and 3-HO-phenylalanine (Figure 1). No hydroxylated amino acids are detectable if apoenzyme is used. The rate of hydroxylation is unchanged when the reaction is carried out in the absence of O<sub>2</sub>. The ratio of tyrosine to 3-HO-phenylalanine produced is different for the three enzymes, with ratios of 1.5, 1.2, and 1.6 for PheH, TyrH, and TrpH, respectively. Controls showed that the PheH does not lose activity

under these conditions. No amino acid products could be detected with sodium periodate, cumene hydroperoxide, peracetic acid, or *tert*-butyl hydroperoxide instead of H<sub>2</sub>O<sub>2</sub>. The yield of hydroxylated amino acids was not affected by the radical quenchers mannitol or benzoate (10 mM). Addition of 2 mM 5-deaza-6-methyltetrahydropterin had no effects on the kinetics or the product distribution, suggesting that an open active site is required for the peroxide-dependent reaction.




**Figure 1.** Peroxide-dependent hydroxylation of phenylalanine by tyrosine hydroxylase. Solid line: TyrH (25  $\mu$ M) was incubated for 15 min at 30  $^{\circ}$ C with 10 mM H<sub>2</sub>O<sub>2</sub>, 400  $\mu$ M ferrous ammonium sulfate, 20 mM phenylalanine, and 100 mM NaCl in 150 mM Hepes buffer, pH 7.0. Dashed line: reaction with 50  $\mu$ M apo-TyrH. Dotted line: Reaction with 400  $\mu$ M ferrous ammonium sulfate but no enzyme. The products of the reaction were analyzed by reversed-phase HPLC, using a mobile phase of 15 mM sodium phosphate, pH 7.0, 1% tetrahydrofuran, with excitation at 270 nm and emission at 310 nm.

TyrH has previously been shown to produce tyrosine and 3-HO-phenylalanine from phenylalanine in PH<sub>4</sub>-dependent turnover,<sup>22</sup> but PheH and TrpH only produce tyrosine.<sup>18,21</sup> Thus, there is some loss of reaction specificity in the peroxide-dependent reaction, possibly due to the absence of the pterin. Tyrosine and tryptophan were also examined as substrates in the peroxide-dependent reactions for TyrH and TrpH, but the expected products 3,4-dihydroxyphenylalanine and 5-HO-tryptophan were not detected. Control reactions showed that these two compounds are not stable to the reaction conditions.

The kinetics of the H<sub>2</sub>O<sub>2</sub>-dependent reactions were examined for comparison with PH<sub>4</sub>-dependent turnover. The initial rate of phenylalanine hydroxylation was directly dependent on the concentration of enzyme (Figure 2). As also shown in Figure 2, the reaction continued for multiple turnovers. When the concentration of H<sub>2</sub>O<sub>2</sub> was varied, the initial rate of the reaction for all three enzymes showed saturation kinetics, with *K<sub>m</sub>* values of  $\sim$ 20 mM for each enzyme. In contrast, the initial rate of the reaction did not show evidence for saturation with phenylalanine at concentrations as high as 50 mM for any of the enzymes. The linear dependence of the rate on the concentration of phenylalanine yields *k<sub>cat</sub>*/*K<sub>phe</sub>* values of 6.4, 5.3, and 4.3 M<sup>-1</sup> min<sup>-1</sup> for the H<sub>2</sub>O<sub>2</sub>-dependent reaction for PheH, TyrH, and TrpH, respectively. These values are

5–6 orders of magnitude smaller than the corresponding values for  $\text{PH}_4$ -dependent turnover.<sup>18,21</sup> As shown in Scheme 1, formation of the  $\text{Fe(IV)O}$  species is proposed to require heterolytic cleavage of the O–O bond in an iron-peroxo-pterin intermediate, with a HO-pterin as the leaving group. Formation of the  $\text{Fe(IV)O}$  in the  $\text{H}_2\text{O}_2$ -dependent reaction would be expected to result from loss of water from an iron peroxide intermediate. The much slower reaction with  $\text{H}_2\text{O}_2$  can be attributed in part to the difference of about 4 units in the  $\text{p}K_a$  values of the leaving groups in the  $\text{H}_2\text{O}_2$ - and  $\text{PH}_4$ -dependent reactions. In addition, the similar kinetic parameters for all three enzymes in the  $\text{H}_2\text{O}_2$ -dependent reactions are consistent with the hydroxylating intermediates having similar reactivities for all three enzymes. The reactivities of the  $\text{Fe(IV)O}$  intermediates in  $\text{PH}_4$ -dependent turnover have previously been shown to be similar.<sup>23</sup>



**Figure 2.** Dependence of the amount of hydroxylated phenylalanine on the concentration of PheH for the peroxide-dependent reaction; p/e, moles of hydroxylated amino acid per mole of enzyme. Conditions same as those for Figure 1.

Active site mutants of TyrH and PheH that affect  $\text{PH}_4$ -dependent turnover were examined in the  $\text{H}_2\text{O}_2$ -dependent reaction. With E332A TyrH, only 2.5% of the reducing equivalents from 6Me $\text{PH}_4$  are used for productive turnover.<sup>20</sup> S395A TyrH forms the 4a-HO-pterin at a normal rate, but the hydroxylating intermediate breaks down unproductively so that only 1% is used to hydroxylate tyrosine.<sup>24</sup> V379D and F263A PheH have low turnover due to uncoupling of  $\text{PH}_4$  oxidation and amino acid hydroxylation.<sup>19,25</sup> With all four mutant enzymes, no tyrosine or 3-HO-phenylalanine could be detected in the  $\text{H}_2\text{O}_2$ -dependent reactions. Thus, amino acid residues required for proper reactivity of the  $\text{Fe(IV)O}$  intermediate in  $\text{PH}_4$ -dependent turnover are also required for  $\text{H}_2\text{O}_2$ -dependent turnover.

The aromatic amino acid hydroxylases have previously been shown to catalyze benzylic<sup>9,23</sup> and aliphatic hydroxylation.<sup>26</sup> To determine if the  $\text{H}_2\text{O}_2$ -dependent reaction is also capable of supporting these nonphysiological reactions, 4- $\text{CH}_3$ -phenylalanine and cyclohexylalanine were examined as substrates. In our hands TyrH and PheH catalyze the  $\text{PH}_4$ -dependent hydroxylation of cyclohexylalanine to form 4-HO-cyclohexylalanine with  $k_{\text{cat}}$  values

of 10 and 5  $\text{min}^{-1}$ , respectively, at 30 °C. Both enzymes also catalyze the same reaction using  $\text{H}_2\text{O}_2$ , with second-order rate constants of 0.17 and 0.28  $\text{M}^{-1} \text{min}^{-1}$ . With 4- $\text{CH}_3$ -phenylalanine as substrate for  $\text{PH}_4$ -dependent turnover, all three enzymes produce a combination of 4- $\text{CH}_2\text{OH}$ -, 3-HO,4- $\text{CH}_3$ -, and 4-HO,3- $\text{CH}_3$ -phenylalanine.<sup>23</sup> In the  $\text{H}_2\text{O}_2$ -dependent reactions, 4- $\text{CH}_2\text{OH}$ -phenylalanine is produced but the other two products could not be detected.

The present results establish that  $\text{H}_2\text{O}_2$  can replace  $\text{PH}_4$  and  $\text{O}_2$  to form the hydroxylating intermediate in the aromatic amino acid hydroxylases. The similar kinetic parameters for all three enzymes in the  $\text{H}_2\text{O}_2$ -dependent reactions are consistent with the hydroxylating intermediates having similar reactivity for all three enzymes. These results provide support for a hydroxylating intermediate such as  $\text{Fe(IV)O}$  that does not involve the pterin.

**Acknowledgment.** This work was supported in part by NIH grants R01 GM047291 (PFF) and F31 GM077092 (JAP) and Welch Foundation Grant A1245 (PFF). We thank Michaela Hyunh for technical assistance.

## References

- (1) Fitzpatrick, P. F. In *Advances in Enzymology and Related Areas of Molecular Biology*; Purich, D. L., Ed.; John Wiley & Sons, Inc.: 2000; Vol. 74, pp 235–294.
- (2) Kappock, T. J.; Caradonna, J. P. *Chem. Rev.* **1996**, *96*, 2659–2756.
- (3) Fitzpatrick, P. F. *Biochemistry* **2003**, *42*, 14083–14091.
- (4) Wang, L.; Erlandsen, H.; Haavik, J.; Knappskog, P. M.; Stevens, R. C. *Biochemistry* **2002**, *41*, 12569–12574.
- (5) Erlandsen, H.; Fusetti, F.; Martinez, A.; Hough, E.; Flatmark, T.; Stevens, R. C. *Nat. Struct. Biol.* **1997**, *4*, 995–1000.
- (6) Goodwill, K. E.; Sabatier, C.; Marks, C.; Raag, R.; Fitzpatrick, P. F.; Stevens, R. C. *Nat. Struct. Biol.* **1997**, *4*, 578–585.
- (7) Hegg, E. L.; Que, L. *Eur. J. Biochem.* **1997**, *250*, 625–629.
- (8) Que, L., Jr. *Nat. Struct. Biol.* **2000**, *7*, 182–184.
- (9) Hillas, P. J.; Fitzpatrick, P. F. *Biochemistry* **1996**, *35*, 6969–6975.
- (10) Moran, G. R.; Derecskei-Kovacs, A.; Hillas, P. J.; Fitzpatrick, P. F. *J. Am. Chem. Soc.* **2000**, *122*, 4535–4541.
- (11) Eser, B. E.; Barr, E. W.; Frantom, P. A.; Saleh, L.; Bollinger, J. M., Jr.; Krebs, C.; Fitzpatrick, P. F. *J. Am. Chem. Soc.* **2007**, *129*, 11334–11335.
- (12) Bollinger, J. M., Jr.; Krebs, C. *J. Inorg. Biochem.* **2006**, *100*, 586–605.
- (13) Krebs, C.; Galonic Fujimori, D.; Walsh, C. T.; Bollinger, J. M., Jr. *Acc. Chem. Res.* **2007**, *40*, 484–492.
- (14) Hrycay, E. G.; Gustafsson, J. A.; Ingelman-Sundberg, M.; Ernster, L. *Biochem. Biophys. Res. Commun.* **1975**, *66*, 209–216.
- (15) Froland, W. A.; Andersson, K. K.; Lee, S. K.; Liu, Y.; Lipscomb, J. D. *J. Biol. Chem.* **1992**, *267*, 17588–17597.
- (16) Neibergall, M. B.; Stubna, A.; Mekmouche, Y.; Munck, E.; Lipscomb, J. D. *Biochemistry* **2007**, *46*, 8004–8016.
- (17) Wolfe, M. D.; Lipscomb, J. D. *J. Biol. Chem.* **2003**, *278*, 829–835.
- (18) Daubner, S. C.; Hillas, P. J.; Fitzpatrick, P. F. *Biochemistry* **1997**, *36*, 11574–11582.
- (19) Pavon, J. A.; Fitzpatrick, P. F. *Biochemistry* **2006**, *45*, 11030–11037.
- (20) Daubner, S. C.; Fitzpatrick, P. F. *Biochemistry* **1999**, *38*, 4448–4454.
- (21) Moran, G. R.; Daubner, S. C.; Fitzpatrick, P. F. *J. Biol. Chem.* **1998**, *273*, 12259–12266.
- (22) Fitzpatrick, P. F. *J. Am. Chem. Soc.* **1994**, *116*, 1133–1134.
- (23) Pavon, J. A.; Fitzpatrick, P. F. *J. Am. Chem. Soc.* **2005**, *127*, 16414–16415.
- (24) Ellis, H. R.; Daubner, S. C.; Fitzpatrick, P. F. *Biochemistry* **2000**, *39*, 4174–4181.
- (25) Daubner, S. C.; Melendez, J.; Fitzpatrick, P. F. *Biochemistry* **2000**, *39*, 9652–9661.
- (26) Carr, R. T.; Balasubramanian, S.; Hawkins, P. C. D.; Benkovic, S. J. *Biochemistry* **1995**, *34*, 7525–7532.

JA900128M