

Characterization of Structurally Unusual Diiron N_xH_y Complexes

Caroline T. Saouma, Peter Müller, and Jonas C. Peters*


Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Received May 15, 2009; E-mail: jcpeters@mit.edu

Mechanistic proposals concerning the pathway of N_2 reduction in biology at the MoFe-cofactor of nitrogenase continue to be advanced.¹ In addition to nitrogen, hydrazine² and diazene^{1a} are nitrogenase substrates, and recent DFT calculations and spectroscopic studies suggest that whereas initial N_2 binding may occur at one iron center, diiron pathways may be involved at certain N_xH_y intermediate stages en route to ammonia formation.^{1a,3} In this broad context, recent work has explored the synthesis and spectroscopic characterization of structurally unusual mono- and bimetallic iron complexes featuring nitrogenous ligand functionalities.^{4,5} The demand for such model complexes continues in light of recent ENDOR and ESEEM spectroscopic data that has been obtained under turnover conditions at the cofactor.^{1a,2,6} To date, there are few synthetic iron systems that feature parent hydrazine (N_2H_4),⁷ hydrazido ($N_2H_2^{2-}$), diazene (N_2H_2),⁸ amide (NH_2^-),⁹ and imide (NH^{2-})^{4d} ligands. Herein we describe the synthesis and characterization of a series of structurally distinct diiron complexes that feature each of these ligand types.

Entry into the N_xH_y chemistry of present interest is realized with the iron(II) alkyl precursors $[PhBP^R_3]FeMe$ ($[PhBP^R_3] = PhB(CH_2PR_2)_3$; R = Ph, CH_2Cy). These high-spin complexes are prepared either by addition of excess Me_2Mg to a benzene solution of $[PhBP^{Ph}_3]FeCl$ or by addition of $MeLi$ to a thawing THF solution of $[PhBP^{CH_2Cy}_3]FeCl$. The room-temperature addition of 1 equiv of hydrazine to yellow THF solutions of either $[PhBP^{Ph}_3]FeMe$ (**1**) or $[PhBP^{CH_2Cy}_3]FeMe$ (**2**) results in the immediate release of methane and clean conversion to red and diamagnetic $\{(PhBP^{Ph}_3)Fe\}_2(\mu-\eta^1:\eta^1-N_2H_4)(\mu-\eta^2:\eta^2-N_2H_2)$ (**3**) or purple and diamagnetic $\{(PhBP^{CH_2Cy}_3)Fe\}_2(\mu-\eta^1:\eta^1-N_2H_4)(\mu-\eta^2:\eta^2-N_2H_2)$ (**4**), respectively (Scheme 1). The solid-state structures of the diiron cores of **3** and **4** are nearly identical [see the Supporting Information (SI)]; that of **3** is shown in Figure 1. Its two iron centers are bridged by N_2H_4 and $N_2H_2^{2-}$ ligands. We assign the end-on bridging ligand as N_2H_4 and the side-bonded bridging ligand as $N_2H_2^{2-}$.¹⁰ The N3–N4 bond distance of 1.429(3) Å in **3** is consistent with an N–N single bond, and hence, this ligand is best assigned as $N_2H_2^{2-}$ rather than $HN=NH$. The 1H NMR spectrum (THF- d_8) of **3** reveals a solution structure similar to that observed in the solid state. A N_2H_4 resonance is noted at 2.54 ppm and a $N_2H_2^{2-}$ resonance at 2.59 ppm. Each of these peaks is split into an apparent doublet when **3** is prepared

Scheme 1



using $^{15}N_2H_4$ ($^1J_{NH} = 59.5$ Hz for $N_2H_2^{2-}$; $^1J_{NH} = 72.0$ Hz for N_2H_4). The broadness of the doublets precludes the resolution of higher-order N–H and H–H coupling.¹¹ The corresponding ^{15}N NMR spectrum shows a doublet at -10.0 ppm for $N_2H_2^{2-}$ and a triplet at 58.37 ppm for N_2H_4 ,¹² both of which collapse into singlets upon 1H decoupling. Complex **4** displays similar ^{15}N NMR characteristics ($N_2H_2^{2-}$: 0.86 ppm, doublet, $J = 45.2$ Hz; N_2H_4 : 50.86 ppm, triplet, $J = 68.9$ Hz).

Whereas Sellmann previously reported a few examples of diiron $\mu-\eta^1:\eta^1-HN=NH$ species,⁸ the bimetallic iron cores of **3** and **4** are unique. Schrock has characterized a ditungsten $(\mu-\eta^1:\eta^1-N_2H_4)(\mu-\eta^2:\eta^2-N_2H_2)$ complex whose core is closely related to those in **3** and **4**.¹³

Whereas solutions of **3** are stable for days at 60 °C, purple solutions of **4** are thermally unstable even at 22 °C ($t_{1/2} = 4$ h) and decay quantitatively to the isolable green diamagnetic product $\{(PhBP^{CH_2Cy}_3)Fe\}_2(\mu-\eta^1:\eta^1-N_2H_2)(\mu-NH_2)_2$ (**5**). Signature NMR data for **5** are as follows: Its 1H NMR spectrum (THF- d_8) features three broad singlets at 16.21, -1.65 , and -3.85 ppm, corresponding to bound $HN=NH$ and diastereotopic NH_2^- protons, respectively. Each of these 1H NMR resonances is split into a doublet upon ^{15}N labeling. Its ^{15}N NMR spectrum contains a doublet at 419.1 ppm ($^1J_{NH} = 65.1$ Hz) that collapses to a singlet upon decoupling of the proton resonance at 16.21 ppm. The NH_2^- nitrogens are observed as a triplet at -58.1 ppm ($J = 59.5$ Hz).

The solid-state structure of **5** is shown in Figure 2. Both the Fe–N1 bond distance of 1.882(8) Å and the N1–N1' bond distance of 1.283(15) Å are similar to those found in Sellmann's $Fe_2(\mu-\eta^1:\eta^1-HN=NH)$ complex.⁸ In addition, the Fe–N bond-distance contraction of 0.17 Å that occurs upon oxidation of hydrazine to diazene is also consistent with other structurally characterized diazene complexes^{8,14} and is likely due to a combination of the smaller covalent radii of sp^2 -hybridized nitrogen relative to sp^3 -hybridized nitrogen and modest back-bonding. Notable in the structure of **5** is the presence of *cis*-diazene. Whereas monosub-




Figure 1. Solid-state molecular structures of the core atoms of **3** and **6**. See the SI for complete structures and details.

stituted diazenes (HN=NR) are known to bind transition metals in a *cis* conformation,¹⁵ to our knowledge all known complexes of HN=NH show *trans* ligation. Hence, **5** appears to be structurally distinct in this context.⁸

The addition of 2.2 equiv of Pb(OAc)₄ to red **3** leads to oxidation of the bound N₂H₄¹⁶ to afford green and diamagnetic {[PhBP^{Ph}]₃-Fe}₂(μ-η¹:η¹-N₂H₂)(μ-η²:η²-N₂H₂) (**6**) as the major product along with release of acetic acid. The solid-state structure of **6** is shown in Figure 1. Most striking is the presence of both an η¹:η¹-bridging *cis*-HN=NH ligand and a μ-η²:η² HN–NH²⁻ ligand. To our knowledge, **6** is the only transition-metal complex featuring the N₂H₂ ligand in each of its limiting states of oxidation. The N1–N2 bond distance of 1.281(5) Å for the ligand assigned as HN=NH is similar to that found for the bridging HN=NH ligand of **5**. The N3–N4 bond distance of 1.458(5) Å for the ligand assigned as N₂H₂²⁻ is slightly elongated compared with that of the N₂H₂²⁻ ligand of **3**. The average Fe–N_{diazene} bond distance of 1.89 Å is appreciably shorter than the average Fe–N_{hydrazido}²⁻ bond distance of 1.99 Å.

As was observed for **3** and **4**, the structure of **6** is preserved in solution. In the ¹H NMR spectrum (C₆D₆) of **6**, broad singlets at 13.20 and 4.16 ppm are assigned to the HN=NH and N₂H₂²⁻ protons, respectively; both of these peaks are split into apparent broad doublets when samples of **6** are prepared using ¹⁵N-enriched **6**. The ¹⁵N NMR spectrum of **6** contains a doublet at 58.0 ppm (¹J_{NH} = 58.1 Hz) for the N₂H₂²⁻ nitrogens and a doublet of doublets at 407.5 ppm (¹J_{NH} = 69.2 Hz, *J* ≈ 20 Hz) corresponding to the HN=NH ligand.

Complex **6** can further be oxidized with *p*-benzoquinone to cleanly generate a new diamagnetic species, {[PhBP^{Ph}]₃Fe}₂(μ-NH)₂ (**7**), along with *p*-hydrobenzoquinone as a byproduct. In order to ascertain the fate of the bound diazene, the reaction was analyzed with a Toepler pump apparatus, and the evolution of 1 equiv of N₂ was confirmed. This complex can alternatively be prepared by addition of 5 equiv of *p*-benzoquinone to **3**. The ³¹P NMR spectrum of **7** displays a single resonance at 32.5 ppm at 22 °C. The ¹H NMR spectrum shows a single set of [PhBP^{Ph}]₃ resonances, with an additional singlet at 25.3 ppm that integrates to one proton per [PhBP^{Ph}]₃; when **7** is prepared using isotopically enriched **3**, the singlet at 25.04 ppm is split into a doublet (¹J_{NH} = 64.0 Hz). The ¹⁵N NMR spectrum displays a doublet at 563.5 ppm. On the basis of these NMR data as well as IR data (see the SI), the solution structure of **7** is assigned as diferric {[PhBP^{Ph}]₃Fe}₂(μ-NH)₂. The spectroscopic data for the bridging NH ligand in **7** is similar to that previously reported for {[PhBP^{Ph}]₃Fe}₂(μ-NH)(μ-H)}{Na}.^{4d}




Figure 2. Solid-state molecular structures of the core atoms of **5** and **7**. See the SI for complete structures and details.

Crystals of **7** can be grown from a THF/cyclopentane solution, and its solid-state structure is shown in Figure 2. The bridging imides are disordered over three positions (see the SI) and modeled satisfactorily with a population ratio of 0.93:0.75:0.32. The average Fe–N distance

of 1.84 Å is ~0.17 Å longer than that found in {[PhBP^{Ph}]₃Fe}₂(μ-NH)(μ-H)}{Na}.^{4d} The average Fe–P distance of 2.33 Å is ~0.075 Å longer than those found in **3** and **6**, consistent with its assignment as an antiferromagnetically coupled diiron(III) complex.

In conclusion, we have characterized a series of structurally fascinating diiron N_xH_y species that contain hydrazine, hydrazido, and *cis*-diazene bridges. Thermal and oxidative transformations also lead to unusual examples of diiron species featuring μ-NH₂ and μ-NH ligands. In certain instances, low-temperature experiments revealed the presence of intermediate species. Ongoing work concerns detailed vibrational characterization of the species described herein, in addition to mechanistic studies.

Acknowledgment. We acknowledge the NIH (GM-070757). Funding for the MIT Department of Chemistry Instrumentation Facility has been provided in part by the NSF (CHE-0234877). C.T.S. is grateful for an NSF graduate fellowship.

Supporting Information Available: Detailed experimental procedures and characterization data for **1–7** and crystallographic details for **3–7** (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Hoffmann, B. M.; Dean, D. R.; Seefeldt, L. C. *Acc. Chem. Res.* **2009**, *42*, 609. (b) Peters, J. C.; Mehn, M. P. In *Activation of Small Molecules*; Tolman, W. B., Ed.; Wiley: New York, 2006; pp 81–120. (c) Howard, J. B.; Rees, D. C. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 17088. (d) Schrock, R. R. *Angew. Chem., Int. Ed.* **2008**, *47*, 5512.
- (2) Barney, B. M.; Yang, T.-C.; Igarashi, R. Y.; Santos, P. C.; Laryukhin, M.; Lee, H.-I.; Hoffman, B. M.; Deadn, D. D.; Seefeldt, L. C. *J. Am. Chem. Soc.* **2005**, *127*, 14960.
- (3) Kästner, J.; Blöchl, P. E. *J. Am. Chem. Soc.* **2007**, *129*, 2998.
- (4) For example, see: (a) Hendrich, M. P.; Gunderson, W.; Behan, R. K.; Green, M. T.; Mehn, M. P.; Betley, T. A.; Lu, C. C.; Peters, J. C. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 17107. (b) Smith, J. M.; Lachicotte, R. J.; Pittard, K. A.; Cundari, T. R.; Lukat-Rodgers, G.; Rodgers, K. R.; Holland, P. L. *J. Am. Chem. Soc.* **2001**, *123*, 9222. (c) Betley, T. A.; Peters, J. C. *J. Am. Chem. Soc.* **2003**, *125*, 10782. (d) Brown, S. D.; Mehn, M. P.; Peters, J. C. *J. Am. Chem. Soc.* **2005**, *127*, 13146. (e) Smith, J. M.; Sadique, A. R.; Cundari, T. R.; Rodgers, K. R.; Lukat-Rodgers, G.; Lachicotte, R. J.; Flaschenriem, C. J.; Vela, J.; Holland, P. L. *J. Am. Chem. Soc.* **2006**, *128*, 756.
- (5) (a) Vogel, C.; Heinemann, F. W.; Sutter, J.; Anthon, C.; Meyer, K. *Angew. Chem., Int. Ed.* **2008**, *47*, 2681. (b) Scepaniak, J. J.; Young, J. A.; Bontchev, R. P.; Smith, J. M. *Angew. Chem., Int. Ed.* **2009**, *48*, 3158.
- (6) Dean, D. R.; Hoffman, B. M.; Seefeldt, L. C. *Biochemistry* **2007**, *46*, 6784.
- (7) (a) Goedken, V. L.; Peng, S.-M.; Molin-Norris, J.; Park, Y.-a. *J. Am. Chem. Soc.* **1976**, *98*, 8391. (b) Sellmann, D.; Shaban, S. Y.; Heinemann, F. W. *Eur. J. Inorg. Chem.* **2004**, 4591. (c) Crossland, J. L.; Zakharov, L. N.; Tyler, D. R. *Inorg. Chem.* **2007**, *46*, 10476. (d) Field, L. D.; Li, H. L.; Dalgarno, S. J.; Turner, P. *Chem. Commun.* **2008**, 1680.
- (8) For a diiron η¹:η¹-HN=NH complex, see: Sellmann, D.; Soglowek, W.; Knoch, F.; Moll, M. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 1271. For a recent example of a monometallic Fe(η²-N₂H₂) complex, see ref 7d.
- (9) (a) For a terminal Fe–NH₂ complex, see: Fox, D. J.; Bergman, R. G. *J. Am. Chem. Soc.* **2003**, *125*, 8985. (b) For bridging Fe₂(NH₂), see: Dahl, L. F.; Costello, W. R.; King, R. B. *J. Am. Chem. Soc.* **1968**, *90*, 5422.
- (10) For all of the structures presented herein, the protons on the N_xH_y ligands were located in the difference Fourier map and refined semi-freely with the help of restraints on the N–H distances and the Fe–N–H angles while constraining the U_{iso} value of each H atom to –1.2 times the U_{eq} value of the N atom connected to that proton.
- (11) The magnitude of these couplings may be expected to be on the order of 0–10 Hz. See: (a) von Philipsborn, W.; Müller, R. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 383. (b) Smith, M. R., III; Cheng, T. Y.; Hillhouse, G. L. *J. Am. Chem. Soc.* **1993**, *115*, 8638.
- (12) All of the ¹⁵N NMR chemical shifts are reported relative to liquid NH₃.
- (13) Blum, L.; Williams, I. D.; Schrock, R. R. *J. Am. Chem. Soc.* **1984**, *106*, 8316.
- (14) Cheng, T. Y.; Ponce, A.; Rheingold, A. L.; Hillhouse, G. L. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 657.
- (15) (a) Albertin, G.; Antoniutti, S.; Lanfranchi, M.; Pelizzi, G.; Bordignon, E. *Inorg. Chem.* **1986**, *25*, 950. (b) Smith, M. R., III; Hillhouse, G. L. *J. Am. Chem. Soc.* **1988**, *110*, 4066. (c) Chen, Y.; Zhou, Y.; Chen, P.; Tao, Y.; Li, Y.; Qu, J. *J. Am. Chem. Soc.* **2008**, *130*, 15250.
- (16) Smith, M. R., III; Cheng, T. Y.; Hillhouse, G. L. *J. Am. Chem. Soc.* **1993**, *115*, 8638.

JA903967Z