

The Reaction Mechanism of the Enantioselective Tsuji Allylation: Inner-Sphere and Outer-Sphere Pathways, Internal Rearrangements, and Asymmetric C–C Bond Formation


John A. Keith,[†] Douglas C. Behenna,[‡] Nathaniel Sherden,[‡] Justin T. Mohr,[‡] Sandy Ma,[‡] Smaranda C. Marinescu,[‡] Robert J. Nielsen,[†] Jonas Oxgaard,[†] Brian M. Stoltz,^{*,‡} and William A. Goddard, III^{*,†}

[†]Materials and Process Simulation Center, Beckman Institute California Institute of Technology, Pasadena, California 91125, United States

[‡]The Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, United States

Supporting Information

ABSTRACT: We use first principles quantum mechanics (density functional theory) to report a detailed reaction mechanism of the asymmetric Tsuji allylation involving prochiral nucleophiles and nonprochiral allyl fragments, which is consistent with experimental findings. The observed enantioselectivity is best explained with an inner-sphere mechanism involving the formation of a 5-coordinate Pd species that undergoes a ligand rearrangement, which is selective with regard to the prochiral faces of the intermediate enolate. Subsequent reductive elimination generates the product and a Pd⁰ complex. The reductive elimination occurs via an unconventional seven-centered transition state that contrasts dramatically with the standard three-centered C–C reductive elimination mechanism. Although limitations in the present theory prevent the conclusive identification of the enantioselective step, we note that three different computational schemes using different levels of theory all find that inner-sphere pathways are lower in energy than outer-sphere pathways. This result qualitatively contrasts with established allylation reaction mechanisms involving prochiral nucleophiles and prochiral allyl fragments. Energetic profiles of all reaction pathways are presented in detail.




INTRODUCTION

The importance of homogeneous asymmetric catalysis increases as the world's need for efficient access to enantioenriched molecules increases.^{1–4} In particular, the synthesis of compounds containing enantioenriched quaternary stereocenters by catalytic methods remains a challenging problem. On the basis of an enantioselective version of the Tsuji decarboxylation allylation reaction (Scheme 1)⁹ we recently reported several related methods to prepare quaternary stereocenters of high ee.^{5–8} In a representative case, allyl enol carbonate **1** undergoes oxidative addition with a Pd⁰ complex and decarboxylation. (For a recent review on decarboxylative

allylic alkylations and their asymmetric variants, see ref 10.) The resulting Pd²⁺ π -allyl complex then reacts with the *in situ*-generated enolate, forming α -allylated ketone **2**. Before our work, the majority of asymmetric enolate allylations focused on reactions of prochiral allyl fragments with malonate nucleophiles,^{11–13} or on prochiral ketones that had a single α -site or whose α -sites had large differences in acidity (e.g., Trost^{12,14–17}). Related to our work, Helmchen¹⁸ and Pfaltz⁴ also used malonates with catalyst ligands similar to those we employed.

Mechanistic studies of these prior asymmetric processes involving more stabilized enolates have established that they proceed via outer-sphere nucleophilic attacks.¹⁹ In more recent work, Trost has shown evidence that reactions of unstabilized enolates catalyzed by his bisphosphine ligands may proceed by either outer-sphere or inner-sphere mechanisms.^{20–23} Lloyd-Jones and Norrby have suggested that hydrogen bonding interactions between the enolate and ligand backbone in Trost's systems can lead to a subtle distinction where an inner-sphere reaction occurs with respect to the complex itself but not with respect to the metal center.²⁴ Insightful mechanistic studies

Scheme 1. Enantioselective Tsuji Allylation



Received: July 19, 2012

Published: October 28, 2012

investigating other allylation mechanisms must explicitly consider the feasibility of different modes of nucleophilic attack. Harder and more basic nucleophiles are known to attack at the metal center before C–C bond formation occurs in a number of processes.^{25–29} A collaboration between our groups determined that the reaction we reported likely reacts via an inner-sphere mechanism that terminates with a nontraditional reductive elimination.³⁰ The present work reports considerably more detail on this mechanism.

Our original experimental study reported the formation of asymmetric quaternary stereocenters from unstabilized prochiral ketone enolate nucleophiles and allyl electrophiles.⁵ The 2-allyl-2-alkyl cyclohexanone products formed are distinct from the vast majority of asymmetric allylic alkylation products. In contrast to previous works that utilized *prochiral* electrophiles,^{4,9,12,14–16,18,19,31–35} we demonstrated enantioselective Tsuji allylation using *nonprochiral* electrophiles and prochiral ketone enolate nucleophiles. This permits the formation of quaternary centers from ketone enolates that have two similarly acidic sites alpha to the ketone. These desirable products are formed in high yields and high enantioselectivities. This process has found successful application in the total synthesis of many natural products to date,^{36–45} as well as in the development of related enantioselective decarboxylative protonation^{46,47} and sequential conjugate addition allylation reactions.⁴⁸ The standard ligand used in these transformations is (*S*)-*t*-Bu-phosphinoxazoline (PHOX)^{49,50} (Scheme 1), which provided variants of ketone **2** with yields from 80 to 99% and ee values ranging from 79 to 92%. Although this level of enantioselectivity is useful for many applications, ee values consistently above 95% would make the process more appealing in industrial setting, and so we sought to understand the fine details of the reaction's mechanism.

Well-established stereochemical probes involving cyclohexenyl electrophiles^{26,27} were employed in attempts to determine the reaction mechanism. However, the extremely low yields, dr and ee values observed in all such attempts indicated that these cyclic substrates with our catalyst likely proceed by a substantially different mechanism from our standard high ee substrates. Extensive efforts to develop other experimental tests to elucidate this mechanism were not successful; thus, we turned to quantum chemistry simulations to investigate the details of the mechanism.

The observation that this reaction is effective using enol carbonates, silyl enol ethers, and β -keto allyl esters^{5,6,51} indicated to us that they likely share a common intermediate. Because most substrates performed equally well in polar solvents such as THF and nonpolar solvents such as benzene, the reaction mechanism appeared not strongly dependent on electrostatic solvation effects. However, we did find that a low solvent dielectric (in the 2–8 range) was important for obtaining high levels of enantioselectivity. We obtained an X-ray crystal structure of a $\text{Pd}(\text{allyl})^+\text{PF}_6^-$ complex, **3** (see Figure 1), that was similar to other examples in the literature^{52–54} and found that it was a competent catalyst for reactions of silyl enol ethers, suggesting that it is an intermediate in the reaction. Two experiments within our group³⁰ suggested the reaction involved only one $\text{Pd}(\text{PHOX})$ complex: (1) experiments varying the ee of the catalyst from 0 to 100% showed no nonlinear effects in the ee of the product formed, and (2) simple kinetic studies showed that the reaction was roughly zero-order in $\text{Pd}(\text{PHOX})$ and starting material.





Figure 1. Superimposed calculated (yellow) and experimentally obtained solid-state (silver) structure for π -allyl complex **3**. The experimental crystal structure was obtained using PF_6^- as the counterion.

Given the current understanding of the Tsuji allylation mechanism and the information stated above, we investigated the enantioselective Tsuji allylation mechanism using first principles quantum mechanics including effects of solvation (in this study, the reaction was modeled in THF). Given the complexity of this mechanism, we focused on the steps following decarboxylation that begin with the association of the enolate and π -allyl complex **3** and report the reaction pathways leading to the enantioenriched product (*S*)-**2**. We also focused calculations using only the PHOX ligand. Our calculations identify an inner-sphere mechanism consistent with current experimental observations and shed light on the enantiodetermining step for this reaction. A key component to this mechanism is a pericyclic reductive rearrangement process that is distinct from the traditional three-centered process seen commonly in other C–C bond-forming reactions mentioned previously. On the basis of the mechanism and the electronic structures of the intermediates and transition states, we suggest strategies for optimizing the catalyst and the reaction conditions for the process (Scheme 2).

Scheme 2. Portion of the Reaction Mechanism Studied in This Work



COMPUTATIONAL METHODS

Density functional theory (DFT) energy calculations, geometry optimizations, and vibrational frequencies were run using Jaguar version 7.0.⁵⁵ Reduced model systems for the PHOX backbone did not adequately describe steric effects; as a result, all calculations involved the full PHOX ligand. Molecular geometries were obtained with the B3LYP hybrid exchange correlation functional,⁵⁶ which uses the Becke generalized gradient approximation⁵⁷ (GGA) in conjunction with a component of exact Hartree–Fock exchange (hybrid DFT) with the Lee, Yang, and Parr correlation functional.⁵⁸ This approach

has been shown quite effective in elucidating important aspects of organometallic reaction mechanisms.^{59–62}

Preliminary geometry optimizations were run with a mixed-basis set approach, henceforth denoted as BS1. In this method, the Pd LACVP electronic core potential^{63,64} (ECP) including 18 explicit electrons was used; the 6-31G* basis set was used on N, P, and the six atoms in the allyl- and enol- fragments; the MIDI! basis set⁶⁵ was used on all other atoms. Vibrational frequencies and their thermodynamic contributions from B3LYP/BS1 were negligibly different from values obtained using the same ECP and with the 6-31G** basis set on all other atoms (henceforth denoted as BS2). We report B3LYP/BS2 geometries.

The correct number of imaginary frequencies confirmed all stationary points. Starting from an optimized transition state, geometry optimizations starting from positions nudged along the imaginary vibrational mode confirmed that the transition states correctly linked stable intermediates along a mechanistic pathway. Thermodynamic energies were added to estimate the free energies and enthalpies at reaction temperature by calculating vibrational frequencies for all intermediates and transition states. Only the vibrational partition function was included for these energies to avoid uncertainties involving translations and rotations for species in aqueous solution. To eschew inconsistent thermal energies due to spuriously low vibrational frequencies, zero-point and thermal energies were calculated after substituting all vibrational frequencies <50 cm⁻¹ with values of 50 cm⁻¹.

Single-point energy calculations of triple- ζ quality (LACV3P, a decontracted version of LACVP, on Pd with 6-311++G** on all other atoms, henceforth denoted as BS3) were calculated and compared to other single-point energies from the hybrid *mpW1PW91*⁶⁶ exchange correlation functional. Calculations include electronic relaxations due to THF solvation (basis set = (LACV3P**, 6-311G**), probe radius = 2.527 Å, $\epsilon = 7.52$) from single-point self-consistent Poisson–Boltzmann implicit solvation calculations.^{67,68} Unless explicitly noted, all reported energies in the text are ΔG_{298} values from B3LYP/BS3//B3LYP/BS2 calculations including solvation corrections. Lastly, single-point dispersion corrections intended to correct deficiencies in B3LYP when calculating nonbonding interactions were included with single-point energy corrections using Grimme's DFT-D3 method⁶⁹ as calculated with GAMESS-US.^{70–72} Overall, assuming the appropriate physics of the chemical reactions is being modeled correctly, our calculations should be accurate to experiment within 2–3 kcal/mol for thermodynamic energies and 2–5 kcal/mol for barrier heights. Differences in barrier heights (which benefit from error cancellation) can be considered to be more accurate (to 1–2 kcal/mol). Molecular illustrations were made from VMD.⁷³

RESULTS AND DISCUSSION

We calculate that the reaction thermodynamics for enol-allyl carbonate **1** converting to (*S*)-2-allyl-2-methyl cyclohexanone, (*S*)-**2**, is exoergic by $\Delta G_{298} = -36.6$ kcal/mol in gas phase and -34.9 kcal/mol in THF. Of this energy, -15.2 kcal/mol comes from releasing CO₂. Experimental thermochemical reaction data is not available; however, the gas phase value corresponds well with those calculated from additivity tables (-35 kcal).⁷⁴

The experimental isolation of π -allyl complex **3**•PF₆ (Figure 1) and its demonstration as a competent catalyst motivated us to start our mechanistic investigation at the point of an ion pair involving the cationic π -allyl intermediate and the enolate, **3**•enolate. The crystal structure of **3**•PF₆ contained a superposition of two conformers, *exo*-**3** and *endo*-**3**, which we label according to Helmchen's conventional notation specifying the orientation of its allyl fragment with respect to the metal-center plane. The isolated product contained nearly equal amounts of *endo*(49%)- and *exo*(51%)-conformations, indicating both structures have essentially the same energy. The C–C–C angle of the allyl fragment (136.6°) in the crystal structure, however, is quite large and is most likely due to

unresolvable superpositioning of the two structures. Our calculations on *exo*-**3** and *endo*-**3** differ in energy by only 0.2 kcal/mol ($\Delta G_{298}(\textit{exo-3}) > \Delta G_{298}(\textit{endo-3})$). As shown in Figure 1, the predicted structure is in very good agreement with experiment, non-CH bond distances have an RMSD of 0.03 Å, and bond angles are accurate within 2.1°. The error in bond angles reduces to 1.3° when omitting the C–C–C angle of the allyl fragment (calculated = 119.4°). Furthermore, the deviations due to phenyl groups projecting in different positions (Figure 1) are largely corrected when geometry optimizations take into account nonbonding interactions due to dispersion.⁷⁵ Despite the improved agreement in the calculated geometry with the crystal structure, the near-equivalent energies of *exo*-**3** and *endo*-**3** remain the same. Figure 2 displays interatomic bond distances for both *exo*-**3** and the enolate molecule.





Figure 2. Structural representations of *exo*-**3** (a Pd²⁺ complex) and the enolate molecule and a table displaying relevant interatomic distances.

External Nucleophilic Attack Product Formation Reactions. Note that throughout the text, figures, and schemes, we use the labeling notation of Prelog and Seebach,⁷⁶ where 'Re' denotes a pathway leading to an 'R' stereocenter, while 'Si' denotes a pathway leading to an 'S' stereocenter. As mentioned in the introduction, asymmetric allylation reactions using PHOX ligands are usually considered to proceed by an external nucleophilic attack as in Scheme 3. Here, the enolate can couple to intermediates (*endo/exo*)-**3** producing the Pd⁰-bound complexes (*R/S*)-**4**, thereby forming products (*R/S*)-**2**.

Scheme 3. Outer-Sphere External Nucleophilic Attack Processes Considered in This Work


The allyl-coordinated complexes (*endo/exo*)-**3** have two reactive carbons (C_1 or C_3) that may undergo nucleophilic attack. Attack at these carbons would result in one of eight different transition states: (*endo/exo*)-(*Re/Si*)-EXT-(1/2), and each would eventually form products (*R/S*)-**2**. The *trans* influence of phosphine leads to a longer Pd– C_3 bond than that for Pd– C_1 . One might then expect that reactions *trans* to phosphine would be favored, calculated results indicate nucleophilic attacks at either C_1 or C_3 are almost equally favored. After identifying 6 of 8 of these transition states we found all have barriers within 1.3 kcal/mol with respect to each other, the two lowest transition states for their respective asymmetric products were *endo*-(*Si*)-EXT-1 and *exo*-(*Re*)-EXT-2. These transition states are only 0.7 kcal/mol different in energy and thus are well within the accuracy of the computational methods employed. Although we were unable to fully optimize the transition states *exo*-(*Si*)-EXT-1 and *endo*-(*Re*)-EXT-1, the small differences in energy between the other six transition states indicated these two transition states should also be quantitatively similar.

Figure 3 shows structural details of the two lowest energy outer-sphere transition states, *endo*-(*Si*)-EXT-1 and *exo*-(*Re*)-EXT-2. Note that despite the differing modes of nucleophilic attack, both have similar energies and interatomic bond distances. All other nonreacting bonds remain essentially constant with respect to those in *exo*-**3**, though Pd–P and Pd–N bonds increase slightly now that the Pd^{2+} center is undergoing a reduction to Pd^0 . The reacting bonds have intermediate bond lengths, where the breaking Pd–C bonds ($R = 2.32 \text{ \AA}$ and 2.43 \AA) are similar while the forming C–C bonds ($R = 3.19 \text{ \AA}$ and 2.80 \AA) are different due to the steric hindrance of the triphenylphosphine group when the enolate approaches *trans* to the nitrogen atom of the PHOX ligand.

endo-(*Si*)-EXT-1 and *exo*-(*Re*)-EXT-2 have barriers of 20.2 and 20.9 kcal/mol, respectively, with respect to the separated ion pair: *exo*-**3** + enolate. Across all six characterized transition states, the barriers range from 20.2–21.5 kcal/mol. Control calculations using *mPW1PW91* electronic energies result in slightly higher barriers ranging from 20.3–22.2 kcal/mol, but this shows the similar barrier heights are not simply an artifact of the B3LYP exchange correlation functional. On the basis of the proximity of the two electron-rich molecular fragments, we considered the possibility that nonbonding interactions would play a significant role in determining these overall barriers. Indeed, the nonbonding interactions due to dispersion lead to a dramatic reduction in these barriers so that they now range from 10.9–13.1 kcal/mol. This nearly 50% reduction in the activation barriers highlights the importance of dispersion in alkylation reaction mechanisms, though we note that the overall accuracy of dispersion-corrected DFT is not well-established for cases of asymmetric alkylation such as this.




Figure 3. Representations of external nucleophilic transition states *endo*-(*Si*)-EXT-1 and *exo*-(*Re*)-EXT-2 and tables displaying relevant interatomic distances.

The experimental observation that (*S*)-**2** predominates suggests the direct product of allylation should be the Pd-olefin complex (*S*)-**4**, whose geometry is presented in Figure 4. In this structure, the five carbon atoms C_1 – C_5 and the carbonyl oxygen are connected sequentially and their bond lengths




Figure 4. Representations of (*S*)-**4** (a Pd^0 complex) and a table consisting of relevant interatomic distances.

indicate a double bond between C₁ and C₂ ($R = 1.40 \text{ \AA}$), three single bonds ($R = 1.52\text{--}1.55 \text{ \AA}$), and a C–O double bond ($R = 1.22 \text{ \AA}$) that is shorter than the C–O bond encountered in the enolate ion or the external nucleophilic transition states ($R \sim 1.28 \text{ \AA}$). While the Pd–P bond length has remained largely constant throughout the mechanistic steps studied so far ($R = 2.35 \text{ \AA}$), the Pd–N bond has lengthened ($R = 2.40 \text{ \AA}$) now that the metal center has been reduced to Pd⁰ and in gas phase structure prefers linear coordination with phosphorus and the η -2 coordinated product.

The overall energy of (S)-4 is -13.4 kcal/mol with respect to the separately solvated *exo*-3 and enolate species. Control calculations with *mPW1PW91* find this structure is slightly more strongly bound ($\Delta G_{298} = -17.6 \text{ kcal/mol}$) and the single-point dispersion correction makes the overall energy approximately $\sim 7 \text{ kcal/mol}$ more exothermic ($\Delta G_{298} = -20.9 \text{ kcal/mol}$). The η -2 bond strength in (S)-4 is challenging to ascertain precisely since it is dependent on the calculation method employed. B3LYP calculations in solvent suggest it is only 2.8 kcal/mol , while control calculations with *mPW1PW91* find it notably higher (8.7 kcal/mol). The difference between the B3LYP and *mPW1PW91* calculations displays a significant degree of uncertainty comparing energies of Pd²⁺ and Pd⁰ complexes with different exchange correlation functionals. The single-point dispersion correction to the B3LYP energy shows a substantially stronger η -2 bond (11.8 kcal/mol), suggesting dispersion forces play a significant role keeping Pd⁰ coordinatively saturated. Regardless of the methodology used in its energy calculation, (S)-4 is always the lowest energy intermediate on the reaction coordinate.

Inner-Sphere Rearrangements. Up to this point, the reaction pathway for the enantioselective Tsuji allylation is unclear because none of the calculated external nucleophilic attack pathways display an enantioselective process that explains experimentally observed ee values. To investigate further, we sought pathways forming intermediates where C–C bond formation can easily occur and may be influenced by the stereocenter on the PHOX ligand. We considered the feasibility of a 5-coordinate complex resulting from the enolate binding to π -allyl complex *exo*-3 at an axial position. The structure for this complex is shown in Figure 5.

We found several orientations that allow the enolate to bond axially with the *exo*-3 complex. The most stable intermediate,




Figure 5. Representations of 5-coordinate *exo*-5 (a Pd²⁺ complex) and a table displaying relevant interatomic distances.


exo-5, has the enolate nestled within the PHOX ligand and anchored to the Pd with a longer than average bond length of 2.59 \AA . The longer bond length is characteristic of ligands axially coordinated to d^8 transition metals. This Pd–O bond has little influence on the bonding of either the *exo*-3 fragment or the enolate as seen in a comparison of bond lengths with those in Figure 2. Other structures also had the enolate axially bound to Pd, but those require the enolate to have less favorable steric interactions with the PHOX ligand. When the enolate is bound to *endo*-3, the energy of the structure is $\sim 4 \text{ kcal/mol}$ higher than when it is bound to *exo*-3. Thus, while *exo*-3 and *endo*-3 have effectively the same energy, axial coordination favors the *exo* form.

As with the energy of (S)-4, the energy of the Pd–O bond is highly dependent on the calculation scheme. Because *exo*-3 is a cation and the enolate is an anion, the separated ionic species are very unstable in vacuum, and bringing both species together results in an electronic energy gain of -72.9 kcal/mol . If zero point energies, thermal corrections, and entropic energy are added, the free energy released when bringing the two species together is less negative, -56.5 kcal/mol . Because the allylation reaction occurs in solution, the energy to form *exo*-5 should be referenced to the free energies in solution of the separately solvated *exo*-3 and enolate species. Under this model, solvated *exo*-5 is now *unstable* by $+16.7 \text{ kcal/mol}$, though it is still $\sim 3.5 \text{ kcal/mol}$ lower in energy than the external attack transition states reported in the previous section. Control calculations with *mPW1PW91* provide a value that is slightly less unstable, $+14.0 \text{ kcal/mol}$. Finally, adding single-point dispersion corrections to energies of solvated species causes *exo*-5 to be effectively isoenergetic (with *exo*-5 favored by merely -0.2 kcal/mol with respect to the solvated ions), and $\sim 10 \text{ kcal/mol}$ more stable than the external attack transition states. Note that in our previous communication on the Tsuji mechanism,³⁰ we reported only electronic energies with solvation corrections, whereas the present work shows that including the free energy penalty to form *exo*-5 essentially cancels out the energy gain captured by dispersion corrections, making *exo*-5 effectively isoenergetic with the separated ions in solution.

With routes to *exo*-5 established, we considered several internal rearrangements, shown in Scheme 4, that lead to 4-coordinate intermediates and can undergo reductive elimination to form Pd⁰ olefin complex (S)-4. One possibility has the enolate O atom exchange into the binding site of the PHOX ligand N atom. On the basis of the long Pd–N bond in (S)-4, it seemed reasonable that the N atom donor from the PHOX ligand would be displaced more readily than the phosphorus donor.

The energies of (Si)-6 and (Re)-6, are $+8.9$ and $+9.3 \text{ kcal/mol}$ with respect to the separated ion pair in solution. The energies are similar to those obtained with *mPW1PW91* control calculations ($+8.6$ and $+9.2 \text{ kcal/mol}$, respectively); however, dispersion energy corrections again greatly reduce this instability making the energies -0.2 and $+0.6 \text{ kcal/mol}$, respectively. Thus, the energies of (Si)-6 and (Re)-6 are almost the same as those of the separated *exo*-3 and enolate ions as well as *exo*-5. The transition states leading to these intermediates, (Si)-INT-1 and (Re)-INT-1 are $+22.4$ and $+24.9 \text{ kcal/mol}$, values that are slightly higher than those for external nucleophilic attack. Even with *mPW1PW91* and dispersion corrections, the barriers remain higher than those of expected racemizing processes. Breaking the Pd–N bond would also be unfavorable from an enantioinduction standpoint because that

Scheme 4. Internal Rearrangement Mechanisms Considered in This Work



side of the PHOX ligand contains the stereocenter that makes the reaction enantioselective.

Another possibility involves the O atom of the enolate exchanging into the binding site of the C₁ atom of the allyl group. During this process the η -3 allyl fragment slips to become an η -1-bonded allyl group, which may be a more favorable position from which to facilitate C–C bond formation. However, the energies for the 4-coordinate species where the enolate becomes *trans* to the N, (*Si*)-7 and (*Re*)-7, as well as those of their corresponding transition states, (*Si*)-INT-2 and (*Re*)-INT-2, are higher than those of the cases where the N atom of the PHOX ligand underwent substitution. Thus, we also ruled out these as feasible enantioselective processes.

The last rearrangement we present was expected to be the most favorable because (1) the PHOX ligand remains intact throughout the reaction and (2) the *trans* influence of the phosphino group should help activate the P–C₃ bond. Indeed, the 4-coordinate species (*Si*)-8 and (*Re*)-8 are more stable than those just mentioned. Again, B3LYP free energies with solvation show the complexes are less stable than the separated *exo*-3 and enolate ions at +5.7 and +5.9 kcal/mol, respectively. As with (*Si*)-7 and (*Re*)-7, *m*PW1PW91 single-point electronic energies make these slightly higher: both are +6.7 kcal/mol. However, adding dispersion corrections now makes these intermediates –4.8 and –4.7 kcal/mol more stable than the separated ions. Unlike all of the other pathways from Scheme 4, this pathway has a significant thermodynamic driving force, and so we now focus our discussion on the transition states that lead to these intermediates, (*Si*)-INT-3 and (*Re*)-INT-3.

The geometries for these transition states are shown in Figure 6. Interestingly, this is the first transition state we have discussed that also shows promise regarding enantioselectivity. Almost all interatomic distances are similar between the two transition states with the notable exception of the breaking Pd–C₃ bond, which is significantly longer in (*Re*)-INT-3 (*R* = 2.61 Å) than it is in (*Si*)-INT-3 (*R* = 2.37 Å). The orientation of the methyl group influences the transition state to be later along the reaction coordinate in (*Re*)-INT-3 than in (*Si*)-INT-3.




Figure 6. Representations of transition states (*Si*)-INT-3 and (*Re*)-INT-3 and a table consisting of relevant interatomic distances.

Figure 7 shows another perspective of these transition states, this time with the π -allyl Pd(PHOX) catalyst rendered with space-filling spheres, while the enolate is rendered with a ball and stick model. The top view of (*Re*)-INT-3 (Figure 7a) shows the enolate's methyl group quite near to a phenyl group of the PHOX ligand, while the top view of (*Si*)-INT-3 (Figure




Figure 7. Top and side view representations for (*Re*)-INT-3 and (*Si*)-INT-3.

7c) has the methyl group pointing away from the phenyl group and poised above the oxazoline ring of the PHOX ligand. Side views of these two transition states (a and b of Figure 7) illustrate that the two faces of the enolate interact differently with the PHOX ligand.

The observed ee values for the enantioselective Tsuji allylation reaction are approximately 90%, corresponding to an (*S*):(*R*) product ratio of 95:5. A Boltzmann relation of this ratio in turn corresponds to a $\Delta\Delta G_{298}$ between the enantiodetermining reaction barriers to be only ~ 1.8 kcal/mol. Given the standard expectation of B3LYP to yield accurate differences between barrier heights within ~ 2 kcal/mol, capturing this energy difference accurately with theoretical calculations is probably not possible. Nevertheless, B3LYP calculated barrier heights are in excellent agreement with this value, calculating the barrier for (*Si*)-INT-3 to be 1.0 kcal/mol lower than the barrier for (*Re*)-INT-3. Barriers using single-point *mPW1PW91* calculations serendipitously makes this difference 1.6 kcal/mol, while inclusion of dispersion energies causes the barrier for (*Re*)-INT-3 to be energetically lower than (*Si*)-INT-3 by 1.4 kcal/mol. Dispersion corrections seem to favor a pathway that would result in asymmetric (*R*)-4 products rather than the observed (*S*)-4 products, but we caution against overinterpretation of barrier heights differing in energy by < 2 kcal/mol.

When considering absolute barrier heights, B3LYP calculates the barriers for (*Si*)-INT-3 and (*Re*)-INT-3 to be 18.4 and 19.4 kcal/mol, respectively. Both values are smaller than the lowest barrier for external nucleophilic attack by 0.8 kcal/mol. Absolute barriers from *mPW1PW91* calculations are 16.5 and 18.1 kcal/mol, respectively, and these are now more than 2 kcal/mol lower than the lowest external nucleophilic attack barrier. Finally, adding dispersion corrections makes the barriers 5.3 and 3.9 kcal/mol, respectively, which are more than 5 kcal/mol lower than the lowest external nucleophilic attack barrier. Thus, while the three different calculation methods disagree within a range of < 2 kcal/mol as to which enantiodetermining process is favored, the three different methods unanimously identify that the internal rearrangement mechanisms, (*Si*)-INT-3 and (*Re*)-INT-3, are energetically lower than external nucleophilic attack processes. Therefore, the internal pathways are most likely to explain this chemistry. NBO charges⁷⁷ between respective atoms on the enolate and the Pd center differ by only ~ 0.02 charge units indicating electronic substitutions will not significantly favor one reaction pathway over another. However, the structural interactions between the enolate and the PHOX ligands differ between the two transition states; thus, the enantioselectivity must be due to steric interactions.

Figure 8 shows the geometric structures for the 4-coordinate species (*Si*)-8; however, the relevant interatomic distances for (*Re*)-8 are the same. These geometries are consistent with expected structures from 4-coordinate Pd²⁺ complexes. The Pd–P bond remains the same length as that found in *exo*-3; however, the Pd–N bond is lengthened due to the *trans* influence of the C₁, which has now formed an η -1 bond. The Pd–C₃ bond in turn is completely dissociated, but the elongated C₁–C₂ and shortened C₂–C₃ bonds show the allyl fragment is no longer in resonance. One can visualize how the allyl and enolate ligands fit next to each other on the basis of the space-filling models depicted in Figure 8.

Rotational barriers of either the allyl fragment (about the Pd–C₁ bond) or the enolate fragment (about the Pd–O bond)





Figure 8. Representations of 4-coordinate (*Si*)-8 and (*Re*)-8 (Pd²⁺ complexes), formed after inner-sphere ligand rearrangement. The table displays relevant interatomic distances.

within this complex are challenging to obtain with quantum mechanics since the potential energy surface will be largely due to non-bonding interactions. Nevertheless, knowing the barrier for the latter rotation is important to know since a free rotation would lead a racemic mixture of (*Re/Si*)-8, and would therefore render the potentially enantiodetermining barriers from the transition states (*Re/Si*)-INT-3 irrelevant to enantioselectivity. Determining such barriers would require dynamical simulations to ensure the potential energy surface was sufficiently sampled, and dispersion will most certainly play a critical role in these interactions as well. We have not yet calculated rotational barriers; however, we note that the reverse barrier from (*Si*)-8 to *exo*-5 via (*Si*)-INT-3 is between 17 and 18 kcal/mol without dispersion corrections and ~ 10 kcal/mol when including the corrections. Overcoming that barrier would result in another opportunity for the reaction to proceed via (*Re*)-INT-3 instead of (*Si*)-INT-3, thereby leading to racemized products. Thus, these barriers are the upper limits of what subsequent barriers in the C–C bond-forming steps would need to be in order to form asymmetric products.

From the 4-coordinate species, (*Re/Si*)-8, we then sought reaction pathways that would lead to asymmetric products (Scheme 5). Focusing first on a process leading to a traditional three-centered reductive elimination we characterized reaction pathways whereby the enolate's carbonyl O atom would exchange with the tertiary α -C atom, (*Re/Si*)-INT-4. We could only fully optimize the (*Re*)-INT-4 transition state; however,

Scheme 5. Pathways for C–C Bond Formation Considered in This Work



we see no reason why that barrier would not be a reasonable approximation for *(Si)*-INT-4 as well. Indeed, this barrier is quite high regardless of the method employed. Calculations using B3LYP with solvation resulted in a barrier of 28.0 kcal/mol, while those from *m*PW1PW91 calculations were very similar: 27.1 kcal/mol. These barriers were ~ 5 kcal/mol higher than the corresponding barriers for external nucleophilic attack. Dispersion energies made this barrier similar to external attack barriers (12.9 kcal/mol) and only a few kcal/mol higher than the reverse process back to *exo*-5.

We also considered traditional 3-centered reductive elimination from the intermediates *(Re/Si)*-9. The energies of these intermediates were roughly the same as those of the 5-coordinate species *exo*-5, and B3LYP and *m*PW1PW91 calculations found the energies of the intermediates approximately +16 kcal/mol less stable than those of the separated intermediates *exo*-3 + enolate; however, dispersion corrections stabilized their energies to about the same as the energies of *exo*-3 + enolate as well as that of *exo*-5. Notably, the energy of *(Si)*-9 is consistently >2 kcal/mol more stable than that of *(Re)*-9. However, in the reductive elimination process *(Si)*-INT-5 has an extraordinarily high barrier: >45 kcal/mol regardless of calculation method.⁴

Having ruled out three-centered reductive elimination mechanisms, we considered different mechanisms analogous to those first reported by Echavarren and co-workers:⁸⁰ coupling allyl fragments together at their terminal carbons, *i.e.*, between C₃ on the allyl fragment and C₄ on the enolate. These mechanisms have also received consideration in other mechanistic studies.^{81–85} While these reactions appear to be a hybrid of the Cope rearrangement and a cheletropic reaction, the process we study resembles a hybrid of a Claisen and reductive cheletropic rearrangement.⁸⁶ This process shown by *(Re/Si)*-INT-6 may also be considered as an intramolecular metallacyclopentane reaction leading to a palladacyclopentane complex, *i.e.* *(R/S)*-4.

The transition states for these processes were *(Re/Si)*-INT-6, and each process was considered a pericyclic reaction process where the allyl and enol fragments were aligned in boat- and chairlike structures. Consistent with sterics for six-membered rings, as well as with the work by Echavarren, the chairlike processes had the lowest energies. Figure 9 shows the geometric structure for *(Si)*-INT-6.




Figure 9. Representations of the seven-membered reductive elimination processes *(Si)*-INT-6 and *(Re)*-INT-6. The tables display relevant interatomic distances.

In this process the Pd–P bond remained intact as it has throughout the entire mechanism, but the Pd–N, Pd–C₁, and Pd–O bonds are longer with respect to those seen in *(Si)*-8. In fact, the Pd–N bond in *(Si)*-INT-6 is slightly shorter than that found in the Pd⁰ complex *(S)*-4, indicating the occupation of antibonding orbitals during the reduction. The transition state has an intermediate C₃–C₄ distance of 2.17 Å, and the bond distances about the Pd center indicate that bond formation and Pd²⁺ reduction to Pd⁰ is a concerted process. The geometry of the *(Re)*-INT-6 transition state is almost exactly the same as that for *(Si)*-INT-6, and interatomic distances vary by less than 0.03 Å.

The energies of the boat- and chair-*(Re/Si)*-INT-6 transition states are likewise almost identical; thus, it is unlikely these steps themselves would be enantiodetermining. The respective barriers for chair-*(Si)*-INT-6 and chair-*(Re)*-INT-6 were +16.9 and +16.8 kcal/mol using B3LYP with solvation, +16.9 and +17.1 kcal/mol with *m*PW1PW91 electronic energies, and dispersion corrections reduced the B3LYP barriers to +7.3 and +7.1 kcal/mol.

A partial picture of the enantioselective Tsuji allylation reaction is now complete. We refer the reader to Figure 10, which summarizes all key energies in the calculated mechanism. In calculations with B3LYP free energies with solvation (Figure 10a), combining *exo*-3 with the enolate is endothermic, and not a stable intermediate in THF solution. (Note that this energy does not include dispersion interactions, but those contributions are included in Figure 10c.) The calculated barriers suggest that internal rearrangement is an enantioselective process favoring the channel leading to *(S)*-2, the experimentally observed major product. Internal rearrangement, *(Si)*-INT-3, is 1.0 kcal/mol lower in energy than *(Re)*-INT-3,




Figure 10. Mechanistic pathways summarizing three different calculation schemes used in this study. Numbers depict relative free energies (in kcal/mol) for relevant intermediates and transition states in the reaction mechanism. (a) Energies calculated from B3LYP/BS3//B3LYP/BS2 including single-point solvation corrections. (b) Same energies as as those for (a) except with mPW1PW91/BS3 single-point electronic energies. (c) Same energies as those for (a) except including single-point dispersion energy corrections. See text for the discussion.

leading to a $\Delta\Delta G_{298}$ value of 1.0 kcal/mol. If racemization of (Si)-8 to (Re)-8 does not occur, the high backward barrier back to *exo*-5 would mean all of the (Re/Si)-8 intermediates would

react via (Re/Si)-INT-6 and the ratio of (S)-2 to (R)-2 would be 84:16. Since outer-sphere reaction barriers are higher, products arising from these pathways are unlikely.

In calculations similar to the previous case except using mPW1PW91 electronic energies (Figure 10b), forming *exo*-5 is still substantially endothermic. The calculated barriers still suggest that internal rearrangement is an enantioselective process favoring the channel leading to S-2, but (Si)-INT-3 is now 1.6 kcal/mol lower in energy than (Re)-INT-3, leading to a $\Delta\Delta G_{298}$ value of 1.6 kcal/mol. The backward barrier to *exo*-5 is substantially lower in this calculation model, but because it is still lower than the (Re)-INT-3 barrier, all (Re/Si)-8 intermediates would react via (Re/Si)-INT-6 resulting in a (S)-2 to (R)-2 ratio of 94:6. Again, no products would arise from outer-sphere attack mechanisms. These results establish a measure of the uncertainties facing the DFT calculations without empirical dispersion corrections, and that a quantitatively similar perspective from theory is presented with two different DFT exchange correlation functionals.

In B3LYP free energy calculations with solvation and single-point dispersion corrections (Figure 10c), a substantially different picture is observed. Formation of *exo*-5 is now very slightly exothermic after changing the relative thermodynamics of single step by almost 17 kcal/mol. This suggests first that dispersion plays a significant role in this mechanism allowing the 5-coordinate *exo*-5 to act as a gateway to the inner-sphere processes. The (Re/Si)-INT-3 processes are affected differently by dispersion; however, both are now substantially lower than the subsequent (Re/Si)-INT-6 processes. At face value, this causes (Re/Si)-INT-3 not to be an enantioselective process, and a roughly racemic mixture of (Re/Si)-8 would be expected during the reaction. However, on the basis of empirically based single-point energy corrections, we note that the current treatment of dispersion will likely bring with it greater uncertainties with energies requiring high precision (such as the barrier heights that determine the enantioselectivity of the reaction).

A natural question is which of the mechanistic profiles shown in Figure 10 is most accurately representing the true chemistry of this process. Since all of the geometries for these mechanisms were obtained using B3LYP optimizations without dispersion corrections, the values presented in Figure 10a will contain the most error cancellation. Thus, the $\Delta\Delta G_{298}$ values for the enantioselective steps are the most meaningful when taken from Figure 10a. Indeed these results match well with experiment, while those in the control calculations shown in Figure 10b serendipitously are in perfect agreement with experiment. In both cases, transition states (Re)-INT-3 and (Si)-INT-3 are enantiodetermining pathways, and these mechanisms predict the experimentally observed products. Figure 10c shows calculations involving dispersion corrections, which are undoubtedly important in this mechanism. Dispersion causes most inner-sphere intermediate energies to be shifted down by 10–15 kcal/mol. Due to this, the lower barriers seen here are probably more appropriate for modeling overall reaction kinetics; however, as noted before, the precision of the enantioselective steps and their corresponding $\Delta\Delta G_{298}$ values will not be as robust, since these calculations have not been taken at stationary points on the DFT + dispersion potential energy surface. To alleviate this problem, a more rigorous calculation scheme involving geometries optimized with dispersion and/or solvation corrections at each iteration could be pursued; however, the computational

cost of such calculations outweighs the marginal gain over the data presented in Figure 10a. Instead, we use this ensemble of schemes to provide two general suggestions to improve the enantioselectivity of this reaction. First, chemical modifications to greater stabilize the (Si)-INT-3 transition state with respect to (Re)-INT-3 should improve enantioselectivities. Furthermore, modifications to make the intermediates (Re/Si)-8 as stable as possible will in turn ensure that those enantioselective products will rapidly convert to final products before reverse or racemizing reactions take place.

CONCLUSIONS

On the basis of our computational results, we propose a mechanistic model to describe the unique performance of the enantioselective Tsuji allylation reaction involving prochiral nucleophiles and nonprochiral allyl fragments. The calculated pathways governing this process are presented in Figure 10. We note that enantioselectivity is only possible when the substrate undergoing allylation is a highly basic enolate nucleophile and/or susceptible to large nonbonding interactions due to dispersion such that it will form the *exo*-5 intermediate. We believe a competition between *Re*- and *Si*-pathways then ensues, and the most likely reaction step with enantiodetermining character is an internal rearrangement from the 5-coordinate *exo*-5 to the 4-coordinate (Si)-8 via the transition state (Si)-INT-3. Following this process, a nontraditional reductive elimination process occurs whereby C–C bond formation occurs at the 3 and 3' positions of the allyl and enolate fragments.

On the basis of these results, we predict that this mechanism will be sensitive to by substitutions at the *ortho*-positions of the phenyl groups on the phosphine ligand to further obstruct (Re)-INT-3 while leaving (Si)-INT-3 largely unaffected. Idealized substitutions may enhance activity, though too large of a substitution runs the risk of interfering too much with the stability of *exo*-5, thus completely shutting down the inner-sphere pathways, leaving only outer-sphere pathways possible and leading to racemic products.⁸⁷ Once the enantioselective step has been optimized, the (Re/Si)-INT-6 barriers must be lower than the backward (Re/Si)-INT-3 barriers to maximize yields. Investigations with modified substrates and ligands are underway to further improve the performance of these catalysts.

ASSOCIATED CONTENT

Supporting Information

Cartesian coordinates and energies of all calculated species mentioned in the text and CIF data for species *exo*-3. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

stoltz@caltech.edu (B.M.S.); wag@wag.caltech.edu (W.A.G.)

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank K. Tani, A. Harned, and J. Enquist for experimental collaboration and discussion, and M. Day and L. Henling for crystallography assistance. J.A.K. thanks A. Tkatchenko for discussions and calculations with dispersion-corrected DFT. This research was partly funded by Chevron-Texaco, and the

facilities used were funded by grants from ARO–DURIP, ONR–DURIP, IBM–SUR, Fannie and John Hertz Foundation (D.C.B.), and Eli Lilly (J.T.M.) with additional support from NSF (CTS-0608889 and CHE-1214158, W.A.G.) and NIH–NIGMS (R01GM080269-01, B.M.S.).

ADDITIONAL NOTE

^aThis high barrier is consistent with earlier work that explained that coupling sp^3 hybridized organic fragments face a high barrier due to the ground state electronic configuration of Pd^0 .^{78,79} The reductive elimination of fragments bound with directional sp^3 bonds would be best facilitated by a metal center with *s*-orbital character in its valence shell. For Pd, reduction from $Pd^{2+} \rightarrow Pd^0$ leads to a change in electronic configuration from $d^8 \rightarrow d^{10}$, so that no *s*-character is available in the reaction. In the case of Pt, however, reduction from $Pt^{2+} \rightarrow Pt^0$ leads to a change in electronic configuration from $d^8 \rightarrow s^1d^9$, so that such reductive eliminations with Pt may be possible.

REFERENCES

- (1) Cornils, B.; Herrmann, W. A., Eds. *Applied Homogeneous Catalysis with Organometallic Compounds: A Comprehensive Handbook in Three Volumes*, 2nd ed.; Wiley-VCH: Weinheim, Germany, 2002.
- (2) Jacobsen, E. N.; Pfaltz, A.; Yamamoto, H., Eds.; *Comprehensive Asymmetric Catalysis*, 1st ed.; Springer: New York, NY, 1999.
- (3) Ojima, I., Ed. *Catalytic Asymmetric Synthesis*, 3rd ed.; John Wiley & Sons, Inc.: Hoboken, NJ, 2010.
- (4) Pfaltz, A.; Lautens, M. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer: New York, NY, 1999; Vol. 2, pp 833–884.
- (5) Behenna, D. C.; Stoltz, B. M. *J. Am. Chem. Soc.* **2004**, *126*, 15044–15045.
- (6) Mohr, J. T.; Behenna, D. C.; Harned, A. M.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2005**, *44*, 6924–6927.
- (7) Mohr, J. T.; Stoltz, B. M. *Chem.—Asian J.* **2007**, *2*, 1476–1491.
- (8) Behenna, D. C.; Liu, Y.; Yurino, T.; Kim, J.; White, D. E.; Virgil, S. C.; Stoltz, B. M. *Nature Chem.* **2011**, *4*, 130–133.
- (9) Tsuji, J.; Minami, I. *Acc. Chem. Res.* **1987**, *20*, 140–145.
- (10) Weaver, J. D.; Recio, A.; Grenning, A. J.; Tunge, J. A. *Chem. Rev.* **2011**, *111*, 1846–1913.
- (11) Auburn, P. R.; Mackenzie, P. B.; Bosnich, B. *J. Am. Chem. Soc.* **1985**, *107*, 2033–2046.
- (12) Trost, B. M. *Acc. Chem. Res.* **1996**, *29*, 355–364.
- (13) Trost, B. M.; Murphy, D. J. *Organometallics* **1985**, *4*, 1143–1145.
- (14) Trost, B. M. *Chem. Pharm. Bull.* **2002**, *50*, 1–14.
- (15) Trost, B. M. *J. Org. Chem.* **2004**, *69*, 5813–5837.
- (16) Trost, B. M.; Lee, C. In *Catalytic Asymmetric Synthesis*, 2nd ed.; Ojima, I., Ed.; Wiley-VCH: New York, NY, 2000; pp 593–649.
- (17) Trost, B. M.; Van Vranken, D. L. *Chem. Rev.* **1996**, *96*, 395–422.
- (18) Helmchen, G. *J. Organomet. Chem.* **1999**, *576*, 203–214.
- (19) Steinhagen, H.; Reggelin, M.; Helmchen, G. *Angew. Chem., Int. Ed.* **1997**, *36*, 2108–2110.
- (20) Trost, B. M.; Xu, J.; Schmidt, T. *J. Am. Chem. Soc.* **2009**, *131*, 18343–18357.
- (21) Trost, B. M.; Thaisrivongs, D. A. *J. Am. Chem. Soc.* **2008**, *130*, 14092–14093.
- (22) Trost, B. M.; Bream, R. N.; Xu, J. *Angew. Chem., Int. Ed.* **2006**, *45*, 3109–3112.
- (23) Trost, B. M.; Xu, J. *J. Am. Chem. Soc.* **2005**, *127*, 17180–17181.
- (24) Butts, C. P.; Filali, E.; Lloyd-Jones, G. C.; Norrby, P.-O.; Sale, D. A.; Schramm, Y. *J. Am. Chem. Soc.* **2009**, *131*, 9945–9957.
- (25) Rayabarapu, D. K.; Tunge, J. A. *J. Am. Chem. Soc.* **2005**, *127*, 13510–13511.
- (26) Sheffy, F. K.; Godschalk, J. P.; Stille, J. K. *J. Am. Chem. Soc.* **1984**, *106*, 4833–4840.

- (27) Sheffy, F. K.; Stille, J. K. *J. Am. Chem. Soc.* **1983**, *105*, 7173–7175.
- (28) Tsuji, Y.; Funato, M.; Ozawa, M.; Ogiyama, H.; Kajita, S.; Kawamura, T. *J. Org. Chem.* **1996**, *61*, 5779–5787.
- (29) Trost, B. M.; Verhoeven, T. R. *J. Am. Chem. Soc.* **1980**, *102*, 4730–4743.
- (30) Keith, J. A.; Behenna, D. C.; Mohr, J. T.; Ma, S.; Marinescu, S. C.; Oxgaard, J.; Stoltz, B. M.; Goddard, W. A., III. *J. Am. Chem. Soc.* **2007**, *129*, 11876–11877.
- (31) Hayashi, T.; Kanehira, K.; Hagihara, T.; Kumada, M. *J. Org. Chem.* **1988**, *53*, 113–120.
- (32) Helmchen, G.; Steinhagen, H.; Reggelin, M.; Kudis, S. In *Selective Reactions of Metal-Activated Molecules: Third Symposium*; Werner, H., Schreier, P., Eds.; Friedrich Vieweg & Sohn Verlagsgesellschaft GmbH: Wiesbaden, 1998; pp 205–215.
- (33) Kuwano, R.; Ito, Y. *J. Am. Chem. Soc.* **1999**, *121*, 3236–3237.
- (34) Sawamura, M.; Nagata, H.; Sakamoto, H.; Ito, Y. *J. Am. Chem. Soc.* **1992**, *114*, 2586–2592.
- (35) Trost, B. M.; Schroeder, G. M. *Chem.—Eur. J.* **2005**, *11*, 174–184.
- (36) McFadden, R. M.; Stoltz, B. M. *J. Am. Chem. Soc.* **2006**, *128*, 7738–7739.
- (37) Levine, S. R.; Krout, M. R.; Stoltz, B. M. *Org. Lett.* **2008**, *11*, 289–292.
- (38) Petrova, K. V.; Mohr, J. T.; Stoltz, B. M. *Org. Lett.* **2008**, *11*, 293–295.
- (39) Mukherjee, H.; McDougal, N. T.; Virgil, S. C.; Stoltz, B. M. *Org. Lett.* **2011**, *13*, 825–827.
- (40) Hagiwara, H.; Fukushima, M.; Kinugawa, K.; Matsui, T.; Hoshi, T.; Suzuki, T. *Tetrahedron* **2011**, *67*, 4061–4068.
- (41) Day, J. J.; McFadden, R. M.; Virgil, S. C.; Kolding, H.; Alleva, J. L.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2011**, *50*, 6814–6818.
- (42) White, D. E.; Stewart, I. C.; Grubbs, R. H.; Stoltz, B. M. *J. Am. Chem. Soc.* **2007**, *130*, 810–811.
- (43) White, D. E.; Stewart, I. C.; Seashore-Ludlow, B. A.; Grubbs, R. H.; Stoltz, B. M. *Tetrahedron* **2010**, *66*, 4668–4686.
- (44) Enquist, J. A., Jr; Stoltz, B. M. *Nature* **2008**, *453*, 1228–1231.
- (45) Enquist, J. A., Jr; Virgil, S. C.; Stoltz, B. M. *Chem.—Eur. J.* **2011**, *17*, 9957–9969.
- (46) Mohr, J. T.; Nishimata, T.; Behenna, D. C.; Stoltz, B. M. *J. Am. Chem. Soc.* **2006**, *128*, 11348–11349.
- (47) Marinescu, S. C.; Nishimata, T.; Mohr, J. T.; Stoltz, B. M. *Org. Lett.* **2008**, *10*, 1039–1042.
- (48) Streuff, J.; White, D. E.; Virgil, S. C.; Stoltz, B. M. *Nature Chem.* **2010**, *2*, 192–196.
- (49) Helmchen, G.; Pfaltz, A. *Acc. Chem. Res.* **2000**, *33*, 336–345.
- (50) Williams, J. M. J. *Synlett* **1996**, *1996*, 705–710.
- (51) Behenna, D. C.; Mohr, J. T.; Sherden, N. H.; Marinescu, S. C.; Harned, A. M.; Tani, K.; Seto, M.; Ma, S.; Novák, Z.; Krout, M. R.; McFadden, R. M.; Roizen, J. L.; Enquist, J. A., Jr; White, D. E.; Levine, S. R.; Petrova, K. V.; Iwashita, A.; Virgil, S. C.; Stoltz, B. M. *Chem.—Eur. J.* **2011**, *17*, 14199–14223.
- (52) Kollmar, M.; Goldfuss, B.; Reggelin, M.; Rominger, F.; Helmchen, G. *Chem.—Eur. J.* **2001**, *7*, 4913–4927.
- (53) Kollmar, M.; Steinhagen, H.; Janssen, J. P.; Goldfuss, B.; Malinovskaya, S. A.; Vázquez, J.; Rominger, F.; Helmchen, G. *Chem.—Eur. J.* **2002**, *8*, 3103–3114.
- (54) Matt, P. V.; Lloyd-Jones, G. C.; Minidis, A. B. E.; Pfaltz, A.; Macko, L.; Neuburger, M.; Zehnder, M.; Rüegger, H.; Pregosin, P. S. *Helv. Chim. Acta* **1995**, *78*, 265–284.
- (55) *Jaguar*; Schrödinger, LLC: New York, NY, 2007.
- (56) Stephens, P. J.; Devlin, F. J.; Chabalowski, C. F.; Frisch, M. J. *J. Phys. Chem.* **1994**, *98*, 11623–11627.
- (57) Becke, A. D. *Phys. Rev. A* **1988**, *38*, 3098–3100.
- (58) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785–789.
- (59) Keith, J. A.; Oxgaard, J.; Goddard, W. A., III. *J. Am. Chem. Soc.* **2006**, *128*, 3132–3133.
- (60) Keith, J. M.; Nielsen, R. J.; Oxgaard, J.; Goddard, W. A., III. *J. Am. Chem. Soc.* **2005**, *127*, 13172–13179.
- (61) Nielsen, R. J.; Goddard, W. A., III. *J. Am. Chem. Soc.* **2006**, *128*, 9651–9660.
- (62) Nielsen, R. J.; Keith, J. M.; Stoltz, B. M.; Goddard, W. A., III. *J. Am. Chem. Soc.* **2004**, *126*, 7967–7974.
- (63) Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 270–283.
- (64) Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 299–310.
- (65) Easton, R. E.; Giesen, D. J.; Welch, A.; Cramer, C. J.; Truhlar, D. G. *Theory Chem. Acta* **1996**, *93*, 281–301.
- (66) Adamo, C.; Barone, V. *J. Chem. Phys.* **1998**, *108*, 664–675.
- (67) Marten, B.; Kim, K.; Cortis, C.; Friesner, R. A.; Murphy, R. B.; Ringnalda, M. N.; Sitkoff, D.; Honig, B. *J. Phys. Chem.* **1996**, *100*, 11775–11788.
- (68) Tannor, D. J.; Marten, B.; Murphy, R.; Friesner, R. A.; Sitkoff, D.; Nicholls, A.; Honig, B.; Ringnalda, M.; Goddard, W. A., III. *J. Am. Chem. Soc.* **1994**, *116*, 11875–11882.
- (69) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. *J. Chem. Phys.* **2010**, *132*, 154104.
- (70) Peverati, R.; Baldrige, K. K. *J. Chem. Theor. Comput.* **2008**, *4*, 2030–2048.
- (71) Schmidt, M. W.; Baldrige, K. K.; Boatz, J. A.; Elbert, S. T.; Gordon, M. S.; Jensen, J. H.; Koseki, S.; Matsunaga, N.; Nguyen, K. A.; Su, S.; Windus, T. L.; Dupuis, M.; Montgomery, J. A., Jr. *J. Comput. Chem.* **1993**, *14*, 1347–1363.
- (72) Gordon, M. S.; Schmidt, M. W. In *Theory and Applications of Computational Chemistry: The First Forty Years*; Dykstra, C., Frenking, G., Kim, K., Scuseria, G., Eds.; Elsevier Science: Amsterdam, 2005; pp 1167–1189.
- (73) Humphrey, W.; Dalke, A.; Schulten, K. *J. Mol. Graphics* **1996**, *14*, 33–38.
- (74) Cohen, N.; Benson, S. W. *Chem. Rev.* **1993**, *93*, 2419–2438.
- (75) Determined with the help of A. Tkachenko and his dispersion-correction code developed at the FHI-Berlin.
- (76) Seebach, D.; Prelog, V. *Angew. Chem., Int. Ed.* **1982**, *21*, 654–660.
- (77) Glendening, E. D.; Badenhop, J. K.; Reed, A. E.; Carpenter, J. E.; Bohman, J. A.; Morales, C. M.; Weinhold, F. *NBO 5.0*; Theoretical Chemistry Institute, University of Wisconsin: Madison, WI, 2001.
- (78) Low, J. J.; Goddard, W. A., III. *Organometallics* **1986**, *5*, 609–622.
- (79) Low, J. J.; Goddard, W. A., III. *J. Am. Chem. Soc.* **1986**, *108*, 6115–6128.
- (80) Méndez, M.; Cuerva, J. M.; Gómez-Bengoia, E.; Cárdenas, D. J.; Echavarren, A. M. *Chem.—Eur. J.* **2002**, *8*, 3620–3628.
- (81) Ariafard, A.; Lin, Z. *J. Am. Chem. Soc.* **2006**, *128*, 13010–13016.
- (82) García-Iglesias, M.; Buñuel, E.; Cárdenas, D. J. *Organometallics* **2006**, *25*, 3611–3618.
- (83) Sieber, J. D.; Liu, S.; Morken, J. P. *J. Am. Chem. Soc.* **2007**, *129*, 2214–2215.
- (84) Waetzig, S. R.; Rayabarapu, D. K.; Weaver, J. D.; Tunge, J. A. *Angew. Chem., Int. Ed.* **2006**, *45*, 4977–4980.
- (85) Wallner, O. A.; Szabó, K. *J. Chem.—Eur. J.* **2006**, *12*, 6976–6983.
- (86) Lutz, R. P. *Chem. Rev.* **1984**, *84*, 205–247.
- (87) Behenna, D. C. Progress toward the synthesis of (+)-zoanthanol and the development of an asymmetric Tsuji allylation reaction. Ph. D. Dissertation, California Institute of Technology: Pasadena, CA, 2007.