

# Azurin as a Protein Scaffold for a Low-coordinate Nonheme Iron Site with a Small-molecule Binding Pocket

Matthew P. McLaughlin,<sup>†</sup> Marius Retegan,<sup>‡</sup> Eckhard Bill,<sup>‡</sup> Thomas M. Payne,<sup>§</sup> Hannah S. Shafaat,<sup>‡</sup> Salvador Peña,<sup>†</sup> Jawahar Sudhamsu,<sup>§</sup> Amy A. Ensign,<sup>†</sup> Brian R. Crane,<sup>§</sup> Frank Neese,<sup>\*,‡</sup> and Patrick L. Holland<sup>\*,†</sup>


<sup>†</sup>Department of Chemistry, University of Rochester, Rochester, New York 14618, United States

<sup>§</sup>Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853, United States

<sup>‡</sup>Max Planck Institute for Chemical Energy Conversion, Mülheim an der Ruhr, Germany

## Supporting Information

**ABSTRACT:** The apoprotein of *Pseudomonas aeruginosa* azurin binds iron(II) to give a 1:1 complex, which has been characterized by electronic absorption, Mössbauer, and NMR spectroscopies, as well as X-ray crystallography and quantum-chemical computations. Despite potential competition by water and other coordinating residues, iron(II) binds tightly to the low-coordinate site. The iron(II) complex does not react with chemical redox agents to undergo oxidation or reduction. Spectroscopically calibrated quantum-chemical computations show that the complex has high-spin iron(II) in a pseudotetrahedral coordination environment, which features interactions with side chains of two histidines and a cysteine as well as the C=O of Gly45. In the  $^5A_1$  ground state, the  $d_{z^2}$  orbital is doubly occupied. Mutation of Met121 to Ala leaves the metal site in a similar environment but creates a pocket for reversible binding of small anions to the iron(II) center. Specifically, azide forms a high-spin iron(II) complex and cyanide forms a low-spin iron(II) complex.




## INTRODUCTION

Nature uses proteins as scaffolds for controlling transition-metal behavior, and chemists have learned to adapt naturally occurring scaffolds for artificial uses.<sup>1,2</sup> This research effort has uncovered routes to functional and spectroscopic models of metalloenzymes, using both rational design and directed evolution strategies.<sup>3–10</sup> Despite many successes in metalloprotein engineering, artificial complexes containing unsaturated iron sites remain a relatively unexplored area.<sup>5,11–16</sup> Preparation of such complexes could further our understanding of nonheme iron enzymes that activate small molecules,<sup>17–20</sup> broaden the applications of iron chemistry, and facilitate the use of unsaturated iron complexes in a water-soluble scaffold.

In this work, *Pseudomonas aeruginosa* (*Pa*) azurin is used as a scaffold to bind iron in a low-coordinate environment. Azurins are “type 1” copper metalloproteins that are utilized in plants and bacteria for electron transfer, in which the copper cofactor cycles between colorless copper(I) and blue copper(II) forms.<sup>21</sup> Azurins and related copper metalloproteins have received attention in the bioinorganic chemistry community due to their interesting spectroscopy and electronic structure in the oxidized copper(II) form, as well as their ability to transfer electrons rapidly.<sup>22,23</sup> The properties of azurins are a result of the unusual geometric structure of the copper binding site, which is between the residues His46, Cys112, His117, Gly45, and Met121 (*P. aeruginosa* numbering). The two histidines and

the cysteine coordinate to copper in a roughly trigonal-planar geometry, and the sulfur of Met121 (3.1–3.3 Å) and the oxygen of Gly45 (2.6–3.1 Å) are more distant (Figure 1).<sup>24</sup>




**Figure 1.** Metal site of the blue copper protein azurin, indicating changes intended to create an unsaturated iron site.

Descriptions of the coordination number at copper vary depending on whether Met121 and Gly45 are considered as donors or not.<sup>21,22</sup> The binding site is considered rigid, because the residues in the active site have a similar geometry in the apoprotein as in the native copper-bound azurin (Figure 2).<sup>23–26</sup> The copper site of azurin can accommodate alternative

Received: August 22, 2012

Published: November 20, 2012

metals, such as  $Zn^{2+}$ ,  $Ni^{2+}$ ,  $Co^{2+}$ ,  $Mn^{2+}$ ,  $Cd^{2+}$ ,  $Au^+$ ,  $Ag^+$ , and  $Hg^{2+}$ .<sup>27–36</sup>



**Figure 2.** Unoccupied metal site in the X-ray crystal structure of apo-azurin (PDB 1E65).<sup>26</sup>

The ease of preparation and the stability of azurin have made it the target of extensive protein engineering efforts. His46Gly, His117Gly, Met121Ala, and Met121Gly variants have open spaces near their copper sites, and exogenous donors bind at the copper site without changes in the protein fold.<sup>21,37–39</sup> Azurin is a durable scaffold, forming stable copper(II) complexes with up to four mutations in the vicinity of its single metal binding site.<sup>40–49</sup>


Despite success in incorporating other first row transition metals, iron binding to apo-azurin has not been demonstrated. Nonheme iron(II) complexes lack easily detected perpendicular-mode EPR signals and do not have intense visible-region electronic absorption bands, but other spectroscopic techniques can reveal electronic and geometric features of the metal ion in the unusual apo-azurin metal-binding site. In this work, we take advantage of <sup>57</sup>Fe Mössbauer and paramagnetic <sup>1</sup>H NMR spectroscopies, which are combined with quantum-chemical computations to confirm the geometric and electronic structure of the high-spin iron(II) center. We also show that a ligand binding pocket may be introduced, enabling the iron(II) species to bind azide and cyanide at the iron center.

## RESULTS AND DISCUSSION

**Preparation of Iron Complexes of Apo-Azurin.** We expressed recombinant *Pa* azurin and its Met121Ala variant in *E. coli*, using a modified literature procedure (see Experimental Section).<sup>45</sup> Protein expression yielded a mixture of apo-azurin, Cu-azurin, and Zn-azurin. The apo-azurin was separated and purified on an anion exchange column, and gel electrophoresis of the protein is shown in Figure S1 (Supporting Information). In the course of these studies, we learned that even small amounts of zinc in media or containers react with apo-azurin to give significant amounts of zinc(II)-azurin, which does not bind iron(II). Commercial plastic releases significant amounts of zinc(II), as we verified by atomic absorption spectroscopy. Therefore, it was essential to treat all glassware and plasticware with nitric acid or EDTA to remove trace zinc(II), and to remove trace metal ions from buffers and water with Chelex resin. Additionally, it was necessary to use high-grade (99.999% metal analysis) acetate during the purification to obtain apoproteins with zinc content below 10%.


The addition of iron(II) sulfate to buffered solutions of wild-type apo-azurin results in the development of UV bands with

maxima at 322 and 364 nm, which are attributed to the iron(II) complex of apo-azurin (Figure 3). Iron binds to the Met121Ala



**Figure 3.** Growth of bands at 322 and 364 nm after the addition of 1 equiv of  $Fe^{2+}$  to wild type apo-azurin (0.18 mM). (Inset) Time course of the absorbance maxima of the two bands. The solid lines are exponential fits to the absorbances.

azurin apoprotein to give a similar UV-vis spectrum (see Figure 4 and text below); iron(II) binding to the mutant is



**Figure 4.** UV-vis spectra of iron(II) azurins (73  $\mu$ M). (Top) Wild type. (Bottom) Met121Ala variant.

much more rapid than to the wild-type protein, as seen for copper(II) binding.<sup>50</sup> In copper systems, this difference has been attributed to the Met121 residue hindering access to the metal site.<sup>37</sup> The intensity of these bands was the same using 1 equiv or excess iron(II), indicating strong binding of the iron(II) ions in solution (a conservative upper limit is  $K_d < 1 \mu$ M).

If iron(II) binds to the copper site, one expects the presence of iron(II) to influence the copper(II) binding affinity. In the literature, copper(II) binding to apo-azurin is relatively rapid ( $t_{1/2} = 14$  min with 0.01 mM apo-azurin and 100 equiv of  $CuSO_4$  at pH = 5.0), yielding copper(II) azurin with a strong band ( $\lambda_{max} = 625$  nm with  $\epsilon = 5.7 \text{ cm}^{-1}\text{mM}^{-1}$ ) that gives the holo-protein an intense blue color.<sup>21–23</sup> Under our conditions of 0.1 mM apo-azurin and 10 equiv of copper (pH = 7.0),

Table 1. LMCT Bands for Metal-substituted Azurins and Model Complexes

M <sup>2+</sup> -azurin	$\lambda_1$ (cm <sup>-1</sup> ) × 10 <sup>-4</sup> ( $\epsilon$ ) (cm <sup>-1</sup> mM <sup>-1</sup> )	$\lambda_2$ (cm <sup>-1</sup> ) × 10 <sup>-4</sup> ( $\epsilon$ ) (cm <sup>-1</sup> mM <sup>-1</sup> )	model complex	$\lambda_1$ (cm <sup>-1</sup> ) × 10 <sup>-4</sup> ( $\epsilon$ ) (cm <sup>-1</sup> mM <sup>-1</sup> )	$\lambda_2$ (cm <sup>-1</sup> ) × 10 <sup>-4</sup> ( $\epsilon$ ) (cm <sup>-1</sup> mM <sup>-1</sup> )
Fe <sup>2+</sup>	2.75 (1.0)	3.11 (2.4)	Fe <sup>2+</sup> L(SC <sub>6</sub> F <sub>5</sub> ) <sup>a</sup>	2.82 (1.4)	3.08 (5.9)
Fe <sup>2+</sup> (Met121Ala)	2.78 (0.34)	3.16 (1.8)			
Fe <sup>2+</sup> (Met121Ala, N <sub>3</sub> <sup>-</sup> )	2.78 (0.34)	3.16 (1.9)			
Co <sup>2+</sup> <sup>b</sup>	2.68 (1.5)	3.00 (4.0)	Co <sup>2+</sup> L(SC <sub>6</sub> F <sub>5</sub> ) <sup>a</sup>	2.94 (2.9)	3.22(5.0)
Ni <sup>2+</sup> <sup>b</sup>	2.28 (3.3)	2.82 (1.6)	Ni <sup>2+</sup> L(SC <sub>6</sub> F <sub>5</sub> ) <sup>a</sup>	2.30 (1.7)	2.80 (1.9)
Cu <sup>2+</sup> <sup>a</sup>	1.60 (5.8)	NA	Cu <sup>2+</sup> L(SC <sub>6</sub> F <sub>5</sub> ) <sup>a</sup>	1.51 (6.0)	NA
Cu <sup>2+</sup> (Met121Ala) <sup>c</sup>	1.59 (NA)	2.17 (NA)			
Cu <sup>2+</sup> (Met121Ala, N <sub>3</sub> <sup>-</sup> ) <sup>c</sup>	1.92 (NA)	2.44 (NA)			

<sup>a</sup>Reference 57. <sup>b</sup>Reference 54. <sup>c</sup>Reference 37.

development of the characteristic 625-nm band upon addition of copper(II) to apo-azurin occurs with a half-life of less than 5 min and copper(II) binding to the apo-Met121Ala protein is complete in less than 1 min. However, the addition of copper(II) to samples of each iron(II)-substituted azurin (0.1 mM) gives less than 20% copper incorporation after 2 days. Therefore, copper(II) binds more strongly than iron(II) to apo-azurin, but iron(II) greatly hinders copper(II) incorporation, most likely by binding to the same site.

Exchanging the buffer of the iron(II)-substituted protein three times resulted in no significant influence to the UV–vis spectrum, as shown by comparing the ratio of the 322 and 364 nm bands to that of the 280 nm band of the protein. This result indicates that the rate of iron(II) dissociation from the protein is very slow. After buffer exchange and digestion of the iron substituted protein with acid, a colorimetric assay to quantify the amount of iron<sup>51,52</sup> indicated 0.93 Fe/protein for the wild-type protein and 0.78 Fe/protein for the Met121Ala variant.

Iron sites in metalloproteins often function through redox reactions, and most iron(II) complexes in four-coordinate and five-coordinate environments are easily oxidized to iron(III). However, the wild-type iron(II)-azurin was surprisingly resistant to redox reactions. The UV–vis, X-band EPR, and <sup>1</sup>H NMR spectra (see below for spectroscopic signatures) do not change for either iron protein variant after treatment with various oxidizing agents (O<sub>2</sub>, Co<sup>3+</sup>-EDTA, ferricyanide, Ce<sup>4+</sup>) or reducing agents (dithionite, titanium citrate, Eu<sup>2+</sup>-DPTA,<sup>53</sup> DTT, sodium borohydride). The UV–vis spectrum of wild-type iron(II) azurin was also unchanged after heating to 70 °C overnight, and under acid/base conditions as low as pH 5.0 and as high as pH 10.5 for days. Thus, the iron-substituted proteins are quite robust to a wide range of oxidation/reduction potential, temperature and pH.

**UV–Vis Characterization.** The wild-type iron(II)-azurin exhibits UV absorption bands with maxima at  $\lambda = 322$  ( $\epsilon = 2400$  M<sup>-1</sup>cm<sup>-1</sup>) and 364 nm ( $\epsilon = 1000$  M<sup>-1</sup>cm<sup>-1</sup>), whereas the Met121Ala variant gives absorptions that are shifted slightly to 316 ( $\epsilon = 1800$  M<sup>-1</sup>cm<sup>-1</sup>) and 360 nm ( $\epsilon = 340$  M<sup>-1</sup>cm<sup>-1</sup>) (Figure 4). The similarity between the energies of the bands in the two metalloproteins suggests that the Fe<sup>2+</sup> ions have a similar coordination environment. Comparing to other reported metal-substituted azurins,<sup>54</sup> there is a consistent trend (Table 1) in the UV–visible absorption energies Cu<sup>2+</sup> < Ni<sup>2+</sup> < Co<sup>2+</sup> < Fe<sup>2+</sup>. These are assigned to ligand-to-metal charge transfer (LMCT) transitions, which are expected to increase in energy as one moves from right to left across the periodic table as a result of the acceptor *d* orbitals shifting to higher energy. In support of this idea, literature thiolate model complexes of these divalent metals have LMCT transitions with

similar energies to the metalloproteins (Table 2).<sup>55–58</sup> Since the model complexes have a pseudotetrahedral geometry, the

Table 2. Mössbauer Isomer Shift and Electric Quadrupole Splitting for Some Four-, Five- and Six-coordinate High-spin Iron(II) Compounds

compound	ligands	$\delta$ (mm/s)	$\Delta E_Q$ (mm/s)	T (K)	reference
Fe–Az	2N,O,S	0.90	–3.17	80	this work
C9S rubredoxin	4S	0.70	–3.25	4.2	68
C42S rubredoxin	3S,O	0.79	–3.27/ +2.95 <sup>a</sup>	4.2	69
desulferoferredoxin					74, 78
center I	4S	0.69	3.51	4.2	
center II	4N,S	1.04	2.87	4.2	
neelaredoxin	4N,S	1.02	2.80	4.2	79, 80
[Fe <sup>II</sup> (cyclam)(tdt)] <sup>b</sup>	4N,2S	1.00	+3.55	80	75
Fe–Az(M121A)	2N,O,S	0.88	2.90	80	this work
Fe–Az(M121A)–N <sub>3</sub>	3N,O,S	1.02	2.74	80	this work
Fe–Az(M121A)–CN	2N,O,C,S	0.37	0.70	80	this work
[(TPP)Fe–CN] <sup>-c</sup>	4N,C	0.36	1.83	100	81
[(TPP)(MeIm)Fe–CN] <sup>-c</sup>	5N,C	0.37	0.61	100	81
cis-(phen) <sub>2</sub> Fe(CN) <sub>2</sub>	4N,2C	0.18 <sup>d</sup>	0.58	298	82

<sup>a</sup>Two forms were observed. <sup>b</sup>tdt = toluenedithiolate. <sup>c</sup>TPP = tetraphenylporphyrin; MeIm = 1-methylimidazole. <sup>d</sup>The value measured at room temperature is expected to be roughly 0.1 mm/s higher at 80 K due to the second-order Doppler shift.

similarity in the energies of the LMCT bands between the iron(II) azurins and small-molecule pseudotetrahedral iron(II) complexes implies that the iron-substituted azurins also have a similar high-spin iron(II) electronic configuration with a pseudotetrahedral geometry. This idea is consistent with the crystal structures and spectroscopic analysis of other metal substituted azurins (Co<sup>2+</sup>, Ni<sup>2+</sup>, Zn<sup>2+</sup>), which have three strongly coordinating residues (Cys112, His46, His117) as well as binding to Gly45.<sup>21</sup> Below, we use quantum-chemical computations to support and enrich our understanding of the coordination environment.


**Proton NMR Characterization.** The binding of iron(II) to apo-azurin was also evaluated using <sup>1</sup>H NMR spectra of 1–2 mM solutions of the proteins in 10% D<sub>2</sub>O/H<sub>2</sub>O. The addition of 1 equiv of iron(II) to apo-azurin causes the appearance of 10–15 resonances that have chemical shifts as extreme as  $\delta$  70

and  $-40$  ppm. A portion of the spectra in the downfield region is shown below in Figure 9, and the full spectra are shown in Figures S2 and S3 (Supporting Information). Large shifts to the resonance frequencies occur in the vicinity of a paramagnetic metal ion, and the relatively small number of such resonances suggests that the paramagnetic center in iron(II)-azurin lies in a well-ordered position in the protein.<sup>59</sup> The number of paramagnetically shifted peaks in the iron(II) azurin is similar to that previously observed in the  $^1\text{H}$  NMR spectra of cobalt(II) and nickel(II) substituted azurin proteins, which are also high-spin.<sup>60–65</sup> Thus the  $^1\text{H}$  NMR spectra show that the iron(II) ion has a high-spin ( $S = 2$ ) rather than a low-spin ( $S = 0$ ) electronic configuration.<sup>59</sup> There is no detectable EPR spectrum (X-band, perpendicular mode,  $7\text{--}77$  K), consistent with an integer-spin system. Many of the peaks in the paramagnetic  $^1\text{H}$  NMR spectra of the wild type and Met121Ala iron substituted proteins have similar chemical shifts, and the UV CD spectra of the two iron-azurin complexes are indistinguishable (Figure S4, Supporting Information), suggesting that there are not significant conformational differences between the two species.

**Mössbauer Characterization.** Addition of 1 equiv of  $^{57}\text{FeSO}_4$  to the wild type and Met121Ala apo-azurin proteins gave  $^{57}\text{Fe}$ -azurin complexes for Mössbauer spectroscopy. The zero-field Mössbauer spectrum of the wild-type azurin complex with iron(II) at 80 K shows a single quadrupole doublet with high isomer shift and large quadrupole splitting ( $\delta = 0.90$  mm/s,  $|\Delta E_Q| = 3.17$  mm/s, Figure 5, top trace). The parameters are typical for iron(II) in a high-spin electronic configuration, and inconsistent with iron(III) or with low-spin iron(II).<sup>66,67</sup>


The isomer shift of iron azurin is significantly higher than that of the tetrahedral  $\text{Fe}^{2+}\text{S}_4$  sites found in rubredoxin,<sup>68,69</sup> desulfoferritin,<sup>70</sup> and the valence-localized  $[\text{2Fe-2S}]^{1+}$  clusters,<sup>71,72</sup> which have  $\delta \approx 0.7$  mm/s (see also Table 2). Replacement of a cysteine sulfur by a serine oxygen donor in mutated rubredoxins causes an increase in the isomer shift of about 0.1 mm/s; a similar trend is observed for the effect of two nitrogens at the ferrous site of the reduced Rieske center (ca. 0.2 mm/s compared to ferredoxins).<sup>73</sup> Thus, the isomer shift for iron azurin is as expected for a high-spin iron(II) ion with a mixture of “soft” cysteine ligation and “hard” histidine ligation. On the other hand, the value for iron azurin is lower than that found for the square-pyramidal center II of desulfoferritin,<sup>74</sup> which has four histidine ligands and one cysteine ( $\delta = 1.04$  mm/s), and six-coordinate  $[\text{Fe}^{\text{II}}(\text{cyclam})(\text{tdt})]$  ( $\delta = 1.0$  mm/s).<sup>75</sup> We attribute this difference to a lower coordination number in iron-azurin. However, the isomer shift is not as low as the values found for three-coordinate iron(II)-diketiminato complexes ( $\delta$  0.48–0.90 mm/s).<sup>76,77</sup> In summary, the isomer shift of iron(II) azurin indicates a high-spin ferrous site with four ligands, some of which bind through sulfur.

Applied-field measurements revealed that the sign of the quadrupole splitting in iron(II) azurin is negative. The quadrupole splitting of ferrous high-spin complexes is usually dominated by a large valence contribution arising from the  $3d^6$  configuration with one doubly occupied  $d$  orbital. The negative sign of the electric field gradient found for iron(II) azurin is consistent with an anisotropic valence charge distribution arising from a low-lying, doubly occupied  $d_z^2$  orbital, and inconsistent with the  $\beta$  electron lying in a  $d_{x^2-y^2}$  or  $d_{xy}$  orbital. This interpretation is consistent with quantum-chemical computations on the active site (see below).



**Figure 5.** Zero-field Mössbauer spectra of frozen 1 mM solutions of (top) wild-type  $^{57}\text{Fe}$ -azurin, (middle)  $^{57}\text{Fe}$ -Met121Ala azurin, and (bottom)  $^{57}\text{Fe}$ -Met121Ala azurin + 100 mM  $\text{N}_3^-$ . The red lines are fits to the data, using the parameters given in the text. All spectra were recorded at 80 K.

Magnetic Mössbauer measurements with fields of 1–7 T applied perpendicular to the  $\gamma$ -rays revealed weak magnetic hyperfine splitting for iron(II) azurin (Figure 6). The



**Figure 6.** Magnetic Mössbauer spectra recorded at 4.2 and 80 K with fields of 1, 4, and 7 T applied perpendicular to the  $\gamma$ -rays. The red line is a spin Hamiltonian simulation for  $S = 2$  with  $D = 14.3$   $\text{cm}^{-1}$ ,  $E/D = -0.37$  (corresponds to  $D = -15.1$   $\text{cm}^{-1}$  and  $E/D = 0.3$  in the usual convention  $0 \leq E/D \leq 1/3$ , obtained by rotations with Euler angles  $\alpha = 90^\circ$ ,  $\beta = 90^\circ$ ),  $A/g_N\beta_N = (-4.9, -5.4, -60.7)$  T,  $\Delta E_Q = -3.19$  mm/s,  $\eta = 0.7$  (efg rotated by  $\alpha = 19^\circ$ ,  $\beta = 12^\circ$ ),  $\delta = 0.93$  mm/s.


corresponding internal field  $\vec{B} = -\vec{A}\langle\vec{S}\rangle$  indicates low and very anisotropic spin expectation values for the ground state, due to large zero-field splitting (zfs) of the  $S = 2$  spin manifold of the ferrous iron. From spin-Hamiltonian simulations of the magnetic spectra a set of almost rhombic zero-field splitting (ZFS) parameters  $D = -15.1 \text{ cm}^{-1}$  and  $E/D = 0.3$  was found.

Attempts to keep the  $D$  tensor, the electric-field gradient (EFG) and the  $A$  tensor collinear in the simulations did not yield acceptable fits. The best result was obtained with  $D$  being rotated around  $z$ - and  $y$ -axes by  $\alpha = 90^\circ$ ,  $\beta = 90^\circ$  with respect to the hyperfine coupling tensor, which has components  $A/g_N\beta_N = (-4.9, -5.4, -60.7) \text{ T}$ . The EFG tensor was also slightly rotated with respect to  $A$  by  $\alpha = 19^\circ$ ,  $\beta = 12^\circ$ , with the asymmetry parameter being  $\eta = 0.7$ . In this model the spin system has easy magnetization for fields along the  $xy$  plane. Since the  $x$ - and  $y$ -components of  $A$  are small, this explains the weak magnetic splitting. The ZFS values,  $D$  and  $E/D$ , are significantly larger than those of tetrahedrally coordinated iron in rubredoxin or its C9S and C42S mutants.<sup>68,69</sup>

The zero-field Mössbauer parameters for the  $^{57}\text{Fe}$ -substituted Met121Ala variant were similar, displaying  $\delta = 0.88 \text{ mm/s}$  and  $\Delta E_Q = 2.90 \text{ mm/s}$ . The relatively small changes in both  $\delta$  and  $\Delta E_Q$  imply that the coordination geometry is not significantly changed by the Met121Ala mutation. Thus both variants most likely have four-coordinate iron(II) with a bond to the oxygen atom of Gly45. The crystal structures of cobalt(II)-, nickel(II)-, and zinc(II) substituted azurins show short M-O distances of 2.1–2.5 Å between the metal and the oxygen of Gly45, supporting this idea.<sup>27–29</sup>

**Crystallographic Characterization.** Colorless crystals of wild-type iron(II) azurin were obtained under conditions established for copper(II) azurin (see Experimental Section). Given the possibility of zinc incorporation, it was important to identify whether iron or zinc is present at the metal-binding site of the azurin protein in the crystal. Diffraction data (2.4–1.8 Å) were collected at X-ray wavelengths of 1.6082 Å and 1.2702 Å, which are on the high-energy side of the K-shell absorption edges for Fe and Zn respectively. Molecular replacement using the coordinates of copper(II) azurin gave refined structures with four independent molecules (A–D) in the asymmetric unit. Only chains B and C displayed significant anomalous diffraction at the active site indicative of metal-ion binding. There is substantially higher anomalous difference density at these metal sites with the Fe-edge data, compared to the Zn-edge data. If Co, Zn, or Cu were the major metal ion present, data from the higher energy (lower wavelength) edge would show greater anomalous diffraction, but this is clearly not the case (see Table S1, Supporting Information).<sup>83</sup> The iron occupancies at the B and C sites were refined to ca. 30% each by comparing the strength of the anomalous diffraction with the expected values (Table S1). Chains A and D do not contain any metal ions in these crystals. Despite the low iron occupancy, it is clear from the X-ray diffraction data that iron binds only to vacant copper sites of the protein. There was no electron density consistent with transition metals elsewhere in the protein.

Overlaying the metal-binding chains B and C of the iron-azurin model and any of the chains of the previously published copper-azurin structure<sup>24</sup> shows no significant difference (Figure 7a). Although the bond lengths could not be precisely quantified due to the low occupancy of the metal-bound form, the metal is more than 2.5 Å from the sulfur of M121, and ~2.6



**Figure 7.** Metal site in the X-ray crystal structure of wild-type iron azurin. (a) Overlay of iron(II) (orange) and copper(II) (blue) azurins. (b) Overlay of chain B of iron(II) azurin (orange) with chain A of apo-azurin (white). The water molecule in apo-azurin is indicated. (c) Overlay of chain B of iron(II) azurin with chain B of apo-azurin.

Å from the oxygen of G45. Thus, there are three firmly bound ligands (H46, H117, C112) and two more distant potential donors (G45, M121), as found in structures of azurin with other metals.<sup>24,25,27–29</sup>

Chains A and D of the iron(II)-azurin structure do not contain a metal ion and display a conformation where His117 shifts from its coordination position and a water molecule occupies the metal-binding site (Figure 7b and c). The structure of the apoprotein contains a similar asymmetric unit with four unique molecules, two which show the shift in His117 and water molecule binding, and two that do not. Overlaying the Fe–Az structure with the structure of the apoprotein<sup>26</sup> reveals that the two proteins have nearly identical conformations in all four pairs of corresponding molecules, with chains A and D of the apoprotein matching chains A and D of the Fe–Az that lack metal ions. Despite the absence of metal, chains B and C of the apoprotein match the Fe-bound conformation of chains B and C in the Fe–Az structure. The consistency in the site conformations between our structure and that of apo-azurin suggests that crystal packing may generate a ligand conformation in chains A and D that is less favorable for iron(II) chelation. We also speculate that the high concentration of acetate (0.1 M) in the crystallization medium may have facilitated loss of iron ions from the protein over time, since iron(II) loss was not observed in other experiments.

**Computational Evaluation of the Active Site.** Quantum mechanics (QM) and hybrid quantum mechanics/molecular mechanics (QM/MM) geometry optimizations were used to characterize details of the iron geometry and electronic structure in the azurin protein. In the QM approach we

Table 3. Bond Distances (Å) in Iron-substituted Azurin<sup>a</sup>

	Fe–O(G45)	Fe–N(H46)	Fe–S(C112)	Fe–N(H117)	Fe–S(M121)
wt, X-ray <sup>b</sup>	2.59 <sup>b</sup>	2.00 <sup>b</sup>	2.29 <sup>b</sup>	2.07 <sup>b</sup>	3.36 <sup>b</sup>
wt, QM	2.12	2.09	2.27	2.05	3.66
wt, QM/MM	2.14 (0.04)	2.09 (0.01)	2.25 (0.01)	2.04 (0.01)	3.48 (0.25)
M121A, QM	2.08	2.06	2.22	2.05	

<sup>a</sup>For the QM/MM calculations, the average bond lengths and their standard deviations (values in parentheses) used five optimized conformations.

<sup>b</sup>The crystallographically determined bond distances should be viewed with caution, due to the low occupancy of the iron-bound form in the crystal structure, and the influence of the crystal lattice on the metal site geometry.

employed a model that included the metal center and the coordinating amino acids. The steric effects of the remaining protein were taken into account by constraining  $C_{\alpha}$  of the amino acids to the crystallographic positions, while the electrostatic effects were modeled by an implicit solvation model. In the QM/MM model, the remainder of the protein was optimized, but at a computationally less demanding MM level.

Relevant structural parameters derived from these calculations are summarized in Table 3. In the wild-type structure, the bond lengths between iron and the in-plane amino acids (H46, C112, H117) are similar for the QM and QM/MM optimized geometries, while larger variations are obtained for the distances to the axial amino acid residues. Optimization of the geometry always led to a tetrahedral iron with an Fe–O(G45) bond (Figure S5, Supporting Information). However, the bond to the amide oxygen was weak, and the energy varied only 1 kcal/mol with a 0.25 Å change in Fe–O distance (Table S2, Supporting Information). The long Fe–S(M121) distance of >3 Å and the large variation of this distance in the QM/MM optimized structures indicates that there is no Fe–S bond. In addition, QM computations on the M121A mutant gave the same coordination geometry as the wild-type protein. The pseudotetrahedral geometry is consistent with the Mössbauer data above. Due to the similarity of the QM and QM/MM structural parameters, all subsequent property calculations were performed on the QM optimized models, unless indicated otherwise.

In order to compare the computational model with the experimental Mössbauer parameters, we calculated the quadrupole splittings<sup>84</sup> for iron of both the wild-type and M121A mutant. These reproduce the experimental values with high accuracy (Table 4). The contributions to the difference

Table 4. Calculated Mössbauer Parameters<sup>a</sup>

	$\delta$ (mm/s)	$\Delta E_Q$ (mm/s)
wt	0.70 (0.90)	–3.15 (–3.17)
wt + Ala121	0.69	–3.09
mutant	0.68 (0.88)	–2.98 (–2.90)

<sup>a</sup>Experimental values are in parentheses.

between the mutant and the wild-type values were quantified by calculating the quadrupole splitting for an unrelaxed M121A mutant of the wild-type structure (“wt + Ala121” in Table 4). Assuming additivity, approximately two-thirds of the difference can thus be attributed to geometry relaxation between the two forms, while one-third is attributable to the mutation *per se*. However, it should be borne in mind that there is only a minor change in the iron geometry and quadrupole splitting upon mutation.

The isomer shifts were also calculated from the electron density at the iron nucleus using a literature correlation,<sup>85</sup> and they reproduce the slight experimental change of –0.02 mm/s in isomer shift upon mutation. Surprisingly, there was a systematic deviation of 0.20 mm/s from the experimental values. A possible cause for this discrepancy could be that the regression parameters from the literature fit are not transferable to the unusual coordination environment in this protein.

Multiconfigurational CASSCF and NEVPT2 *ab initio* calculations were carried out on the QM optimized geometries to model the zero field splitting parameters. (At the DFT level of theory the calculated  $D$  value was in poor agreement with experimental data (Table S3, Supporting Information), as seen in other systems.<sup>86–89</sup>) The spin–spin coupling (SSC) and spin–orbit coupling (SOC) contributions were calculated using quasi-degenerate perturbation theory (QDPT), and the  $D$  and  $E$  values were subsequently calculated using effective Hamiltonian theory.<sup>90</sup> The individual contribution of each excited state to  $D^{\text{SOC}}$  was obtained using second-order perturbation theory.<sup>91</sup>

The calculated ground state is  $^5A_1$ , as expected for a tetrahedron that is flattened along the  $z$ -axis. As depicted in Figure 8, the doubly occupied orbital is  $d_{z^2}$ . In the wild type




Figure 8. Metal  $d$ -based molecular orbitals for the wild-type iron(II)-azurin from a CASSCF calculation. The right side of the figure shows single excitations that lead to spin–orbit coupling, analyzed under approximate  $D_{2d}$  symmetry.

iron-azurin, the dominant SOC contribution to the ZFS comes from a low lying  $^5B_1$  excited state that results from spin-conserving single excitation (Table 5). The CASSCF(6,5) calculation predicts a SOC contribution of  $-17.1 \text{ cm}^{-1}$ , approximately 87% of the total  $D^{\text{SOC}}$ . Inclusion of dynamic correlation in the NEVPT2 approach results in an increase of

**Table 5. Calculated Transition Energies ( $\text{cm}^{-1}$ ) for the Spin Conserving  $d-d$  Excitations and Their Contributions to the  $D^{\text{SOC}}$  Values Obtained from Second Order Perturbation Theory in the Case of the Wild-type Structure<sup>a</sup>**

	state	CASSCF		NEVPT2	
		trans. energy	$D$ contr.	trans. energy	$D$ contr.
2nd order	${}^5B_1 (d_z^2 \rightarrow d_{x^2-y^2})$	643	-17.1	780	-14.1
	${}^5E (d_z^2 \rightarrow d_{xz})$	3586	1.4	4281	1.1
	${}^5E (d_z^2 \rightarrow d_{yz})$	4812	0.6	5546	0.5
	${}^5B_2 (d_z^2 \rightarrow d_{xy})$	7161	-2.4	8363	-2.0
	$D^{\text{SOC}}$		-19.5		-16.6
QDPT	$D^{\text{SS}}$		0.5		0.5
	$D^{\text{SOC}}$		-16.4		-14.6
	$D$		-16.6		-14.8
	$E/D$		0.25		0.25
	$D$				-15.1
Exp	$D$				-15.1
	$E/D$				0.3


<sup>a</sup> $D^{\text{SS}}$ ,  $D^{\text{SOC}}$ ,  $D$  and  $E/D$  calculated with the effective Hamiltonian method are also presented.

the  ${}^5A_1 \rightarrow {}^5B_1$  transition energy, thus reducing the contribution of the excited state to  $-14.1 \text{ cm}^{-1}$ . The treatment of the spin-orbit coupling to infinite order by means of the QDPT method results in a  $D^{\text{SOC}}$  of  $-14.6 \text{ cm}^{-1}$ , close to the one obtained from second order perturbation theory. Inclusion of the SSC contribution results in a  $D$  value of  $-14.8 \text{ cm}^{-1}$ , in good agreement with the experimental  $D$  value of  $-15.1 \text{ cm}^{-1}$ . Note that the high rhombicity measured experimentally ( $E/D = 0.3$ ) and confirmed computationally ( $E/D \approx 0.25$ ) renders the sign of  $D$  ambiguous.

It is also important to gauge the sensitivity of the spectroscopic parameters to small changes that could be imposed by the protein structure.<sup>92-95</sup> Therefore, we calculated the ZFS parameters for five QM/MM optimized structures that resulted from different low-energy protein conformations. At the NEVPT2 level of theory,  $|D^{\text{SOC}}|$  is consistently predicted at  $11.8 \pm 0.3 \text{ cm}^{-1}$ , despite the significant variations in Fe-Met121 distances in these structures (see above). The minimal variation of the ZFS suggests that Met121 has little influence on the iron electronic structure. Interestingly, substituting the MM point charges from the QM/MM calculation with an implicit COSMO model for the protein also has little influence on the calculated  $|D^{\text{SOC}}|$  value ( $12.9 \pm 0.9 \text{ cm}^{-1}$ ), while a gas phase calculation gave a value of  $17.1 \pm 2.6 \text{ cm}^{-1}$ . So, though the ZFS parameters are sensitive to bulk dielectric effects inside the active site cavity, they are not very sensitive to specific second-sphere influences from the protein.

**Small Molecule Binding.** The Met121Ala mutation was anticipated to create an open site in the axial position of the metal atom. Previous work on copper(II) azurins has demonstrated the binding of exogenous ligands to the metal ion.<sup>21,37-39</sup> To test this hypothesis, we surveyed several potential exogenous ligands that could bind to a coordinatively unsaturated iron(II) center.

The addition of sodium azide to the Met121Ala iron substituted azurin at pH 7 resulted in the immediate appearance of a new set of highly shifted peaks in  ${}^1\text{H}$  NMR spectra (Figures 9 and S6, Supporting Information), suggesting that azide binds rapidly at or near the paramagnetic iron(II) center. With moderate amounts of azide (1 to 100 mM), the resonances for the azide-free form coexist with the new species,



**Figure 9.** Portions of the  ${}^1\text{H}$  NMR spectra of iron(II)-substituted Met121Ala azurin (1.2 mM) with addition of  $\text{NaN}_3$ . After the sequential additions, the azide was removed to generate the bottom spectrum.


indicating that azide exchange is slow on the NMR time scale. At high concentrations of azide ( $[\text{N}_3^-] > 100 \text{ mM}$ ) there was complete conversion to the new species. Though the baseline correction and overlap in the  ${}^1\text{H}$  NMR spectra prevent quantitatively reliable integrations, it is possible to estimate  $K_d \approx 0.02 \text{ M}$ . In control experiments, (a) addition of azide to the wild-type iron azurin resulted in no change to its  ${}^1\text{H}$  NMR spectrum and (b) addition of azide to iron-Met121Ala azurin caused no changes to its UV CD spectrum (Figure S4, Supporting Information), suggesting that there is no conformational change in the protein. These results suggest that, as in the  $\text{Cu}^{2+}$ -Met121Ala azurin,<sup>37,96</sup> the Met121Ala mutant of the  $\text{Fe}^{2+}$ -azurin has a small molecule binding pocket over the metal site, in the space vacated by the methionine residue. In the adduct, the iron(II) ion remains in a high-spin electronic configuration.

The isomer shift in Mössbauer spectra increases upon coordination of additional ligands in high-spin complexes, but is not very sensitive to the precise geometry around the iron atom.<sup>66</sup> Thus it is an ideal method for confirming the idea that added azide coordinates to iron(II). The addition of 100 equiv of sodium azide (120 mM) to  ${}^{57}\text{Fe}^{2+}$ -substituted Met121Ala azurin resulted in a shift in  $\delta$  from 0.88 to 1.02 mm/s and  $\Delta E_Q$  from 2.90 to 2.74 mm/s (Figure 5, see above). The significant change of 0.14 mm/s in the isomer shift is consistent with coordination of an additional ligand, and is too large to explain through a conformational change. The higher isomer shift also suggests that the metal remains coordinated to Gly45.

Potassium cyanide was tested as another potential ligand for iron(II), and these studies were done above pH = 10.0 (100 mM CAPS buffer) to avoid excessive formation of HCN. CD spectra (Figure S4, Supporting Information) verified that there is no significant structural change in the protein at this pH. The addition of KCN to iron-substituted Met121Ala azurin resulted in a rapid,  $[\text{KCN}]$ -dependent loss of the paramagnetically shifted signals in the  ${}^1\text{H}$  NMR spectrum (Figure S7, Supporting Information). The UV spectrum was masked by large UV peaks from the added KCN. After the addition of 50 mM of potassium cyanide (50 equiv relative to protein), no paramagnetically shifted signals were detectable in the  ${}^1\text{H}$  NMR spectrum. When the excess cyanide was removed from the solution via repeated buffer exchanges, the UV-vis and  ${}^1\text{H}$  NMR spectra returned, and were nearly identical to the original iron-substituted Met121Ala azurin.<sup>97</sup> The absence of paramagnetically shifted  ${}^1\text{H}$  NMR signals in the adduct indicates that the iron(II) ion is low-spin in the iron azurin cyanide complex. Mössbauer spectroscopy was used to support this

conclusion. Addition of cyanide to  $^{57}\text{Fe}^{2+}$ -substituted Met121-Ala azurin gave partial conversion to a new species with Mössbauer parameters of  $\delta = 0.37$  mm/s and  $\Delta E_{\text{Q}} 0.70$  mm/s (Figure S8, Supporting Information). These values are drastically different from the high-spin iron(II) complexes above and similar to those previously observed for low-spin iron(II) monocyano complexes (Table 2).<sup>81,82</sup>

To gain further insight into the cyanide binding, FTIR spectra were collected on iron(II)-substituted Met121Ala protein solutions at pH 10 with 5 mM and with 88 mM cyanide ( $\sim 10$  equiv and  $\sim 150$  equiv, respectively) (Figure 10).



**Figure 10.** FTIR spectra of solutions of 1.1 mM Met121Ala iron(II)-azurin in 25 mM MOPS buffer at pH 10, with addition of 5 mM (purple) and 88 mM (red) cyanide. For comparison, the spectra of Met121Ala iron(II) azurin (green) and cyanide (blue dashed) are overlaid on the spectrum. The peak at 2080  $\text{cm}^{-1}$  is due to free cyanide. The peak at 2043  $\text{cm}^{-1}$  at low  $[\text{CN}^-]$  fits to the sum of two Lorentzian peaks (Figure S9, Supporting Information), consistent with a superposition of the residual protein peak at 2048  $\text{cm}^{-1}$  and the C–N stretching band of the cyanide adduct at 2039  $\text{cm}^{-1}$ .

Though there is a small interfering protein band at 2048  $\text{cm}^{-1}$ , the cyanide species has a new  $[\text{CN}^-]$ -dependent band at 2039  $\text{cm}^{-1}$ . This signal, which is roughly 100 times more intense than that for free cyanide, is assigned to the cyanide adduct of Met121Ala iron(II) azurin. This CN stretching frequency is significantly lower than the value of 2080  $\text{cm}^{-1}$  in free cyanide, and the decrease is attributable to backbonding from iron(II) into the  $\pi^*$  orbitals of cyanide. We tentatively attribute this unusually low<sup>81</sup> stretching frequency to the lower coordination number in the iron(II)-azurin, which gives enhanced backbonding as recently described in tetrahedral synthetic iron-carbonyl complexes.<sup>98</sup> However, the need to add excess cyanide to achieve full binding implies that cyanide binding to iron(II) ion within the vacant space created by the Met121Ala mutation is not exceptionally strong (we estimate  $K_{\text{d}}(\text{CN}^-) = 0.001\text{--}0.005$  M based on the NMR and IR titrations). So, although cyanide acts as a strong-field ligand with respect to the ligand-field splitting, it does not bind especially strongly to the iron(II) ion.

Addition of *N*-methylformamide (200 mM), hydrazine (100 mM), or carbon monoxide (1 atm) to the iron(II) complex of Met121Ala apo-azurin resulted in no significant changes to the paramagnetically shifted resonances in its  $^1\text{H}$  NMR spectra. The inability to bind these species could be due to the size, shape, lack of charge, or inherent binding ability of each of

these small molecules. Previous studies on thiolate-bound, low coordinate iron complexes have shown that the variations in binding strength for additional ligands can be attributed to a mixture of steric effects and electronic effects (both  $\sigma$ -donation and  $\pi$ -backbonding).<sup>99</sup>

## CONCLUSIONS

Iron(II) binds strongly to apo-azurin to form a iron(II)-azurin complex that is surprisingly stable to oxidation. Considering that the type 1 copper binding site prefers copper(I) (high  $\text{Cu}^{2+/1+}$  reduction potential),<sup>100</sup> it is likely that the high charge of iron(III) is not compatible with the azurin binding site, and this prevents the oxidation of iron(II). The stability of the iron(II)-protein complex has enabled its characterization using UV-vis,  $^1\text{H}$  NMR, and Mössbauer spectroscopies, as well as X-ray crystallography. These techniques, in combination with computations, give strong evidence for a low-coordinate iron(II) site in which the three biological copper(II) ligands (His46, Cys112, and His117) are accompanied by an interaction with Gly45. There is no significant interaction with the axial Met121. The ferrous site has a high-spin electronic configuration, with the sixth valence electron in the  $d_{z^2}$  orbital.

Removal of a nearby side chain through an Met121Ala mutation does not significantly change the coordination environment or electronic structure of the iron(II) ion, but creates a metal binding site that accommodates azide and cyanide. The azide adduct is high-spin and five-coordinate, but the cyanide adduct is low-spin. Binding of each of these exogenous ligands is reversible. These studies indicate that protein engineering is a viable strategy for introducing a reactive iron(II) binding site into azurin, and suggest that further engineering will lead to novel iron chemistry. In addition, we emphasize the usefulness of Mössbauer spectroscopy and calculations to accurately define the geometry and electronic structure of the iron site in an engineered protein.

## EXPERIMENTAL SECTION

All manipulations involving iron were carried out using standard Schlenk or glovebox techniques under a dinitrogen atmosphere. Unless otherwise noted, solutions were made with water purified with a Barnstead NANOpure DIAMOND Life Sciences (UV/UF) ultrapure water system. Purified water and protein solutions were deoxygenated under vacuum for 12 h and sparged with nitrogen for 15 min before use. Glassware was soaked in 1 M nitric acid and plastics were soaked in 1 M EDTA for 24 h before use. Buffer solutions with a pH above 4 were stirred with Chelex 100 sodium form (Sigma-Aldrich) overnight to remove trace metals. UV-vis measurements were taken with either a Cary 50 Bio UV/vis spectrometer or a HP 8452A Diode Array Spectrophotometer. All buffers were the highest grade available on a per metals basis and purchased from EMD Biosciences, except the acetate buffers which were synthesized from sodium hydroxide, ammonium acetate, and acetic acid purchased from Sigma-Aldrich. The iron(II) sulfate, potassium cyanide, and sodium azide were 98%+ from Sigma-Aldrich. **CAUTION:** Azide and cyanide salts are hazardous. We used azide solutions only at or above pH = 7, and cyanide solutions at or above pH = 10, to avoid exposure to gaseous  $\text{HN}_3$  or HCN.

**Preparation and Purification of Apoazurin.** *Pseudomonas aeruginosa* (Pa) apo-azurin was overexpressed and purified using methods analogous to a previous protocol.<sup>45</sup> The *P. aeruginosa* azurin gene in a pET9a vector, obtained from Professor John H. Richards,<sup>101</sup> was expressed at 25 °C in BL21\*(DE3) *E. coli* (Invitrogen) in 6 L of 2xYT media (16 h, 25 °C). The overexpression of protein was induced with 450 mg of Isopropyl-beta-D thiogalactopyranoside (IPTG) and shaken for an additional 4 h at 25 °C. The cells were collected by



centrifugation (9000 rpm, 10 min) followed by resuspension in buffer (20% sucrose (w/v), 1 mM EDTA, 30 mM TrisHCl, pH = 8.0, 1 L) and shaken at 4 °C for 40 min. The cells were collected by centrifugation and the supernatant was decanted. The cells were resuspended in a solution composed of 4 mM NaCl and 1 mM DTT (1 L) and shaken vigorously at 4 °C for 10 min. The insoluble cell fragments were collected by centrifugation and the protein was decanted into a cool 2-L Erlenmeyer flask. The pH of the solution was reduced by the dropwise addition of 100 mL of 500 mM acetate buffer at pH = 4.1. The precipitated protein was removed by centrifugation and the supernatant was mixed with SP Sepharose fast flow beads (GE Healthcare) and stirred at 4 °C for 5 min. The beads were collected on a frit and washed with buffer (50 mM acetate buffer, pH = 4.1) until the UV detector had a flat baseline. The contaminating zinc azurin was removed by washing the column with 192 mL of pH = 5.34 acetate buffer. Apo-azurin was eluted with 448 mL of pH = 6.01 acetate buffer. The protein was pure by gel electrophoresis. When titrated with copper(II) sulfate, it showed the expected 625/280 nm ratio of 0.58.<sup>21</sup> The buffer was exchanged into 50 mM 3-(*N*-morpholino)propanesulfonic acid (MOPS) buffer (pH = 7.0) using a PD10 desalting column (GE Healthcare, Sephadex G-25 Medium, volume 13.5 mL). Following this protocol, the zinc content was shown to be less than 7% relative to the protein (see "Zinc Analysis" below). The protein concentration was estimated from the 280 nm absorbance using the literature extinction coefficient of 9.0 mM<sup>-1</sup>cm<sup>-1</sup>.<sup>21</sup>

**Site-directed Mutagenesis.** The pET9a (Kan<sup>r</sup>) plasmid containing the azurin gene was used as a template for site directed mutagenesis. The Met121Ala mutation was accomplished according to the QuickChange II method (Stratagene, La Jolla, CA) using the following primer in which the mutated site is underlined: 5'-GGG-TCACTCCGCACTGGCGAAAGGTACCCTGACTC-3'

The parental DNA was digested with Dpn1 at 37 °C for 2 h and the final product was transformed into XL1-Blue supercompetent cells (Stratagene, La Jolla, CA). DNA sequencing confirmed the mutations. The plasmids were transformed into BL21(DE3)\* cells (Invitrogen) for expression, and the protein was overexpressed and purified using methods described above.

**Preparation of Iron(II) Azurin.** FeSO<sub>4</sub> or <sup>57</sup>FeSO<sub>4</sub> was dissolved in H<sub>2</sub>O (17 mg in 500 μL, 0.12 M). A stock solution of 20 mM 3-(*N*-morpholino)propanesulfonic acid (MOPS) was prepared by the addition of 419 mg of MOPS to 100 mL water, and adjusted to pH 7.0 with 2 M sodium hydroxide. A solution of apoprotein (89 μL of a 2.82 mM solution) was diluted to 450 μL with the 20 mM MOPS buffer to give a 0.56 mM solution. The iron(II) metalloderivatives were prepared by the addition of 3.1 μL (252 nmol, 1 equiv) of the FeSO<sub>4</sub> solution to the 450 μL solution of apoprotein. Mixtures were left for at least 12 h before purification to allow complete loading. Loading was monitored by observing the UV-vis spectrum: the wild type protein showed absorbance at 322 and 364 nm ( $\epsilon = 2.4, 1.0$  mM<sup>-1</sup>cm<sup>-1</sup>) whereas the Met121Ala mutant had absorbance at 360 and 316 nm ( $\epsilon = 1.8, 0.34$  mM<sup>-1</sup>cm<sup>-1</sup>). The extinction coefficients were determined by relating the absorbance to the estimated Fe-azurin extinction coefficient of 9.0 mM<sup>-1</sup>cm<sup>-1</sup> at 280 nm.<sup>21</sup> Excess iron was removed by washing the protein with buffer on a Centricon concentrator (GE Healthcare, 3 kDa filter).

**Iron Determination.** Iron concentrations were determined following a literature procedure.<sup>51</sup> This colorimetric iron determination indicated 93% and 78% iron for the wild type and Met121Ala substituted proteins, respectively. The protein concentrations were calculated from the intensity of the absorbance bands at 280 nm using  $\epsilon = 9.0$  mM<sup>-1</sup>cm<sup>-1</sup>.<sup>21</sup>

**<sup>1</sup>H NMR Spectroscopy of Iron(II) Azurin.** NMR spectra were collected on a Varian Inova 600-MHz spectrometer. The spectra were acquired using a super-WEFT pulse sequence,  $d_1$ -180°- $\Delta$ -90°- $t_a$  where  $\Delta = 21.7$  ms and  $t_a + d_1 = 117$  ms. The spectra were acquired with a spectral width of 120 kHz and processed using 30 Hz of Gaussian line broadening. Delays were selected to optimize water suppression, rather than diamagnetic protein suppression.

**Mössbauer Spectroscopy.** Mössbauer spectra were recorded on a conventional spectrometer with alternating constant acceleration of

the  $\gamma$ -source. The minimum experimental line width was 0.24 mm/s (full width at half-height). The sample temperature was maintained constant in an Oxford Instruments Variox or in an Oxford Instruments Mössbauer-Spectromag cryostat with split-pair magnet system; the latter was used for measurements with applied fields up to 7 T with the field at the sample being oriented perpendicular to the  $\gamma$ -beam. The  $\gamma$ -source (<sup>57</sup>Co/Rh, 1.8 GBq) was kept at room temperature. By using a re-entrant bore tube the  $\gamma$ -source could be positioned inside the gap of the magnet coils at a position with zero field. Isomer shifts are quoted relative to iron metal at 300 K. Magnetic Mössbauer spectra were simulated with the program MX (by E.B.) by diagonalization of the spin Hamiltonian for  $S = 2$ :

$$\hat{H} = g\beta\hat{S} \cdot \vec{B} + D[\hat{S}_z^2 - 1/3S(S+1) + E/D(\hat{S}_x^2 - \hat{S}_y^2)] \quad (1)$$

where  $g$  is the average electronic  $g$  value, and  $D$  and  $E/D$  are the axial zero-field splitting and rhombicity. The hyperfine interaction for <sup>57</sup>Fe was calculated by using the usual nuclear Hamiltonian.<sup>66</sup>

**FTIR Spectroscopy.** Infrared measurements were carried out on solutions of 0.6 mM protein with 100 mM buffer (MOPS at pH 7.0 for N<sub>3</sub><sup>-</sup> or CAPS at pH 10.0 for CN<sup>-</sup>), using a Bruker IFS 66v/s FTIR spectrometer with 2 cm<sup>-1</sup> resolution. The detector was a photovoltaic mercury cadmium telluride (MCT) element. Data recording used the OPUS package (Bruker Optics), while analysis and further processing used MATLAB 7.0 (Mathworks).

**Zinc Analysis.** Zinc concentrations were determined using a Perkin-Elmer AAnalyst 600 atomic absorption spectrometer equipped with longitudinal Zeeman background correction and a transverse heated graphite furnace (Perkin-Elmer Life and Analytical Sciences, Shelton, CT). A PerkinElmer Lumina Zn hollow cathode lamp was used. Spectrometer settings: Wavelength, 213.9 nm; Slit width, 0.7 nm; Zn lamp current, 15 mA; Background correction, Zeeman-effect; Integration time, 5 s; Injection volume, 10 μL.

**Crystallographic Characterization.** Crystals were obtained by hanging-drop vapor diffusion against well solutions of 3.0–3.5 M ammonium sulfate, 0.5 M lithium nitrate, and 0.1 M sodium acetate at pH 5.2–5.4. Iron(II)-azurin (40 mg/mL) was mixed 1:1 (v/v) to a total volume of 2 μL and set up against 500 μL of well solution. Colorless diamond-shaped plates 200 μm in length appeared amidst precipitate within 2–3 weeks. X-ray diffraction data sets were collected at CHESS on beamlines A1 and F2. Data were indexed and scaled with HKL2000<sup>102</sup> and phases were determined by molecular replacement with the Phenix software suite<sup>103</sup> using the coordinates of Cu-Az (PDB code 1AZU) as a probe. The models were fit to the density using XtalView<sup>104</sup> and COOT,<sup>105</sup> and the structures were refined to 2.4 – 1.8 Å resolution with the CNS<sup>106</sup> software suite (Table S6, Supporting Information).

**QM Models.** The initial coordinates for the wild-type model were based on subunit C of the crystallographic structure. The QM model included the iron atom and amino acids G45, H46, C112, H117 and M121. Geometry optimizations were performed with the BP86 functional<sup>107,108</sup> and the segmented all-electron relativistically contracted (SARC) def2-TZVP(-f) basis set.<sup>109,110</sup> Resolution of identity with the def2-TZVP/J basis set<sup>111</sup> was used to speed up the calculation. In addition, relativistic corrections were included by means of the zeroth-order regular approximation (ZORA) Hamiltonian and dispersion effects were modeled with the latest parametrization of Grimme's group.<sup>112</sup> During the geometry optimization the steric and electrostatic effects of the missing protein were approximated by freezing the  $\alpha$  carbons of the amino acids and by using a COSMO model<sup>113</sup> with a dielectric constant of 4, respectively. In the case of the mutant, the initial coordinates were obtained by replacing M121 with an alanine residue. All calculations (geometry optimizations and spectroscopic parameters) were performed with the ORCA program.<sup>114</sup>

**QM/MM Models.** MM simulations were performed with the pDynamo<sup>115</sup> library that provides the MM description of the system and a convenient interface with the ORCA program for the QM treatment. The entire subunit C was included in the calculations, which were limited to the wild-type form.

The protonation state of residues were assigned using PROPKA 3.0,<sup>116</sup> and were verified by visual inspection. The positions of the hydrogen atoms were refined by a short classical energy minimization while constraining the heavy atoms to their crystallographic positions. The system was then “soaked” in a pre-equilibrated TIP3P<sup>117</sup> water box of dimensions 58 × 58 × 62 Å and was neutralized by adding two Na<sup>+</sup> ions. Afterward the system was equilibrated at 298.15 K using Langevin-Verlet molecular dynamics. The electrostatic and Lennard-Jones interactions were calculated using an atom-based force-switching method with inner and outer cutoffs of 8 and 12 Å, respectively. During the equilibration procedure the position of the iron atom and the coordinating amino acids was constrained to the crystallographic coordinates. Five snapshots were extracted for subsequent QM/MM geometry optimizations. For the selected snapshots only the water molecules within 5 Å of the protein were kept, in order to reduce the number of point charges that enter the QM Hamiltonian. The reported geometrical parameters are averaged over the five QM/MM optimized geometries.

The OPLS-AA force field<sup>118</sup> was used to describe the protein, while the iron atom, the C and O atoms from the peptide bond of residues M44 and H46, the side chain of C112, H117 and M121 residues, and the entire G45 and H46 residues were included in the QM region and were treated at the same level of theory as in the previously described QM calculations.

**Calculation of Mössbauer Parameters.** The quadrupole splitting parameter for <sup>57</sup>Fe of the nuclear excited state (*I* = 3/2) is given by

$$\Delta E_Q = \frac{1}{2}e_0^2 Q V_{zz} \sqrt{1 + \frac{1}{3}\eta^2}$$

where  $e_0$  is the elementary charge,  $Q$  is the quadrupole moment of the <sup>57</sup>Fe nucleus and in this study is equal to 0.16 barn, and  $\eta = (V_{yy} - V_{xx})/V_{zz}$  is the asymmetry parameter with the principal component of the electric field gradient (EFG)  $|V_{zz}| \geq |V_{yy}| \geq |V_{xx}|$ . The EFG tensor ( $V$ ) was calculated with the B3LYP functional,<sup>119,120</sup> def2-TZVP(-f) basis set, ZORA approximation and a COSMO model with a dielectric constant of 4.

The isomer shift depends linearly on the electron density at the iron nucleus,  $\rho(0)$ :

$$\delta = \alpha[\rho(0) - C] + \beta$$

where  $\alpha$  and  $\beta$  are parameters determined by a linear regression, and the number  $C$  is a scaling parameter introduced for convenience. Since the density at the nucleus varies with the density functional and basis set used, each combination requires a new calibration procedure. All isomer shift calculations reported in the present paper were done with the B3LYP functional, TZVP basis set and ZORA. For this particular functional/basis set combination the regression parameters are:  $\alpha = -0.312$ ,  $\beta = 4.103$ ,  $C = 13770$ .<sup>85</sup>

**Calculation of Zero-Field Splitting Parameters.** The effects of the zero-field splitting (ZFS) are usually described by the following spin Hamiltonian:

$$\hat{H}_{ZFS} = \hat{S} \cdot \mathbf{D} \cdot \hat{S}$$

where  $\hat{S}$  is the fictitious spin of the ground state and  $\mathbf{D}$  is the ZFS tensor. In the absence of an external magnetic field and for systems with  $S > 1/2$ , the ZFS term lifts the degeneracy of the 2S+1 magnetic sublevels. The ZFS tensor consists of spin–spin coupling (SSC) and spin–orbit coupling (SOC) contributions, the latter being assumed to be dominant for the present systems.

The SOC contributions to the  $\mathbf{D}$  tensor were calculated using both the well-established perturbation treatment<sup>91</sup> and the quasi-degenerate perturbation theory (QDPT). The Breit–Pauli two-electron SOC operator was represented by the spin–orbit mean field (SOMF) approximation.<sup>121,122</sup> In QDPT one starts from approximate solutions to the Born–Oppenheimer Hamiltonian obtained from *ab initio* methods such as CASSCF. A minimal active space consisting of six electrons distributed into five 3d iron based orbitals, that is, CASSCF(6,5), was used. The matrix elements of the SOC operator

were calculated in the basis of all the spin-multiplets obtained from 5 quintet, 18 triplet and 13 singlet state average CASSCF roots. Diagonalization of the SOC matrix yields the energy levels and the eigenvectors from which the entire  $\mathbf{D}$  tensor was then extracted by employing the effective Hamiltonian approach.<sup>90</sup> The SSC contributions were estimated using the Breit–Pauli spin–spin Hamiltonian in conjunction with first-order perturbation theory.<sup>123</sup>

In addition, the influence of dynamic correlation on the calculated ZFS values was verified with second-order N-electron valence perturbation theory (NEVPT2) calculations<sup>124,125</sup> on top of the state-averaged CASSCF reference wave functions. All calculations used the def2-TZVP(-f) basis set, and were performed either in gas phase, in the presence of MM point charges or with an implicit COSMO model with a dielectric constant of 4 (see main text for details).

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

NMR spectra and crystallographic details. Coordinates for iron azurin are deposited in the Protein Data Bank under accession number 4HZ1. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

[holland@chem.rochester.edu](mailto:holland@chem.rochester.edu)

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was supported by the National Institutes of Health (GM-065313 to P.L.H. and TM-GM-008267 to T.M.P.), the Fulbright Foundation (P.L.H.), the Alexander von Humboldt Foundation (H.S.S.), the Max-Planck-Gesellschaft (F.N.), and the National Science Foundation (CHE-0749997 to B.R.C.). We thank Robert Gelein for atomic absorption analysis, Scott Kennedy and Sally Rocks for assistance with <sup>1</sup>H NMR spectroscopy, and Thomas Darrah and Robert Poreda for ICP/MS analysis. We thank Kara Bren and her students (especially Ravinder Kaur) and Yi Lu and his students (especially Nicholas Marshall and Nathan Sieracki) for assistance with protein expression and purification. We thank Wolfgang Lubitz for the use of the FTIR spectrometer. We thank John Richards and Yi Lu for providing plasmids.

## ■ REFERENCES

- (1) Lu, Y. *Angew. Chem., Int. Ed.* **2006**, *45*, 5588.
- (2) Pordea, A.; Ward, T. R. *Chem. Commun.* **2008**, 4239.
- (3) Lu, Y.; Berry, S. M.; Pfister, T. D. *Chem. Rev.* **2001**, *101*, 3047.
- (4) Steinreiber, J.; Ward, T. R. *Coord. Chem. Rev.* **2008**, *252*, 751.
- (5) Ueno, T.; Abe, S.; Yokoi, N.; Watanabe, Y. *Coord. Chem. Rev.* **2007**, *251*, 2717.
- (6) Ward, T. R. *Acc. Chem. Res.* **2010**, *44*, 47.
- (7) Brustad, E. M.; Arnold, F. H. *Curr. Opin. Chem. Biol.* **2011**, *15*, 201.
- (8) Reetz, M. T. *Angew. Chem., Int. Ed.* **2011**, *50*, 138.
- (9) Lu, Y.; Garner, D. K.; Zhang, J.-L. Artificial metalloproteins: design and engineering. In *Encyclopedia of Chemical Biology*; Wiley: New York, 2009; Vol. 1, p 124.
- (10) Tracewell, C. A.; Arnold, F. H. *Curr. Opin. Chem. Biol.* **2009**, *13*, 3.
- (11) Bell, C. B.; Calhoun, J. R.; Bobyr, E.; Wei, P.-P.; Hedman, B.; Hodgson, K. O.; DeGrado, W. F.; Solomon, E. I. *Biochemistry* **2008**, *48*, 59.
- (12) Benson, D. E.; Wisz, M. S.; Hellinga, H. W. *Proc. Natl. Acad. Sci.* **2000**, *97*, 6292.

- (13) Benson, D. E.; Wisz, M. S.; Liu, W.; Hellinga, H. W. *Biochemistry* **1998**, *37*, 7070.
- (14) Lin, Y.-W.; Yeung, N.; Gao, Y.-G.; Miner, K. D.; Lei, L.; Robinson, H.; Lu, Y. *J. Am. Chem. Soc.* **2010**, *132*, 9970.
- (15) Lin, Y.-W.; Yeung, N.; Gao, Y.-G.; Miner, K. D.; Tian, S.; Robinson, H.; Lu, Y. *Proc. Natl. Acad. Sci.* **2010**, *107*, 8581.
- (16) Yeung, N.; Lin, Y.-W.; Gao, Y.-G.; Zhao, X.; Russell, B. S.; Lei, L.; Miner, K. D.; Robinson, H.; Lu, Y. *Nature* **2009**, *462*, 1079.
- (17) Wallar, B. J.; Lipscomb, J. D. *Chem. Rev.* **1996**, *96*, 2625.
- (18) Hegg, E. L.; Que, L. *Eur. J. Biochem.* **1997**, *250*, 625.
- (19) Burgess, B. K.; Lowe, D. J. *Chem. Rev.* **1996**, *96*, 2983.
- (20) Holland, P. L. Nitrogen Fixation. In *Comprehensive Coordination Chemistry II*; McCleverty, J., Meyer, T. J., Eds.; Elsevier: Oxford, 2004; Vol. 8, p 569.
- (21) Kolczak, U.; Dennison, C.; Messerschmidt, A.; Canters, G. W. Azurin and Azurin Mutants. In *Handbook of Metalloproteins*; Wiley: New York, 2001; Vol. 2, p 1170.
- (22) Lu, Y. Cupredoxins. In *Comprehensive Coordination Chemistry 2*; McCleverty, J., Meyer, T. J., Eds.; Elsevier: Oxford, 2004; Vol. 8.
- (23) Gray, H. B.; Malmström, B. G.; Williams, R. J. P. *J. Biol. Inorg. Chem.* **2000**, *5*, 551.
- (24) Crane, B. R.; Di Bilio, A. J.; Winkler, J. R.; Gray, H. B. *J. Am. Chem. Soc.* **2001**, *123*, 11623.
- (25) Nar, H.; Messerschmidt, A.; Huber, R.; Van de Kamp, M.; Canters, G. W. *J. Mol. Biol.* **1991**, *221*, 765.
- (26) Nar, H.; Messerschmidt, A.; Huber, R.; Van de Kamp, M.; Canters, G. W. *FEBS Lett.* **1992**, *306*, 119.
- (27) Nar, H.; Huber, R.; Messerschmidt, A.; Filippou, A. C.; Barth, M.; Jaquinod, M.; Van de Kamp, M.; Canters, G. W. *Eur. J. Biochem.* **1992**, *205*, 1123.
- (28) Moratal, J. M.; Romero, A.; Salgado, J.; Perales-Alarcon, A.; Jimenez, H. R. *Eur. J. Biochem.* **1995**, *228*, 653.
- (29) Bonander, N.; Vanngard, T.; Tsai, L.-C.; Langer, V.; Nar, H.; Sjoelin, L. *Proteins* **1997**, *27*, 385.
- (30) Blaszk, J. A.; Ulrich, E. L.; Markley, J. L.; McMillin, D. R. *Biochemistry* **1982**, *21*, 6253.
- (31) Engeseth, H. R.; McMillin, D. R.; Otvos, J. D. *J. Biol. Chem.* **1984**, *259*, 4822.
- (32) Jimenez, H. R.; Salgado, J.; Moratal, J. M.; Morgenstern-Badarau, I. *Inorg. Chem.* **1996**, *35*, 2737.
- (33) Klemens, A. S.; McMillin, D. R.; Tsang, H. T.; Penner-Hahn, J. E. *J. Am. Chem. Soc.* **1989**, *111*, 6398.
- (34) McCleskey, T. M.; Mizoguchi, T. J.; Richards, J. H.; Gray, H. B. *Inorg. Chem.* **1996**, *35*, 3434.
- (35) McMillin, D. R.; Rosenberg, R. C.; Gray, H. B. *Proc. Natl. Acad. Sci.* **1974**, *71*, 4760.
- (36) Wittung-Stafshede, P. *Inorg. Chem.* **2004**, *43*, 7926.
- (37) Tsai, L.-C.; Bonander, N.; Harata, K.; Karlsson, G.; Vaenngaard, T.; Langer, V.; Sjoelin, L. *Acta Cryst. D* **1996**, *52*, 950.
- (38) Canters, G. W.; Gilardi, G. *FEBS Lett.* **1993**, *325*, 39.
- (39) Vidakovic, M.; Germanas, J. P. *Angew. Chem., Int. Ed.* **1995**, *34*, 1622.
- (40) Mizoguchi, T. J.; Di Bilio, A. J.; Gray, H. B.; Richards, J. H. *J. Am. Chem. Soc.* **1992**, *114*, 10076.
- (41) Garner, D. K.; Vaughan, M. D.; Hwang, H. J.; Savelieff, M. G.; Berry, S. M.; Honek, J. F.; Lu, Y. *J. Am. Chem. Soc.* **2006**, *128*, 15608.
- (42) Berry, S. M.; Ralle, M.; Low, D. W.; Blackburn, N. J.; Lu, Y. *J. Am. Chem. Soc.* **2003**, *125*, 8760.
- (43) Hwang, H. J.; Berry, S. M.; Nilges, M. J.; Lu, Y. *J. Am. Chem. Soc.* **2005**, *127*, 7274.
- (44) Hwang, H. J.; Carey, J. R.; Brower, E. T.; Gengenbach, A. J.; Abramite, J. A.; Lu, Y. *J. Am. Chem. Soc.* **2005**, *127*, 15356.
- (45) Hay, M.; Richards, J. H.; Lu, Y. *Proc. Natl. Acad. Sci.* **1996**, *93*, 461.
- (46) Germanas, J. P.; Di Bilio, A. J.; Gray, H. B.; Richards, J. H. *Biochemistry* **1993**, *32*, 7698.
- (47) Marshall, N. M.; Garner, D. K.; Wilson, T. D.; Gao, Y.-G.; Robinson, H.; Nilges, M. J.; Lu, Y. *Nature* **2009**, *462*, 113.
- (48) Lancaster, K. M.; George, S. D.; Yokoyama, K.; Richards, J. H.; Gray, H. B. *Nat. Chem.* **2009**, *1*, 711.
- (49) Lancaster, K. M.; Sproules, S.; Palmer, J. H.; Richards, J. H.; Gray, H. B. *J. Am. Chem. Soc.* **2010**, *132*, 14590.
- (50) Blaszk, J. A.; McMillin, D. R.; Thornton, A. T.; Tennent, D. L. *J. Biol. Chem.* **1983**, *258*, 9886.
- (51) Haigler, B. E.; Gibson, D. T. *J. Bacteriol.* **1990**, *172*, 457.
- (52) Braunschweig, J.; Bosch, J.; Heister, K.; Kuebeck, C.; Meckenstock, R. U. *J. Microbiol. Methods* **2012**, *89*, 41.
- (53) Vincent, K. A.; Tilley, G. J.; Quammie, N. C.; Streeter, I.; Burgess, B. K.; Cheesman, M. R.; Armstrong, F. A. *Chem. Commun.* **2003**, 2590.
- (54) Tennent, D. L.; McMillin, D. R. *J. Am. Chem. Soc.* **1979**, *101*, 2307.
- (55) Matsunaga, Y.; Fujisawa, K.; Ibi, N.; Miyashita, Y.; Okamoto, K.-I. *Inorg. Chem.* **2004**, *44*, 325.
- (56) Chowdhury, A.; Peteanu, L. A.; Holland, P. L.; Tolman, W. B. *J. Phys. Chem. B* **2002**, *106*, 3007.
- (57) Gorelsky, S. I.; Basumallick, L.; Vura-Weis, J.; Sarangi, R.; Hodgson, K. O.; Hedman, B.; Fujisawa, K.; Solomon, E. I. *Inorg. Chem.* **2005**, *44*, 4947.
- (58) Czernuszewicz, R. S.; Fraczkiewicz, G.; Zareba, A. A. *Inorg. Chem.* **2005**, *44*, 5745.
- (59) Ming, L.-J. Nuclear magnetic resonance of paramagnetic metal centers in proteins and synthetic complexes. In *Physical Methods in Bioinorganic Chemistry*; Que, L., Ed.; University Science Books: Sausalito, CA, 2000; p 375.
- (60) Blaszk, J. A.; Ulrich, E. L.; Markley, J. L.; McMillin, D. R. *Biochemistry* **1982**, *21*, 6253.
- (61) Donaire, A.; Salgado, J.; Moratal, J.-M. *Biochemistry* **1998**, *37*, 8659.
- (62) Moratal, J.-M.; Salgado, J.; Donaire, A.; Jimenez, H. R.; Castells, J. *J. Chem. Soc., Chem. Commun.* **1993**, 110.
- (63) Moratal, J. M.; Salgado, J.; Donaire, A.; Jimenez, H. R.; Castells, J.; Martinez-Ferrer, M. J. *Magn. Reson. Chem.* **1993**, *31*, S41.
- (64) Moratal, J. M.; Salgado, J.; Donaire, A.; Jimenez, H. R.; Castells, J. *Inorg. Chem.* **1993**, *32*, 3587.
- (65) Salgado, J.; Jimenez, H. R.; Donaire, A.; Moratal, J. M. *Eur. J. Biochem.* **1995**, *231*, 358.
- (66) Gütlich, P.; Bill, E.; Trautwein, A. X. *Mössbauer Spectroscopy and Transition Metal Chemistry*; Springer Verlag: Berlin, 2011.
- (67) Neese, F.; Petrenko, T. Quantum Chemistry and Mössbauer Spectroscopy. In *Mössbauer Spectroscopy and Transition Metal Chemistry*; Gütlich, P., Bill, E., Trautwein, A. X., Eds.; Springer: Berlin, 2011; p 137.
- (68) Schulz, C. E.; Debrunner, P. G. *J. Phys. Coll.* **1976**, *37*, 153.
- (69) Yoo, S. J.; Meyer, J.; Achim, C.; Peterson, J.; Hendrich, M. P.; Münck, E. *J. Biol. Inorg. Chem.* **2000**, *5*, 475.
- (70) Moura, I.; Huynh, B. H.; Hausinger, R.; Gall, J. L.; Xavier, A. V.; Münck, E. *J. Biol. Inorg. Chem.* **1980**, *255*, 2493.
- (71) Beinert, H.; Holm, R. H.; Münck, E. *Science* **1997**, *277*, 653.
- (72) Dunham, W. R.; Bearden, A. J.; Salmeen, I. T.; Plamer, G.; Sands, R. H.; Orme-Johnson, W. J.; Beinert, H. *Biochim. Biophys. Acta* **1971**, *253*, 134.
- (73) Fee, J. A.; Findling, K. L.; Yoshida, T.; Hille, R.; Tarr, G. E.; Hearshen, D. O.; Dunham, W. R.; Day, E. P.; Kent, T. A.; Münck, E. *J. Biol. Chem.* **1984**, *259*, 124.
- (74) Moura, I.; Tavares, P.; Moura, J. J.; Ravi, N.; Huynh, B. H.; Liu, M. Y.; LeGall, J. *J. Biol. Chem.* **1990**, *265*, 21596.
- (75) Milsman, C.; Patra, G. K.; Bill, E.; Weyhermüller, T.; George, S. D.; Wieghardt, K. *Inorg. Chem.* **2009**, *48*, 7430.
- (76) Andres, H.; Bominaar, E.; Smith, J. M.; Eckert, N. A.; Holland, P. L.; Münck, E. *J. Am. Chem. Soc.* **2002**, *124*, 3012.
- (77) Stoian, S. A.; Vela, J.; Smith, J. M.; Sadique, A. R.; Holland, P. L.; Münck, E.; Bominaar, E. L. *J. Am. Chem. Soc.* **2006**, *128*, 10181.
- (78) Ascenso, C.; Rusnak, F.; Cabrito, I.; Lima, M. J.; Naylor, S.; Moura, I.; Moura, J. J. G. *J. Biol. Inorg. Chem.* **2000**, *5*, 720.

- (79) Jovanovic, T.; Ascenso, C.; Hazlett, K. R. O.; Sikkink, R.; Krebs, C.; Litwiller, R.; Benson, L. M.; Moura, I.; Moura, J. J. G.; Radolf, J. D.; Huynh, B. H.; Naylor, S.; Rusnak, F. *J. Biol. Chem.* **2000**, *275*, 28439.
- (80) Brioukhanov, A. *Appl. Biochem. Microbiol.* **2008**, *45*, 335.
- (81) Li, J.; Noll, B. C.; Schulz, C. E.; Scheidt, W. R. *Angew. Chem., Int. Ed.* **2009**, *48*, 5010.
- (82) Berrett, R. R.; Fitzsimmons, B. W. *J. Chem. Soc. A* **1967**, 525.
- (83) Brennan, S.; Cowan, P. L. *Rev. Sci. Instrum.* **1992**, *63*, 850.
- (84) Neese, F. *Curr. Opin. Chem. Biol.* **2003**, *7*, 125.
- (85) Römelt, M.; Ye, S.; Neese, F. *Inorg. Chem.* **2008**, *48*, 784.
- (86) Maganas, D.; Grigoropoulos, A.; Staniland, S. S.; Chatziefthimiou, S. D.; Harrison, A.; Robertson, N.; Kyritsis, P.; Neese, F. *Inorg. Chem.* **2010**, *49*, 5079.
- (87) Maganas, D.; Sottini, S.; Kyritsis, P.; Groenen, E. J. J.; Neese, F. *Inorg. Chem.* **2011**, *50*, 8741.
- (88) Maganas, D.; Krzystek, J.; Ferentinos, E.; Whyte, A. M.; Robertson, N.; Psycharis, V.; Terzis, A.; Neese, F.; Kyritsis, P. *Inorg. Chem.* **2012**, *51*, 7218.
- (89) Ye, S.; Neese, F. *J. Chem. Theory Comput.* **2012**, *8*, 2344.
- (90) Maurice, R.; Bastardis, R.; deGraaf, C.; Suaud, N.; Mallah, T.; Guihéry, N. *J. Chem. Theory Comput.* **2009**, *5*, 2977.
- (91) Neese, F.; Solomon, E. I. *Inorg. Chem.* **1998**, *37*, 6568.
- (92) Zein, S.; Duboc, C.; Lubitz, W.; Neese, F. *Inorg. Chem.* **2007**, *47*, 134.
- (93) Bühl, M.; Reimann, C.; Pantazis, D. A.; Bredow, T.; Neese, F. *J. Chem. Theory Comp.* **2008**, *4*, 1449.
- (94) Duboc, C.; Ganyushin, D.; Sivalingam, K.; Collomb, M.-N.; Neese, F. *J. Phys. Chem. A* **2010**, *114*, 10750.
- (95) Carmieli, R.; Larsen, T. M.; Reed, G. H.; Zein, S.; Neese, F.; Goldfarb, D. *J. Am. Chem. Soc.* **2007**, *129*, 4240.
- (96) Fraczkiewicz, G.; Bonander, N.; Czernuszewicz, R. S. *J. Raman Spectrosc.* **1998**, *29*, 983.
- (97) This experiment shows that the disappearance of the  $^1\text{H}$  NMR signal is not because cyanide extracts iron from the protein; also note that  $[\text{Fe}(\text{CN})_6]_4^-$  was not observed in our system by Mössbauer or IR spectroscopy.
- (98) Lee, Y.; Peters, J. C. *J. Am. Chem. Soc.* **2011**, *133*, 4438.
- (99) Chiang, K. P.; Barrett, P. M.; Ding, F.; Smith, J. M.; Kingsley, S.; Brennessel, W. W.; Clark, M. M.; Lachicotte, R. J.; Holland, P. L. *Inorg. Chem.* **2009**, *48*, 5106.
- (100) Solomon, E. I.; Szilagy, R. K.; DeBeer, S.; Basumallick, L. *Chem. Rev.* **2004**, *104*, 419.
- (101) Chang, T. K.; Iverson, S. A.; Rodrigues, C. G.; Kiser, C. N.; Lew, A. Y.; Germanas, J. P.; Richards, J. H. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 1325.
- (102) Otwinowski, Z.; Minor, W. *Methods Enzymol.* **1997**, *276*, 307.
- (103) Adams, P. D.; Afonine, P. V.; Bunkóczi, G.; Chen, V. B.; Davis, I. W.; Echols, N.; Headd, J. J.; Hung, L.-W.; Kapral, G. J.; Grosse-Kunstleve, R. W.; McCoy, A. J.; Moriarty, N. W.; Oeffner, R.; Read, R. J.; Richardson, D. C.; Richardson, J. S.; Terwilliger, T. C.; Zwart, P. H. *Acta Cryst. D* **2010**, *66*, 213.
- (104) McRee, D. J. *Mol. Graph.* **1992**, *10*, 44.
- (105) Emsley, P.; Lohkamp, B.; Scott, W. G.; Cowtan, K. *Acta Cryst. D* **2010**, *66*, 486.
- (106) Brünger, A. T.; Adams, P. D.; Clore, G. M.; DeLano, W. L.; Gros, P.; Grosse-Kunstleve, R. W.; Jiang, J.-S.; Kuszewski, J.; Nilges, M.; Pannu, N. S.; Read, R. J.; Rice, L. M.; Simonson, T.; Warren, G. L. *Acta Cryst. D* **1998**, *54*, 905.
- (107) Becke, A. D. *Phys. Rev. A* **1988**, *38*, 3098.
- (108) Perdew, J. P. *Phys. Rev. B* **1986**, *33*, 8822.
- (109) Weigend, F.; Ahlrichs, R. *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297.
- (110) Pantazis, D. A.; Chen, X.-Y.; Landis, C. R.; Neese, F. *J. Chem. Theory Comp.* **2008**, *4*, 908.
- (111) Weigend, F. *Phys. Chem. Chem. Phys.* **2006**, *8*, 1057.
- (112) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. *J. Chem. Phys.* **2010**, *132*, 154104.
- (113) Klamt, A.; Schuurmann, G. *J. Chem. Soc., Perkin 2* **1993**, 799.
- (114) Neese, F. *WIREs Comput. Mol. Sci.* **2012**, *2*, 73.
- (115) Field, M. J. *J. Chem. Theory Comp.* **2008**, *4*, 1151.
- (116) Olsson, M. H. M.; Søndergaard, C. R.; Rostkowski, M.; Jensen, J. H. *J. Chem. Theory Comp.* **2011**, *7*, 525.
- (117) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. *J. Chem. Phys.* **1983**, *79*, 926.
- (118) Jorgensen, W. L.; Maxwell, D. S.; Tirado-Rives, J. *J. Am. Chem. Soc.* **1996**, *118*, 11225.
- (119) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785.
- (120) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648.
- (121) Hess, B. A.; Marian, C. M.; Wahlgren, U.; Gropen, O. *Chem. Phys. Lett.* **1996**, *251*, 365.
- (122) Neese, F. *J. Chem. Phys.* **2005**, *122*, 034107.
- (123) Ganyushin, D.; Neese, F. *J. Chem. Phys.* **2006**, *125*, 024103.
- (124) Angeli, C.; Cimraglia, R.; Evangelisti, S.; Leininger, T.; Malrieu, J. P. *J. Chem. Phys.* **2001**, *114*, 10252.
- (125) Angeli, C.; Cimraglia, R.; Malrieu, J.-P. *J. Chem. Phys.* **2002**, *117*, 9138.

#### NOTE ADDED AFTER ASAP PUBLICATION

This paper was published ASAP on November 20, 2012. Equation 1 has been corrected. The revised version was posted on November 26, 2012.