

Nickel-Catalyzed Carbon–Carbon Bond-Forming Reactions of Unactivated Tertiary Alkyl Halides: Suzuki Arylations

Susan L. Zultanski^{†,‡} and Gregory C. Fu^{*,†,‡}

[†]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

[‡]Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, United States


Supporting Information

ABSTRACT: The first Suzuki cross-couplings of unactivated tertiary alkyl electrophiles are described. The method employs a readily accessible catalyst ($\text{NiBr}_2\text{-diglyme}/4,4'\text{-di-}tert\text{-butyl-2,2'\text{-bipyridine}}$, both commercially available) and represents the initial example of the use of a group 10 catalyst to cross-couple unactivated tertiary electrophiles to form C–C bonds. This approach to the synthesis of all-carbon quaternary carbon centers does not suffer from isomerization of the alkyl group, in contrast with the umpolung strategy for this bond construction (cross-coupling of a tertiary alkylmetal with an aryl electrophile). Preliminary mechanistic studies are consistent with the generation of a radical intermediate along the reaction pathway.

During the past decade, substantial progress has been accomplished in the development of metal-catalyzed cross-coupling reactions of alkyl electrophiles with organometallic reagents to form C–C bonds.^{1,2} Nearly all reports have focused on couplings of primary and secondary electrophiles; in contrast, there have been only isolated examples of successful reactions with unactivated tertiary electrophiles. In particular, Oshima has described cross-couplings with indenyllithium (Ag catalyst), cyclopentadienylmagnesium (Cu), benzylmagnesium (Ag), allylmagnesium (Co, Cu, and Ag), and allyl- and benzylzinc (Ag) reagents.³

The difficulty in coupling tertiary alkyl electrophiles can be attributed to a variety of factors, including a low propensity to undergo oxidative addition via an S_N2 or a direct-insertion pathway. We and others have demonstrated that, for an array of Ni-catalyzed cross-coupling reactions of unactivated primary and secondary alkyl halides, oxidative addition likely proceeds through an inner-sphere electron-transfer pathway.^{4–6} In view of the relative stability of tertiary radicals, this mechanism could be conducive to reactions of tertiary alkyl electrophiles. Indeed, we recently established that carbon–boron bond formation can be achieved by coupling tertiary halides with diboron reagents.⁷ In this report, we expand the scope of cross-coupling reactions of unactivated tertiary alkyl electrophiles by describing the first method for accomplishing carbon–carbon bond formation with the aid of a group 10 catalyst (eq 1).

Early in this investigation, we decided to focus on organoboron reagents as coupling partners for two primary reasons. First, the Suzuki reaction is the most widely used cross-coupling method for the formation of C–C bonds.⁸ Second, we have



employed organo-substituted 9-borabicyclo[3.3.1]nonane (9-BBN) reagents (in conjunction with an alkoxide activator) as partners in an array of Ni-catalyzed Suzuki couplings of *primary* and *secondary* electrophiles,⁹ and we have further established that the conditions are not highly Brønsted-basic, because of complexation of the alkoxide to the trivalent borane.⁴ The propensity of many tertiary alkyl electrophiles to undergo elimination under basic or acidic conditions is a significant impediment to cross-coupling this family of reaction partners.

When we applied our recently reported method for Ni-catalyzed carbon–boron bond formation with tertiary alkyl bromides to a corresponding Suzuki cross-coupling, we observed essentially no C–C bond formation (eq 2). Similarly, the conditions that we had developed for Suzuki arylations of primary and secondary alkyl bromides were not effective for tertiary bromides (<2% yield).¹⁰

Notwithstanding this result, we continued our pursuit of a method for the synthesis of all-carbon quaternary centers via the cross-coupling of unactivated tertiary halides. In view of the steric demand of tertiary electrophiles, we chose to focus on the use of smaller ligands. Ultimately, we determined that a Ni complex with the ligand 4,4'-di-*tert*-butyl-2,2'-bipyridine (**1**) serves as an effective catalyst for the coupling of 1-bromo-1-methylcyclohexane with phenyl-(9-BBN), furnishing the target compound in 88% yield (Table 1, entry 1). A variety of bioactive compounds, including oxycodone (OxyContin),¹¹ include an all-carbon quaternary center with one aryl substituent.

As illustrated in Table 1, in the absence of $\text{NiBr}_2\text{-diglyme}$, ligand **1**, or LiOt-Bu , little or no Suzuki cross-coupling of the unactivated tertiary alkyl bromide occurs (entries 2–4).

Received: November 29, 2012

Published: January 2, 2013








Table 1. Suzuki Arylation of an Unactivated Tertiary Alkyl Bromide: Effect of the Reaction Parameters




^aThe yields were determined by GC analysis versus a calibrated internal standard and are averages of two experiments. ^b1.7 LiOt-Bu, 1.7 *i*-BuOH.



The use of 2,2'-bipyridine (**2**) or bathophenanthroline (**3**)^{10a} rather than **1** as the ligand leads to a somewhat lower yield (entries 5 and 6). For every other Ni-catalyzed Suzuki reaction of 9-BBN reagents that we have described (couplings of primary and secondary electrophiles), a 1,2-diamine was the ligand of choice;⁹ however, this is not the case for this Suzuki cross-coupling of a tertiary alkyl electrophile (entry 7). The counterion of the base plays a critical role in the efficiency of the coupling process (entry 8). C–C bond formation proceeds in lower yield in solvents such as toluene, cyclohexane, and Et₂O (entries 9–11), and the use of less organoborane or less catalyst leads to less product (entries 12 and 13). Finally, the cross-coupling is not highly moisture-sensitive (entry 14).

We established that this method can be applied to Suzuki reactions of an array of unactivated tertiary alkyl bromides, generating the desired C–C bond in good yield (Table 2).¹² Thus, both 1-bromo-1-methylcyclohexane and *tert*-butyl bromide

Table 2. Suzuki Cross-Couplings of Unactivated Tertiary Alkyl Bromides: Scope with Respect to the Electrophile



^aYields of purified products (averages of two experiments). ^bBecause of the volatility of the product, the yield in parentheses was determined by GC analysis with the aid of a calibrated internal standard.

are suitable substrates (entries 1 and 2), as is 3-bromo-3-ethylpentane, which is significantly more hindered (entry 3). Furthermore, a tertiary cyclobutyl bromide can be cross-coupled (entry 4). Importantly, functional groups such as an aromatic ring, ether, alkene, and primary alkyl chloride are compatible with this method for Suzuki cross-coupling of tertiary alkyl bromides (entries 5–8).

Although this method for cross-coupling of unactivated tertiary alkyl halides is versatile with respect to the electrophile (Table 2), it has limitations with respect to the nucleophile (Table 3).¹³ Specifically, ortho- and para-substituted aryl-(9-BBN) reagents generally do not couple in useful yields; on the other hand, a range of meta-substituted compounds serve as suitable Suzuki cross-coupling partners. Fortunately, this set of products is particularly useful, since they are not available through direct Friedel–Crafts alkylation.


1-Iodoadamantane can also be cross-coupled, generating 1-aryladamantanes, a motif found in a number of pharmaceuticals (eq 3).^{14–16} Furthermore, in a preliminary study, we determined that C–C bond formation can be achieved with an unactivated tertiary alkyl chloride as the electrophile (eq 4; yield determined by GC analysis).¹⁶

A possible mechanism for this Ni-catalyzed Suzuki cross-coupling, illustrated for the reaction of 1-bromo-1-methylcyclohexane with Ph-(9-BBN), is provided at the top of Figure 1.^{4–6} As with our couplings of unactivated secondary alkyl halides, our initial mechanistic observations are consistent with a radical pathway for the oxidative addition of an unactivated tertiary halide to Ni. For example, when we conducted a Suzuki reaction in toluene rather than benzene, we obtained a significant amount of diphenylmethane (eq 5), likely due to H-atom abstraction from toluene by radical B to form a benzyl radical, which then enters the catalytic cycle (bottom of Figure 1).


Table 3. Suzuki Cross-Couplings of Unactivated Tertiary Alkyl Bromides: Scope with Respect to the Nucleophile

entry	Ar	yield (%) ^a
1	- <chem>c1ccccc1</chem>	60
2	- <chem>c1ccccc1</chem> <i>i</i> -Pr	61
3	- <chem>c1ccccc1</chem> Ph	74
4	- <chem>c1ccccc1</chem> OTBS	57
5	- <chem>c1ccccc1</chem> O <i>i</i> -Pr	54

^aYields of purified products (averages of two experiments).



Furthermore, a Ni-catalyzed Suzuki reaction of a single diastereomer of a tertiary alkyl bromide leads to the formation of a mixture of diastereomeric cross-coupling products (eq 6). This observation can also be accommodated by a radical pathway for oxidative addition.



In conclusion, we have developed a method for Suzuki arylation of tertiary alkyl bromides, using commercially available catalyst components ($\text{NiBr}_2\text{-diglyme}$ and $4,4'\text{-di-}tert\text{-butyl-2,2'-bipyridine}$). To the best of our knowledge, this reaction is the




Figure 1. (top) Outline of a possible mechanism for the Ni-catalyzed Suzuki cross-coupling of an unactivated tertiary alkyl bromide. (bottom) Possible pathway for the formation of diphenylmethane when toluene is employed as the solvent (the early steps of the catalytic cycle are as illustrated at the top and have been omitted for simplicity).

first example of the use of organoboron reagents as cross-coupling partners with unactivated tertiary electrophiles as well as the first to employ a group 10 catalyst in cross-couplings of unactivated tertiary alkyl halides to form C–C bonds. In contrast with the umpolung approach to this bond construction (coupling of a tertiary organometallic reagent with an aryl halide), our method does not suffer from isomerization of the alkyl group.¹⁷ Preliminary observations are consistent with a radical intermediate along the pathway of this new cross-coupling process. Additional studies directed at expanding the scope of this reaction and enhancing our understanding of the mechanism are underway.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures and compound characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author
gcfu@caltech.edu

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Support was provided by the National Institutes of Health (National Institute of General Medical Sciences, R01-GM62871) and Merck Research Laboratories (summer fellowship for S.L.Z.). We thank Dr. Alexander S. Dudnik for preliminary observations.

REFERENCES

- (1) For reviews and leading references, see: (a) Glasspoole, B. W.; Cradden, C. M. *Nat. Chem.* **2011**, *3*, 912. (b) Glorius, F. *Angew. Chem., Int. Ed.* **2008**, *47*, 8347. (c) Rudolph, A.; Lautens, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 2656. (d) Terao, J.; Kambe, N. *Acc. Chem. Res.* **2008**, *41*, 1545. (e) Hu, X. *Chimia* **2010**, *64*, 231.
- (2) For a recent application of an asymmetric Negishi cross-coupling of an alkyl electrophile in a total synthesis of carolacton, see: Schmidt, T.; Kirschning, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 1063.
- (3) For examples of cross-couplings (not Heck reactions or other related processes) that provide >50% yield with an unactivated tertiary alkyl halide other than a 1-haloadamantane (which cannot eliminate HX), see: (a) Allylmagnesium (Co): Tsuji, T.; Yorimitsu, H.; Oshima, K. *Angew. Chem., Int. Ed.* **2002**, *41*, 4137. Ohmiya, H.; Tsuji, T.; Yorimitsu, H.; Oshima, K. *Chem.—Eur. J.* **2004**, *10*, 5640. (b) Allyl- and benzylmagnesium (Ag): Someya, H.; Ohmiya, H.; Yorimitsu, H.; Oshima, K. *Org. Lett.* **2008**, *10*, 969. Mitamura, Y.; Someya, H.; Yorimitsu, H.; Oshima, K. *Synlett* **2010**, 309. (c) Cyclopentadienylmagnesium (Cu): Sai, M.; Someya, H.; Yorimitsu, H.; Oshima, K. *Org. Lett.* **2008**, *10*, 2545. (d) Allylmagnesium (Cu): Sai, M.; Yorimitsu, H.; Oshima, K. *Bull. Chem. Soc. Jpn.* **2009**, *82*, 1194. (e) Allyl- and benzylzinc (Ag): Mitamura, Y.; Asada, Y.; Murakami, K.; Someya, H.; Yorimitsu, H.; Oshima, K. *Chem.—Asian J.* **2010**, *5*, 1487. (f) Indenyllithium (Ag): Someya, H.; Yorimitsu, H.; Oshima, K. *Tetrahedron* **2010**, *66*, 5993.
- (4) For leading references to mechanistic studies of Ni-catalyzed Suzuki reactions of unactivated alkyl electrophiles, see: Wilsily, A.; Tramutola, F.; Owston, N. A.; Fu, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 5794.
- (5) For early mechanistic studies of Ni-catalyzed Negishi reactions of unactivated alkyl electrophiles, see: (a) Jones, G. D.; Martin, J. L.; McFarland, C.; Allen, O. R.; Hall, R. E.; Haley, A. D.; Brandon, R. J.; Konovalova, T.; Desrochers, P. J.; Pulay, P.; Vicic, D. A. *J. Am. Chem. Soc.* **2006**, *128*, 13175. (b) Phapale, V. B.; Buñuel, E.; García-Iglesias, M.; Cárdenas, D. J. *Angew. Chem., Int. Ed.* **2007**, *46*, 8790. (c) Lin, X.; Phillips, D. L. *J. Org. Chem.* **2008**, *73*, 3680.
- (6) For discussions and leading references, see: (a) Hu, X. *Chem. Sci.* **2011**, *2*, 1867. (b) Phapale, V. B.; Cárdenas, D. J. *Chem. Soc. Rev.* **2009**, *38*, 1598.
- (7) Dudnik, A. S.; Fu, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 10693.
- (8) For leading references, see: (a) Suzuki, A. *Angew. Chem., Int. Ed.* **2011**, *50*, 6722. (b) *Metal-Catalyzed Cross-Coupling Reactions*; de Meijere, A., Diederich, F., Eds.; Wiley-VCH: New York, 2004. (c) *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E.-i., Ed.; Wiley Interscience: New York, 2002.
- (9) For recent examples and leading references, see ref 4 and: Zultanski, S. L.; Fu, G. C. *J. Am. Chem. Soc.* **2011**, *133*, 15362.
- (10) (a) Ligand = 3: Zhou, J.; Fu, G. C. *J. Am. Chem. Soc.* **2004**, *126*, 1340 (footnote 15: “Under our standard conditions, primary and tertiary alkyl bromides...are not suitable substrates.”). (b) Ligand = amino alcohol: González-Bobes, F.; Fu, G. C. *J. Am. Chem. Soc.* **2006**, *128*, 5360.
- (11) For leading references, see: James, K. S.; Colette, R.; Forbes, K.; Hanks, G. *Palliative Med.* **2011**, *25*, 454.
- (12) Under our standard conditions: (a) when the cross-coupling illustrated in entry 1 of Table 2 was conducted in a capped vial under an atmosphere of air, the desired product was generated in 31% yield; (b) an unactivated primary or secondary alkyl bromide/iodide, 1-
- bromoadamantane, and an acyclic tertiary iodide were not suitable cross-coupling partners; (c) PhBpin, PhB(OH)₂, and PhBF₃K did not afford significant amounts of coupling product (<5% yield).
- (13) Several other meta-substituted aryl-(9-BBN) reagents, including several halogenated compounds, were not effective coupling partners.
- (14) Adapalene (Differin) is an example of a pharmaceutical that includes a 1-aryladamantane.
- (15) For a review of the use of the adamantyl group in medicinal chemistry, see: Lamoureux, G.; Artavia, G. *Curr. Med. Chem.* **2010**, *17*, 2967.
- (16) Essentially no cross-coupling product was observed in the absence of NiBr₂-diglyme.
- (17) For two recent advances and leading references, see: (a) Joshi-Pangu, A.; Wang, C.-Y.; Bischof, M. R. *J. Am. Chem. Soc.* **2011**, *133*, 8478. (b) Lohre, C.; Dröge, T.; Wang, C.; Glorius, F. *Chem.—Eur. J.* **2011**, *17*, 6052.