

# Asymmetric Palladium-Catalyzed Directed Intermolecular Fluoroarylation of Styrenes

Eric P. A. Talbot,<sup>\*,†,‡,§</sup> Talita de A. Fernandes,<sup>‡</sup> Jeffrey M. McKenna,<sup>\*,†</sup> and F. Dean Toste<sup>\*,‡</sup>

<sup>†</sup>Novartis Institutes for Biomedical Research, Horsham, West Sussex, RH12 5AB, United Kingdom


<sup>‡</sup>Department of Chemistry, University of California, Berkeley, California 94720, United States

**S** Supporting Information

**ABSTRACT:** A mild catalytic asymmetric direct fluoroarylation of styrenes has been developed. The palladium-catalyzed three-component coupling of Selectfluor, a styrene and a boronic acid, provides chiral monofluorinated compounds in good yield and in high enantiomeric excess. A mechanism proceeding through a Pd(IV)-fluoride intermediate is proposed for the transformation and synthesis of an sp<sup>3</sup> C–F bond.

Fluorine substituents have become widespread in their use in drugs,<sup>1</sup> their presence in a plethora of agrochemicals<sup>2</sup> and in high-performance materials. The introduction of fluorine affects a multitude of properties, and it is well-known that C–(F)<sub>x</sub> substituents can induce conformational changes, alter pK<sub>a</sub> and dipole moments, increase the lipophilicity, and beneficially alter the oral bioavailability of a drug.<sup>2,3</sup> Given this perspective, their introduction has attracted considerable attention from the synthetic community, and the safe and selective introduction of these moieties into the desired scaffold is key to the further expansion in these areas of research. The asymmetric synthesis of sp<sup>3</sup> C–F bonds<sup>4</sup> is most commonly achieved through α-fluorination of ketones and aldehyde derivatives with the use of organocatalysts,<sup>5</sup> phase transfer catalysis,<sup>6</sup> or by way of ring-opening of strained heterocycles.<sup>7</sup> Few examples of enantioselective transition-metal-catalyzed fluorination exist, and where they do, the metal generally serves as a Lewis acid.<sup>8</sup> Herein, we report a palladium-catalyzed arylation fluorination and its application toward the enantioselective construction of sp<sup>3</sup> C–F bonds.


During the course of our studies on the Selectfluor promoted gold-catalyzed intramolecular aminoarylation of styrene **1a** with *p*-tolylboronic acid,<sup>9</sup> we noted that small amounts of benzylic fluoride **1b** was formed when bimetallic gold–palladium complexes were employed as catalysts. Switching to palladium chloride as a catalyst and employing bipyridine (BiPy) as a ligand allowed for **1b** to be formed as the major product. We envisioned that this product arose from a pathway in which the sulfonamide<sup>10</sup> served as chelating group, rather than a nucleophile, to a high-valent palladium intermediate (eq 1).



We had previously shown that sp<sup>3</sup> C–F occurred from gold(III)-fluorides<sup>11</sup> and Ritter and Sanford have elegantly demonstrated that C–F bonds could be generated from the reductive elimination of high-valent palladium.<sup>12</sup> Therefore, we reasoned that the directing group could control the regioselectivity and stabilize the high-valent metal intermediate in conjunction with the bipyridine ligand, diverting it from an oxidative Heck-type coupling reaction toward C–F bond formation.<sup>13</sup>

On the basis of this hypothesis, and starting from commercially available 2-vinylbenzoic acid, we sought an effective directing group (DG) that could provide the desired control of regiochemistry and product distribution. The PdCl<sub>2</sub>/BiPy-catalyzed reaction of **1a** with *p*-tolylboronic acid and Selectfluor,<sup>14</sup> in wet CH<sub>2</sub>Cl<sub>2</sub><sup>15</sup> containing *tert*-butylcatechol,<sup>16</sup> afforded the benzylic fluoride **1b** in 57% yield (Scheme 1).

## Scheme 1. Screening of Suitable Directing Group<sup>a</sup>



Moreover, we were pleased to observe that under identical conditions, the desired fluorinated product was formed when the DG was 2,6-difluoroaniline, pentafluoroaniline,<sup>17</sup> or 8-aminoquinoline<sup>18</sup> (**2a–4a**, Scheme 1). In contrast, no product was detected with **5a**, and only cyclized nonarylated isoxindole-type products were observed with **6a** and **7a**. Finally, decomposition and trace of product was observed with **8a**, likely a result of competing oxidation of the arylthioether.

Received: December 18, 2013

Published: March 11, 2014

With these results in hand, we decided to focus our investigation using 8-aminoquinoline (AQ) as the directing group. This versatile directing group had proven easy to remove and offered more flexibility to extend the chemistry.<sup>18</sup> Screening of several palladium sources revealed Pd(OAc)<sub>2</sub> as the optimal choice, with Pd(TFA)<sub>2</sub>, PdCl<sub>2</sub>, and PdBr<sub>2</sub> all producing increased amounts of the undesired competing oxidative Heck process (see Supporting Information for product distribution) at the expense of the desired fluoride (Table 1, entries 1–4). *N,N*-

**Table 1. Optimization of Fluoroarylation of Styrene 4a**


entry	catalyst	ligand <sup>c</sup>	solvents	additive <sup>d</sup>	yield 4b <sup>a</sup>
1	PdCl <sub>2</sub>	L1	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.1)	none	54
2	Pd(OAc) <sub>2</sub>	L1	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.1)	none	67
3	PdBr <sub>2</sub>	L1	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.1)	none	54
4	Pd(TFA) <sub>2</sub>	L1	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.1)	none	61
5	Pd(OAc) <sub>2</sub>	–	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.1)	none	0
6	Pd(OAc) <sub>2</sub>	L2	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.1)	none	56
7	Pd(OAc) <sub>2</sub>	L3	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.1)	none	76
8	Pd(OAc) <sub>2</sub>	L3	CH <sub>2</sub> Cl <sub>2</sub>	none	23
9	Pd(OAc) <sub>2</sub>	L3	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.2)	none	82
10	Pd(OAc) <sub>2</sub>	L3	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.2)	P1: 30 mol %	91 (86) <sup>b</sup>
11	Pd(OAc) <sub>2</sub>	L5	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.2)	P1: 50 mol %	82
12	Pd(OAc) <sub>2</sub>	L3	CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O (1/0.2)	P2: 30 mol %	74

<sup>a</sup>Determined by correlation between HPLC-MS and crude <sup>1</sup>H-NMR analysis. <sup>b</sup>Value in parentheses reflects isolated yield. <sup>c</sup>L1: bipyridine; L2: 4,4'-dimethoxy-2,2'-bipyridine, L3: 4,4'-di-*tert*-butyl-2,2'-bipyridine. <sup>d</sup>P1: bis(2-ethylhexyl) hydrogen phosphate; P2: dibenzyl hydrogen phosphate.

Bipyridine-type ligands were vital to achieving the fluorinative reaction manifold, with 4,4'-di-*tert*-butylpyridine proving optimal (entry 7); absence of coordinating ligand produced solely the conventional Heck product (entry 5).<sup>19</sup> Solvent affects were dramatic with methylene chloride being the solvent of choice; no coupling products were formed in toluene, DMF, THF, CH<sub>3</sub>CN, and EtOAc. Water was found to increase the reaction rate, and therefore the ideal solvent consisted of a biphasic mixture of 1.0:0.2 CH<sub>2</sub>Cl<sub>2</sub>:water (entries 7–9). Addition of an organic phosphate (30 mol %) as a phase transfer catalyst further increased the rate of the fluoroarylation and diminished the impurities seen in the reaction.<sup>18k</sup> Under the optimized reaction conditions, the palladium-catalyzed oxidative arylation of styrene 4a furnished fluoride 4b in 86% isolated yield after 24 h at room temperature (entry 10).

Having the optimized conditions in hand, a range of boronic acids<sup>19</sup> were subjected to the oxidative conditions using Selectfluor in CH<sub>2</sub>Cl<sub>2</sub>/water (5/1:v/v) to evaluate the scope of the reaction. As shown in Scheme 2, this protocol was efficient with various arylboronic acids bearing electron-donating groups


**Scheme 2. Substrate Scope for AQ Directing Group<sup>a</sup>**



<sup>a</sup>Yields after chromatographic purification.

such as methyl, methoxy, and amino afforded the corresponding coupling products in good yields. Benzylic fluoride 13 was generated in 45% yield after 24h with 2,6-dimethylphenylboronic acid as the substrate, indicating that steric hindrance has an effect on the reaction. The inductive electron-withdrawing/mesomerically donating chloro group was also well tolerated, and 10 was formed in 81% yield after 13h. Electron-poor ester-substituted boronic acids reacted slowly but still afforded the desired fluorides in modest to good yields. It should also be noted that reaction of *p*-tolylboronic acid and Selectfluor under the optimal conditions with the (*E*)-propenyl amide (R<sub>2</sub> = Me) yielded 20 as a single diastereoisomer. On the other hand, the palladium-catalyzed reaction of alkynyl, alkenyl, or alkyl boronic acids failed to deliver the desired product.<sup>20</sup>


In light of the above results, having defined conditions to control the regiochemistry and product distribution, we undertook the evaluation of a range of different *N,P*- and *N,N*-chiral ligands. Unfortunately, using these ligands resulted in erosion of the yield and slow conversion to the desired product; attempting to heat the reaction failed to increase the conversion. The most encouraging *N,N*-chiral ligand (pyridyl-oxazolidine ligand L\*) in combination with the quinolone-based directing, afforded 4b in good enantiomeric excess (81%; er 90.5:9.5) but at the expense of yield (15%) (Scheme 3). Given the issues that were emerging using the quinolone-based directing group in conjunction with the *N,P*- and *N,N*-chiral ligands, we reinvestigated the requirements of our directing group. To that end, we were encouraged to find that the palladium-catalyzed enantioselective fluoroarylation of the 2,6-difluoroanilide and the 2,4,6-trimethylphenyl derivatives gave a much improved yield without significant changes to perturbing the enantioselectivity. The dramatic improvement was seen with the simplified anilides; an aromatic group with no *ortho*-substituent or even a simple methylamide now gave ee in excess of 90% and with the 4-methoxyanilide directing group giving fluoride 27 in 74% yield

Scheme 3. Substrate Scope for Enantioselective Fluoroarylation<sup>a</sup>

<sup>a</sup>Yields after chromatographic purification and enantioselectivities determined by chiral SFC or HPLC. Absolute Configuration assigned by analogy to X-ray structure of 33.


and 96% ee. Following these results, a range of boronic acids were tested on the 4-methoxyaniline substrate, and pleasingly, desired products (**26–32**) were obtained in good yield and high ee with the exception of the bulky 2,6-dimethylboronic acid.

To demonstrate the potential of this fluorinated compound as a useful precursor for synthesis, deprotection of the aminoquinoline **4b** was achieved using Boc anhydride with DMAP, followed by mild deprotection with LiOH and hydrogen peroxide in water/THF to give the desired acid **34** in 62% yield over two steps (eq 2).<sup>18h</sup> Additionally, the 4-methoxyanilide directing group employed in the enantioselective variant was converted to amide **35** by reaction with ceric ammonium nitrate (CAN) (eq 3).



A proposed mechanism for the reaction is outlined in Scheme 4. As already noted, fluoroarylation was only achieved in the presence of an *N,N*-ligand; thus, the catalytic cycle is initiated through formation of an *N,N*-ligated palladium(II) intermedi-

Scheme 4. Proposed Mechanism for the Directed Fluoroarylation of Styrene



ate.<sup>21</sup> Transmetalation with the boronic acid and thereafter coordination and insertion yields a  $\beta$ -arylated Pd(II) species. The regiocontrol of insertion is provided by the directing group coordination and the electronics of the substrate. The coordination of the directing group and the *N,N*-ligand may also stabilize this intermediate and retard the competing  $\beta$ -hydride elimination process thereby promoting oxidation.<sup>22</sup> Oxidation to the high-valent Pd(IV) intermediate is achieved using Selectfluor. Reductive elimination yields the desired monofluorinated product and the catalytic [N–N–Pd(II)] complex.

In conclusion, using amide-based directing groups, we have developed a palladium-catalyzed fluoroarylation of styrenes. The reaction allows for the synthesis of a range of enantioenriched benzylic fluorides by a three-component coupling of styrenes, Selectfluor, and boronic acids.

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

Experimental procedures and compound characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Authors

jeff.mckenna@novartis.com  
eric.p.talbot@gsk.com  
fdtoste@berkeley.edu

### Present Address

<sup>§</sup>GlaxoSmithKline, Gunnels Wood Road Stevenage Herts SG1 2NY, U.K.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The authors thank the Novartis Education, Diversity and Inclusion (ED&I) office for a presidential postdoctoral fellowship (E.P.A.T.). F.D.T. thanks NIHGMs (RO1 GM073932) for financial support, and T.A.D. is grateful for a postdoctoral fellowship from CAPES. We are grateful to H. Douglas, G.

Browne and C. Page (Novartis) for analytical support. We acknowledge A. DiPasquale (UC Berkeley) for assistance with X-ray crystallography and support from NIH Shared Instrument Grant (S10-RR027172).

## REFERENCES

- (1) (a) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320. (b) Hagmann, W. K. *J. Med. Chem.* **2008**, *51*, 4359. (c) Böhm, H.-J.; Banner, B.; Bendels, S.; Kansy, M.; Kuhn, B.; Müller, K.; Obst-Sander, U.; Stahl, M. *ChemBioChem* **2004**, *5*, 637. (d) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881.
- (2) Jeschke, P. *ChemBioChem* **2004**, *5*, 570.
- (3) (a) Ojima, I. *Fluorine in Medicinal Chemistry and Chemical Biology*; Wiley-Blackwell: Chichester, U.K., 2009. (b) Morgenthaler, M.; Schweizer, E.; Hoffmann-Röder, A.; Benini, F.; Martin, R. E.; Jaeschke, G.; Wagner, B.; Fischer, H.; Bendels, S.; Zimmerli, D.; Schneider, J.; Diederich, F.; Kansy, M.; Müller, K. *ChemMedChem* **2007**, *2*, 1100.
- (4) (a) Ma, J.-A.; Cahard, D. *Chem. Rev.* **2004**, *104*, 6119. (b) Ma, J.-A.; Cahard, D. *Chem. Rev.* **2008**, *108*, PR1. (c) Lectard, S.; Hamashima, Y.; Sodeoka, M. *Adv. Synth. Catal.* **2010**, *352*, 2708. (d) Bobbio, C.; Gouverneur, V. *Org. Biomol. Chem.* **2006**, *4*, 2065. (e) Cahard, D.; Xu, X.; Couve-Bonnaire, S.; Pannecoucke, X. *Chem. Soc. Rev.* **2010**, *39*, 558.
- (5) (a) Cahard, D.; Audouard, C.; Plaquevent, J.-C.; Roques, N. *Org. Lett.* **2000**, *2*, 3699. (b) Shibata, N.; Suzuki, E.; Asahi, T.; Shiro, M. *J. Am. Chem. Soc.* **2001**, *123*, 7001. (c) Ishimaru, T.; Shibata, N.; Horikawa, T.; Yasuda, N.; Nakamura, S.; Toru, T.; Shiro, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 4157. (d) Lozano, O.; Blessley, G.; Martinez del Campo, T.; Thompson, A. L.; Giuffredi, G. T.; Bettati, M.; Walker, M.; Borman, R.; Gouverneur, V. *Angew. Chem., Int. Ed.* **2011**, *50*, 8105. (e) Kwiatkowski, P.; Beeson, T. D.; Conrad, J. C.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2011**, *133*, 1738. (f) Beeson, T. D.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2005**, *127*, 8826. (g) Enders, D.; Hüttl, M. R. M. *Synlett* **2005**, 991. (h) Marigo, M.; Fielenbach, D.; Braunton, A.; Kjærsgaard, A.; Jørgensen, K. A. *Angew. Chem., Int. Ed.* **2005**, *44*, 3703. (i) Steiner, D. D.; Mase, N.; Barbas, C. F., III. *Angew. Chem., Int. Ed.* **2005**, *44*, 3706.
- (6) (a) Rauniyar, V.; Lackner, A. D.; Hamilton, G. H.; Toste, F. D. *Science* **2011**, *334*, 1681. (b) Wang, Y.-M.; Wu, J.; Hoong, C.; Rauniyar, V.; Toste, F. D. *J. Am. Chem. Soc.* **2012**, *134*, 12928. (c) Phipps, R. J.; Hiramatsu, K.; Toste, F. D. *J. Am. Chem. Soc.* **2012**, *134*, 8376. (d) Honjo, T.; Phipps, R. J.; Rauniyar, V.; Toste, F. D. *Angew. Chem., Int. Ed.* **2012**, *51*, 9684. (e) Phipps, R. J.; Toste, F. D. *J. Am. Chem. Soc.* **2013**, *135*, 1268. (f) Shunatona, H. P.; Fruh, N.; Wang, Y.-M.; Rauniyar, V.; Toste, F. D. *Angew. Chem., Int. Ed.* **2013**, *52*, 7724. (g) Woltenshulme, J. R.; Rosenqvist, J.; Lozano, O.; Ilupeju, J.; Engle, K. M.; Pidgeon, G.; Moore, P. R.; Sandford, G.; Gouverneur, V. *Angew. Chem., Int. Ed.* **2013**, *52*, 9697.
- (7) (a) Kalow, J. A.; Doyle, A. G. *Tetrahedron* **2013**, *69*, 5702. (b) Kalow, J. A.; Doyle, A. G. *J. Am. Chem. Soc.* **2011**, *133*, 16001. (c) Kalow, J. A.; Doyle, A. G. *J. Am. Chem. Soc.* **2010**, *132*, 3268. (d) Zhu, J.; Tsui, G. C.; Lautens, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 12353.
- (8) (a) Hintermann, L.; Togni, A. *Angew. Chem., Int. Ed.* **2000**, *39*, 4359. (b) Shibata, N.; Suzuki, E.; Takeuchi, Y. *J. Am. Chem. Soc.* **2000**, *122*, 10728. (c) Hamashima, Y.; Yagi, K.; Takano, H.; Tamás, L.; Sodeoka, M. *J. Am. Chem. Soc.* **2002**, *124*, 14530. (d) Paull, D. H.; Scerba, M. T.; Alden-Danforth, E.; Widger, L. R.; Lectka, T. *J. Am. Chem. Soc.* **2008**, *130*, 17260. (e) Suzuki, T.; Hamashima, Y.; Sodeoka, M. *Angew. Chem., Int. Ed.* **2007**, *46*, 5435. (f) Katcher, M. H.; Sha, A.; Doyle, A. G. *J. Am. Chem. Soc.* **2011**, *133*, 15902. (g) Katcher, M. H.; Doyle, A. G. *J. Am. Chem. Soc.* **2010**, *132*, 17402.
- (9) (a) Ball, L. T.; Lloyd-Jones, G. C.; Russell, C. A. *Chem.—Eur. J.* **2012**, *18*, 2931. (b) Melhado, A. D.; Brenzovich, W. E., Jr.; Lackner, A. D.; Toste, F. D. *J. Am. Chem. Soc.* **2010**, *132*, 8885. (c) Brenzovich, W. E., Jr.; Brazeau, J. F.; Toste, F. D. *Org. Lett.* **2010**, *12*, 4728. (d) Brenzovich, W. E., Jr.; Benitez, D.; Lackner, A. D.; Shunatona, H. P.; Tkatchouk, E.; Goddard, W. A., III; Toste, F. D. *Angew. Chem., Int. Ed.* **2010**, *49*, 5519. (e) Zhang, G.; Cui, L.; Wang, Y.; Zhang, L. *J. Am. Chem. Soc.* **2010**, *132*, 1474. (f) Ball, L. T.; Green, M.; Lloyd-Jones, G. C.; Russell, C. A. *Org. Lett.* **2010**, *12*, 4724.
- (10) Dai, H.-X.; Stepan, A. F.; Plummer, M. S.; Zhang, Y.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 7222.
- (11) Mankad, N. P.; Toste, F. D. *Chem. Sci.* **2012**, *3*, 72.
- (12) (a) Furuya, T.; Ritter, T. *Angew. Chem., Int. Ed.* **2008**, *47*, 5993. (b) Ball, N. D.; Sanford, M. S. *J. Am. Chem. Soc.* **2009**, *131*, 3796. (c) Furuya, T.; Benitez, D.; Tkatchouk, E.; Strom, A. E.; Tang, P.; Goddard, W. A., III; Ritter, T. *J. Am. Chem. Soc.* **2010**, *132*, 3793. (d) For review, see: Engle, K. M.; Mei, T.-S.; Wang, X.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2011**, *50*, 1478.
- (13) (a) Karimi, B.; Behzadnia, H.; Elhamifar, D.; Akhavan, P. F.; Esfahani, F. K.; Zamani, A. *Synthesis* **2010**, 1399. (b) Beccalli, E. M.; Broggini, G.; Martinelli, M.; Sottocornola, S. *Chem. Rev.* **2007**, *107*, 5318. (c) Su, Y.; Jiao, N. *Curr. Org. Chem.* **2011**, *15*, 3362. (d) Wu, T.; Yin, G.; Liu, G. *J. Am. Chem. Soc.* **2009**, *131*, 16354. (e) Wu, T.; Cheng, J.; Chen, P.; Liu, G. *ChemComm.* **2013**, *49*, 8707. (f) Liu, G. *Org. Biomol. Chem.* **2012**, *10*, 6243. (g) Zhao, S.-B.; Becker, J. J.; Gagné, M. R. *Organometallics* **2011**, *30*, 3926. (h) For a platinum-catalyzed asymmetric fluorination, see: Cochrane, N. A.; Nguyen, H.; Gagné, M. R. *J. Am. Chem. Soc.* **2013**, *135*, 628.
- (14) Different electrophilic fluorine sources were tested (*N*-fluorobenzenesulfonamide, *N*-fluoro-2,4,6-trimethylpyridinium triflate, *N*-fluoropyridinium triflate, *N*-fluoropyridinium tetrafluoroborate) without success.
- (15) Li, Z.; Song, L.; Li, C. *J. Am. Chem. Soc.* **2013**, *135*, 4640.
- (16) *tert*-Butylcatechol was used as a radical scavenger to prevent polymerization of starting material and also to negate any radical pathway involving the boronic acid. See Martinez, C.; Muniz, K. *Angew. Chem., Int. Ed.* **2012**, *51*, 7031.
- (17) For selected examples, see: (a) Wasa, M.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2009**, *131*, 9886. (b) Wasa, M.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 3680. (c) Chan, K. S. L.; Wasa, M.; Wang, X.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2011**, *50*, 9081. (d) Zhang, X.-G.; Dai, H.-X.; Wasa, M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 11948. (e) He, J.; Wasa, M.; Chan, K. S. L.; Yu, J.-Q. *J. Am. Chem. Soc.* **2013**, *135*, 3387.
- (18) For review see: (a) Corbet, M.; De Campo, F. *Angew. Chem., Int. Ed.* **2013**, *52*, 2. For selected examples, see: (b) Zaitsev, G.; Shabashov, D.; Daugulis, O. *J. Am. Chem. Soc.* **2005**, *127*, 13154. (c) Reddy, B. V. S.; Reddy, L. R.; Corey, E. J. *Org. Lett.* **2006**, *8*, 3391. (d) Gou, F. R.; Wang, X.-C.; Huo, P.-F.; Bi, H.-P.; Guan, Z.-H.; Liang, Y.-M. *Org. Lett.* **2009**, *11*, 5726. (e) Feng, Y.; Chen, G. *Angew. Chem., Int. Ed.* **2010**, *49*, 958. (f) Ano, Y.; Tobisu, M.; Chatani, N. *J. Am. Chem. Soc.* **2011**, *133*, 12984. (g) Tran, L. D.; Daugulis, O. *Angew. Chem., Int. Ed.* **2012**, *51*, 5188. (h) Gutekunst, W. R.; Gianatassio, R.; Baran, P. S. *Angew. Chem., Int. Ed.* **2012**, *51*, 7507. (i) Nishino, M.; Hirano, K.; Satoh, T.; Miura, Y. M. *Angew. Chem., Int. Ed.* **2013**, *52*, 4457. (j) Shang, R.; Iliés, L.; Matsumoto, A.; Nakamura, E. *J. Am. Chem. Soc.* **2013**, *135*, 6030. (k) Zhang, Z.-Y.; Li, Q.; He, Q.; Nack, W. N.; Chen, G. *J. Am. Chem. Soc.* **2013**, *135*, 12135.
- (19) No coupling product was produced when 2,6-dimethylbipyridine was used as a ligand, suggesting that the steric hindrance may inhibit the coupling reaction. Phosphine ligands (dppm, dppe, and dppf) also failed to give the desired products.
- (20) Aryl boronic ester as well as MIDA boronate were found inactive under the optimized conditions. Slow conversion was observed with aryl trifluoroborate salt. A range of heterocyclic boronic acid were tested without success. See also: Kikushima, K.; Holder, J. C.; Gatti, M.; Stoltz, B. M. *J. Am. Chem. Soc.* **2011**, *133*, 69.
- (21) Yoo, K. S.; Park, C. P.; Yoon, C. H.; Sakaguchi, S.; O'Neill, J.; Jung, K. W. *Org. Lett.* **2007**, *9*, 3933.
- (22) (a) Zhang, Q.; Lu, X.; Han, X. *J. Org. Chem.* **2001**, *66*, 7676. (b) Zhang, Q.; Lu, X. *J. Am. Chem. Soc.* **2000**, *122*, 7604.