

Reconstitution of Peptidoglycan Cross-Linking Leads to Improved Fluorescent Probes of Cell Wall Synthesis

Matthew D. Lebar,^{†,⊥} Janine M. May,^{†,⊥} Alexander J. Meeske,[‡] Sara A. Leiman,[§] Tania J. Lupoli,^{†,‡} Hirokazu Tsukamoto,[†] Richard Losick,[§] David Z. Rudner,[‡] Suzanne Walker,^{*,‡} and Daniel Kahne^{*,†}

[†]Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, United States

[‡]Department of Microbiology and Immunobiology, Harvard Medical School, Boston, Massachusetts 02115, United States

[§]Department of Molecular and Cellular Biology, Harvard University, Cambridge, Massachusetts 02138, United States

Supporting Information

ABSTRACT: The peptidoglycan precursor, Lipid II, produced in the model Gram-positive bacterium *Bacillus subtilis* differs from Lipid II found in Gram-negative bacteria such as *Escherichia coli* by a single amidation on the peptide side chain. How this difference affects the cross-linking activity of penicillin-binding proteins (PBPs) that assemble peptidoglycan in cells has not been investigated because *B. subtilis* Lipid II was not previously available. Here we report the synthesis of *B. subtilis* Lipid II and its use by purified *B. subtilis* PBP1 and *E. coli* PBP1A. While enzymes from both organisms assembled *B. subtilis* Lipid II into glycan strands, only the *B. subtilis* enzyme cross-linked the strands. Furthermore, *B. subtilis* PBP1 catalyzed the exchange of both D-amino acids and D-amino carboxamides into nascent peptidoglycan, but the *E. coli* enzyme only exchanged D-amino acids. We exploited these observations to design a fluorescent D-amino carboxamide probe to label *B. subtilis* PG in vivo and found that this probe labels the cell wall dramatically better than existing reagents.

Bacterial cells are surrounded by a cell wall composed of layers of peptidoglycan (PG). This mesh-like macromolecule stabilizes cell membranes against high internal osmotic pressures and is essential for survival. Peptidoglycan is assembled from the lipid-linked disaccharide pentapeptide precursor Lipid II (Figure 1a, 1).¹ Peptidoglycan glycosyltransferases polymerize Lipid II into glycan strands and enzymes known as penicillin-binding proteins (PBPs) cross-link and process the peptide side chains to produce mature PG.² Because PG is highly conserved and essential for cell survival, it is a target for antibiotics. Indeed, the most successful class of antibiotics in history, the β -lactams, inhibits the transpeptidases (TPs) that cross-link the carbohydrate strands of PG.³

TPs contain a conserved active-site serine that attacks the terminal D-Ala-D-Ala amide bond of the pentapeptide attached to the glycan polymer, forming a covalent enzyme–substrate (E-S) intermediate and releasing D-Ala (Figure 1b, activation step). An amino group on the peptide side chain of another glycan strand can then attack the E-S intermediate to form a cross-link and regenerate the enzyme (Figure 1b, cross-linking




Figure 1. Reactions catalyzed by transpeptidases (TPs) that cross-link peptidoglycan. (a) Structures of Lipid II used by *E. coli* and *B. subtilis*.⁴ (b) Reactions catalyzed by TPs proceed via an enzyme–substrate intermediate that can be resolved by water to generate a tetrapeptide side chain, by a side chain on another glycan strand to yield a cross-link, or by a D-amino acid to give a pentapeptide with a new terminal amino acid.

step). While the structure of Lipid II is highly conserved, the identity of the amino acid containing the reactive amino group

Received: June 5, 2014

Published: July 18, 2014

varies by species.^{1a,5} As the Lipid II substrate is difficult to obtain,^{6,7} it is not well understood how these differences in the stem peptide affect TP cross-linking chemistry.^{8,9}

The canonical Lipid II used by Gram-negative bacteria, including *Escherichia coli*, contains the pentapeptide L-Ala- γ -D-Glu-*m*-DAP-D-Ala-D-Ala (1).¹⁰ The third residue, *m*-DAP, which functions as the nucleophile in the cross-linking reaction, has a D-stereocenter in the side chain. *Bacillus subtilis* Lipid II is similar but contains a carboxamide instead of a carboxylic acid on the *m*-DAP side chain (2).^{11,12} Cell wall precursor 1 has been synthesized previously.^{8d} We synthesized the *B. subtilis* Lipid II substrate 2 in 32 steps by adapting the synthesis of Gram-negative Lipid II^{8d,13} to introduce the carboxamide-containing *m*-DAP residue.¹⁴

Access to both 1 and 2 has allowed us to investigate the substrate preferences of TPs from *E. coli* and *B. subtilis*. *E. coli* PBP1A and *B. subtilis* PBP1¹⁵ were purified as described in the Supporting Information, and each enzyme was then incubated with substrate 1 or substrate 2. These enzymes contain an N-terminal glycosyltransferase domain that polymerizes disaccharide subunits and a C-terminal TP domain that cross-links polymeric glycan strands. To assess cross-linking, we treated reaction mixtures with the glycosylhydrolase mutanolysin followed by NaBH₄ and then used LC/MS to identify the products (Figure 2a).^{8d,16} As expected, each enzyme is able to polymerize and cross-link its native substrate (Figure 2b, traces i and ii). In addition, *B. subtilis* PBP1 is able to cross-link the canonical Gram-negative Lipid II (1, Figure 2b, trace iii), but *E. coli* PBP1A is not able to cross-link *B. subtilis* Lipid II (2, Figure 2b, trace iv), implying that the *E. coli* PBP discriminates against the carboxamide on the *m*-DAP.

We have previously shown that *E. coli* PBP1A can exchange the terminal D-Ala in cell wall precursors with other D-amino acids during in vitro PG synthesis.^{13a,16,17} To further assess the substrate scope of the *E. coli* and *B. subtilis* enzymes, we examined their ability to incorporate either D-Phe or D-Phe carboxamide (D-Phe-NH₂) into synthetic PG (Figure 3a). *E. coli* PBP1A incorporated D-Phe but not D-Phe-NH₂ (Figure 3b, traces i and ii), whereas *B. subtilis* PBP1 incorporated both (Figure 3b, traces iii and iv).¹⁸ Hence, the cross-linking and D-amino acid incorporation experiments are consistent in showing that the *E. coli* PBP1A TP domain discriminates against carboxamide substrates but the *B. subtilis* PBP1 TP domain does not.

A variety of D-amino acid probes have recently been developed to fluorescently label PG in living cells,¹⁹ but labeling is poor in *B. subtilis* unless the cell wall hydrolase DacA, which removes terminal D-amino acids from PG, is absent.^{19a} To determine if D-amino carboxamides can be stably incorporated into PG during *B. subtilis* growth, we grew cells to early log phase in medium supplemented with 500 μ M D-Phe or D-Phe-NH₂ and then analyzed the composition of the pentapeptide in PG fragments following enzymatic degradation. LC/MS analysis showed that D-Phe and D-Phe-NH₂ were incorporated exclusively at the fifth position of the stem peptide, as observed previously for D-amino acids,²⁰ but the levels of D-Phe-NH₂ were much higher (Figure S1). These results led us to examine the utility of fluorescent D-amino carboxamide probes for imaging cell wall biosynthesis in *B. subtilis*.

We prepared previously reported fluorescent probe 3 (FDL)^{19a} as well as new probe 4 (FDL-NH₂) by appending fluorescein to D-Lys and D-Lys carboxamide, respectively




Figure 2. *B. subtilis* transpeptidases (TPs) can cross-link amidated peptidoglycan precursors in vitro. (a) Protocol for detecting cross-linking by *E. coli* and *B. subtilis* TPs in vitro. Fragments A/A' are derived from unmodified polymer; the remaining products are derived from polymer that has been acted on by a TP. (b) LC/MS-extracted ion chromatograms showing unprocessed (A/A'), hydrolyzed (B/B'), and cross-linked products (C/C', D/D') formed in the presence of *E. coli* PBP1A (i,iv) or *B. subtilis* PBP1 (ii,iii) and substrate 1 (i,iii) or 2 (ii,iv).

(Figure 4a). Wild-type *B. subtilis* (dacA+) was grown in medium supplemented with 100 μ M of either 3 or 4 for approximately four generations, and cells were then examined by fluorescence microscopy. FDL-NH₂ 4 efficiently labeled both new septa and the cylindrical walls of the cells. By contrast, FDL 3 inefficiently labeled the cells (Figure 4b). Even FDL-NH₂ concentrations as low as 10 μ M yielded strong labeling (Figure S2). Use of the fluorescent carboxamide probe FDL-NH₂ may be preferable for imaging cell wall synthesis, as it does not require mutations (e.g., *dacA* deletion) that perturb peptidoglycan processing.

To test whether D-amino carboxamides label PG in *E. coli*, we grew cells to early log phase in medium with 500 μ M of either D-Phe or D-Phe-NH₂, harvested the PG, and analyzed the PG fragments using LC/MS as before. As reported previously for D-amino acids,^{20b,21} D-Phe incorporation was detected exclusively in the fourth position of the stem peptide (Figure S3b, trace ii). Strikingly, almost no D-Phe-NH₂ incorporation was observed




Figure 3. *B. subtilis* TPs can incorporate D-amino carboxamides into PG in vitro. (a) Protocol for detecting TP-mediated D-amino acid or carboxamide exchange in vitro. (b) LC/MS-extracted ion chromatograms of products generated by reaction of *E. coli* PBP1A (i,ii) or *B. subtilis* PBP1 (iii,iv) with Lipid II and 1 mM D-Phe (i,iii) or D-Phe-NH₂ (ii,iv). Products E/E' contain D-Phe; product F' contains D-Phe-NH₂.




Figure 4. Fluorescent D-amino carboxamide probes dramatically improve detection of PG synthesis. (a) Structures of fluorescent probes. (b) *B. subtilis* was grown with 100 μM probe to mid log phase. Images were adjusted to the same intensity scale to allow comparison. Scale bars: 1 μm.

(Figure S3b, trace iii). It has previously been suggested that another subset of TPs called the L_D-transpeptidases are responsible for fourth-position incorporation,^{20b} and these data suggest that, like the *E. coli* PBPs, *E. coli* L_D-transpeptidases discriminate against carboxamides. Likewise, *E. coli* grown with 500 μM of either probe 3 or 4 for four generations incorporated 3 but not 4 (Figure S4), indicating that the ability to incorporate D-amino carboxamides may be specific to organisms like *B. subtilis* that cross-link substrates containing an amino group alpha to a carboxamide.

The ability of enzymes from *B. subtilis* to incorporate both D-amino acids and D-amino carboxamides suggests that there may

be two trajectories for nucleophiles to interact with the *B. subtilis* transpeptidase enzyme–substrate intermediate: one for D-amino acids, which may constitute the reverse of the activation step in which a D-amino acid is the leaving group, and the other for forming cross-links to incoming *m*-DAP-NH₂ side chains as well as D-amino carboxamides that resemble these side chains. For *E. coli*, the incoming side chain is essentially a D-amino acid, and other modifications are not well tolerated. The dramatically increased efficiency of incorporation of D-amino carboxamides in *B. subtilis* cells could be due to more efficient incorporation of this nucleophile, less efficient removal by endogenous PG hydrolases once installed, or both. D-amino carboxamide probes may also be useful for studying pathogenic bacteria that contain an amidated *m*-DAP residue in their peptidoglycan, such as *Mycobacterium tuberculosis* and *Corynebacterium diphtheriae*.^{19b,22} Using this general strategy, it may be possible to design probes that mimic cross-linking residues for other bacteria.

■ ASSOCIATED CONTENT

📄 Supporting Information

Experimental procedures, synthesis of substrates and compound analysis, protein purification protocols, and LC/MS analysis of PG polymers. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

✉ Corresponding Authors

suzanne_walker@hms.harvard.edu
kahne@chemistry.harvard.edu

Author Contributions

¹M.D.L. and J.M.M. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by the NIH (R01 GM076710 to S.W.; R01 GM066174 to D.K.; R01 GM073831 to D.Z.R.; R01 GM18568 to R.L.; CETR (U19 AI109764) to D.K., S.W., and D.Z.R.; F32 GM103056 to M.D.L.) and NSF (DGE-1144152 to J.M.M.). All high-resolution LC/MS data were acquired on an Agilent 6520 Q-TOF, spectrometer supported by the Taplin Funds for Discovery Program (P.I.: S.W.).

■ REFERENCES

- (1) (a) Vollmer, W.; Blanot, D.; de Pedro, M. A. *FEMS Microbiol. Rev.* **2008**, *32*, 149–167. (b) Lovering, A. L.; Safadi, S. S.; Strynadka, N. C. J. *Annu. Rev. Biochem.* **2012**, *81*, 451–478. (c) Typas, A.; Banzhaf, M.; Gross, C. A.; Vollmer, W. *Nat. Rev. Microbiol.* **2012**, *10*, 123–136.
- (2) Sauvage, E.; Kerff, F.; Terrak, M.; Ayala, J. A.; Charlier, P. *FEMS Microbiol. Rev.* **2008**, *32*, 234–258.
- (3) (a) Waxman, D. J.; Strominger, J. L. *Annu. Rev. Biochem.* **1983**, *52*, 825–869. (b) Walsh, C. *Antibiotics: Actions, Origins, Resistance*; ASM Press: Washington, DC, 2003.
- (4) A heptaprenyl lipid chain was used rather than the most commonly found natural lipid (undecaprenyl) because the former does not aggregate as extensively and thus is a better substrate for biochemical studies (ref 7d).
- (5) Bouhss, A.; Trunkfield, A. E.; Bugg, T. D. H.; Mengin-Lecreux, D. *FEMS Microbiol. Rev.* **2008**, *32*, 208–233.
- (6) (a) Anderson, J. S.; Matsushashi, M.; Haskin, M. A.; Strominger, J. L. *Proc. Natl. Acad. Sci. U.S.A.* **1965**, *53*, 881–889. (b) Breukink, E.; van Heusden, H. E.; Vollmerhaus, P. J.; Swiezewska, E.; Brunner, L.;

- Walker, S.; Heck, A. J.; de Kruijff, B. *J. Biol. Chem.* **2003**, *278*, 19898–19903. (c) El Ghachi, M.; Bouhss, A.; Barreateau, H.; Touzé, T.; Auger, G.; Blanot, D.; Mengin-Lecreulx, D. *J. Biol. Chem.* **2006**, *281*, 22761–22772. (d) Lloyd, A. J.; Gilbey, A. M.; Blewett, A. M.; De Pascale, G.; El Zoeiby, A.; Levesque, R. C.; Catherwood, A. C.; Tomasz, A.; Bugg, T. D. H.; Roper, D. I.; Dowson, C. G. *J. Biol. Chem.* **2008**, *283*, 6402–6417. (e) Patin, D.; Barreateau, H.; Auger, G.; Magnet, S.; Crouvoisier, M.; Bouhss, A.; Touzé, T.; Arthur, M.; Mengin-Lecreulx, D.; Blanot, D. *Biochimie* **2012**, *94*, 985–990.
- (7) (a) Men, H.; Park, P.; Ge, M.; Walker, S. *J. Am. Chem. Soc.* **1998**, *120*, 2484–2485. (b) Ha, S.; Chang, E.; Lo, M.-C.; Men, H.; Park, P.; Ge, M.; Walker, S. *J. Am. Chem. Soc.* **1999**, *121*, 8415–8426. (c) Schwartz, B.; Markwalder, J. A.; Wang, Y. *J. Am. Chem. Soc.* **2001**, *123*, 11638–11643. (d) Ye, X.-Y.; Lo, M.-C.; Brunner, L.; Walker, D.; Kahne, D.; Walker, S. *J. Am. Chem. Soc.* **2001**, *123*, 3155–3156. (e) Vannieuwenhze, M. S.; Mauldin, S. C.; Zia-Ebrahimi, M.; Winger, B. E.; Hornback, W. J.; Saha, S. L.; Aikins, J. A.; Blaszczyk, L. C. *J. Am. Chem. Soc.* **2002**, *124*, 3656–3660. (f) Shih, H.-W.; Chang, Y.-F.; Li, W.-J.; Meng, F.-C.; Huang, C.-Y.; Ma, C.; Cheng, T.-J. R.; Wong, C.-H.; Cheng, W.-C. *Angew. Chem., Int. Ed.* **2012**, *51*, 10123–10126.
- (8) Reconstitution of cross-linking has only been reported with enzymes from *E. coli* and *S. pneumoniae*, see: (a) Bertsche, U.; Breukink, E.; Kast, T.; Vollmer, W. *J. Biol. Chem.* **2005**, *280*, 38096–38101. (b) Born, P.; Breukink, E.; Vollmer, W. *J. Biol. Chem.* **2006**, *281*, 26985–26993. (c) Banzhaf, M.; van den Berg van Saparoea, B.; Terrak, M.; Fraipont, C.; Egan, A.; Philippe, J.; Zapun, A.; Breukink, E.; Nguyen-Distèche, M.; den Blaauwen, T.; Vollmer, W. *Mol. Microbiol.* **2012**, *85*, 179–194. (d) Lebar, M. D.; Lupoli, T. J.; Tsukamoto, H.; May, J. M.; Walker, S.; Kahne, D. *J. Am. Chem. Soc.* **2013**, *135*, 4632–4635. (e) Zapun, A.; Philippe, J.; Abrahams, K. A.; Signor, L.; Roper, D. I.; Breukink, E.; Vernet, T. *ACS Chem. Biol.* **2013**, *8*, 2688–2696. (f) For computational modeling of cross-linking, see: Shi, Q.; Meroueh, S. O.; Fisher, J. F.; Mobashery, S. *J. Am. Chem. Soc.* **2011**, *133*, 5274–5283.
- (9) (a) While it has been difficult to characterize the transpeptidases, the substrate specificities of the low-molecular-mass (carboxypeptidase and endopeptidase) PBP classes have been studied. For a review, see: Pratt, R. F. *Cell. Mol. Life Sci.* **2008**, *65*, 2138–2155. (b) For a discussion of how amidated substrates are processed by a *B. subtilis* low-molecular-mass PBP, see: Nemmara, V. V.; Adediran, S. A.; Dave, K.; Duez, C.; Pratt, R. F. *Biochemistry* **2013**, *52*, 2627–2637.
- (10) (a) Primosigh, J.; Pelzer, H.; Maass, D.; Weidel, W. *Biochim. Biophys. Acta* **1961**, *46*, 68–80. (b) Glauner, B.; Höltje, J. V.; Schwarz, U. *J. Biol. Chem.* **1988**, *263*, 10088–10095.
- (11) (a) Warth, A. D.; Strominger, J. L. *Biochemistry* **1971**, *10*, 4349–4358. (b) Atrih, A.; Bacher, G.; Allmaier, G.; Williamson, M. P.; Foster, S. J. *J. Bacteriol.* **1999**, *181*, 3956–3966.
- (12) In *B. subtilis* spore PG, *m*-DAP is not amidated. See: Warth, A. D.; Strominger, J. L. *Proc. Natl. Acad. Sci. U.S.A.* **1969**, *64*, 528–535.
- (13) (a) Lupoli, T. J.; Tsukamoto, H.; Doud, E. H.; Wang, T.-S. A.; Walker, S.; Kahne, D. *J. Am. Chem. Soc.* **2011**, *133*, 10748–10751. (b) Tsukamoto, H.; Kahne, D. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 5050–5053. (c) Heseck, D.; Lee, M.; Zajiček, J.; Fisher, J. F.; Mobashery, S. *J. Am. Chem. Soc.* **2012**, *134*, 13881–13888.
- (14) Experimental procedures and characterization described in the SI.
- (15) (a) Jackson, G. E.; Strominger, J. L. *J. Biol. Chem.* **1984**, *259*, 1483–1490. (b) Popham, D. L.; Setlow, P. *J. Bacteriol.* **1995**, *177*, 326–335.
- (16) Lupoli, T. J.; Lebar, M. D.; Markovski, M.; Bernhardt, T.; Kahne, D.; Walker, S. *J. Am. Chem. Soc.* **2014**, *136*, 52–55.
- (17) PBPs also have been shown to transfer D-amino acids onto peptide substrates, see: (a) Pollock, J. J.; Ghuysen, J. M.; Linder, R.; Salton, M. R.; Perkins, H. R.; Nieto, M.; Leyh-Bouille, M.; Frère, J. M.; Johnson, K. *Proc. Natl. Acad. Sci. U.S.A.* **1972**, *69*, 662–666. (b) Adam, M.; Damblon, C.; Jamin, M.; Zorzi, W.; Dusart, V.; Galleni, M.; el Kharroubi, A.; Piras, G.; Spratt, B. G.; Keck, W. *Biochem. J.* **1991**, *279*, 601–604. (c) Kumar, I.; Pratt, R. F. *Biochemistry* **2005**, *44*, 9961–9970.
- (18) This result provides direct evidence that PBPs are responsible for D-amino acid incorporation into *B. subtilis* PG in vivo (see ref 20b).
- (19) (a) Kuru, E.; Hughes, H. V.; Brown, P. J.; Hall, E.; Tekkam, S.; Cava, F.; de Pedro, M. A.; Brun, Y. V.; Vannieuwenhze, M. S. *Angew. Chem., Int. Ed.* **2012**, *51*, 12519–12523. (b) Siegrist, M. S.; Whiteside, S.; Jewett, J. C.; Aditham, A.; Cava, F.; Bertozzi, C. R. *ACS Chem. Biol.* **2013**, *8*, 500–505. (c) Liechti, G. W.; Kuru, E.; Hall, E.; Kalinda, A.; Brun, Y. V.; VanNieuwenhze, M.; Maurelli, A. T. *Nature* **2014**, *506*, 507–510. (d) Shieh, P.; Siegrist, M. S.; Cullen, A. J.; Bertozzi, C. R. *Proc. Natl. Acad. Sci. U.S.A.* **2014**, *111*, 5456–5461.
- (20) (a) Lam, H.; Oh, D.-C.; Cava, F.; Takacs, C. N.; Clardy, J.; de Pedro, M. A.; Waldor, M. K. *Science* **2009**, *325*, 1552–1555. (b) Cava, F.; de Pedro, M. A.; Lam, H.; Davis, B. M.; Waldor, M. K. *EMBO J.* **2011**, *30*, 3442–3453. (c) Cava, F.; Lam, H.; de Pedro, M. A.; Waldor, M. K. *Cell. Mol. Life Sci.* **2011**, *68*, 817–831. (d) Leiman, S. A.; May, J. M.; Lebar, M. D.; Kahne, D.; Kolter, R.; Losick, R. *J. Bacteriol.* **2013**, *195*, 5391–5395.
- (21) (a) Tsuruoka, T.; Tamura, A.; Miyata, A.; Takei, T.; Iwamatsu, K.; Inouye, S.; Matsushashi, M. *J. Bacteriol.* **1984**, *160*, 889–894. (b) Caparros, M.; Pisabarro, A. G.; de Pedro, M. A. *J. Bacteriol.* **1992**, *174*, 5549–5559.
- (22) (a) Schleifer, K. H.; Kandler, O. *Bacteriol. Rev.* **1972**, *36*, 407–477. (b) Linnett, P. E.; Strominger, J. L. *J. Biol. Chem.* **1974**, *249*, 2489–2496. (c) Lavollay, M.; Arthur, M.; Fourgeaud, M.; Dubost, L.; Marie, A.; Veziriz, N.; Blanot, D.; Gutmann, L.; Mainardi, J.-L. *J. Bacteriol.* **2008**, *190*, 4360–4366.