

A Chromo-Fluorogenic Synthetic “Canary” for CO Detection Based on a Pyrenylvinyl Ruthenium(II) Complex

María E. Moragues,^{†,‡,§} Anita Toscani,^{||} Félix Sancenón,^{†,‡,§} Ramón Martínez-Mañez,^{*,†,‡,§} Andrew J. P. White,^{||} and James D. E. T. Wilton-Ely^{*,||}

[†]Centro de Reconocimiento Molecular y Desarrollo Tecnológico (IDM), Unidad Mixta Universidad Politécnica de Valencia-Universidad de Valencia, Camino de Vera s/n, 46022 Valencia, Spain

[‡]Departamento de Química, Universidad Politécnica de Valencia, Camino de Vera s/n, 46022 Valencia, Spain

[§]CIBER de Bioingeniería, Biomateriales y Nanomedicina (CIBER-BBN), Spain

^{||}Department of Chemistry, Imperial College London, South Kensington Campus, London SW7 2AZ, U.K.

S Supporting Information

ABSTRACT: The chromo-fluorogenic detection of carbon monoxide in air has been achieved using a simple, inexpensive system based on ruthenium(II). This probe shows exceptional sensitivity and selectivity in its sensing behavior in the solid state. A color response visible to the naked eye is observed at 5 ppb of CO, and a remarkably clear color change occurs from orange to yellow at the onset of toxic CO concentrations (100 ppm) in air. Even greater sensitivity (1 ppb) can be achieved through a substantial increase in turn-on emission fluorescence in the presence of carbon monoxide, both in air and in solution. No response is observed with other gases including water vapor. Immobilization of the probe on a cellulose strip allows the system to be applied in its current form in a simple optoelectronic device to give a numerical reading and/or alarm.

The development of electronic household detectors for harmful gases dates from the 1980s and 1990s. Back in 1911, canaries were traditionally used in coal mines as an early detection system against life-threatening gases such as carbon dioxide, carbon monoxide, and methane. The canary, normally a very songful bird, would stop singing and eventually die in the presence of these gases, signaling to the miners to exit the mine. Although considered as moderately toxic in comparison to other highly poisonous gases, carbon monoxide (CO) detection has always been of vital importance, as it lacks color, odor, or taste and is present in numerous natural and artificial environments.¹ Carbon monoxide is a temporary atmospheric pollutant in some urban areas, mainly arising from the exhaust of internal combustion engines (such as vehicles, portable generators, lawn mowers and power washers) but also from incomplete combustion of other fuels (including wood, coal, charcoal, oil, paraffin, propane, natural gas, and waste). Many technologies and devices (electrochemical,² optical,³ and those based on metal-oxide semiconductors⁴) have been developed to detect, monitor, and alert the leakage of carbon monoxide. These systems suffer from drawbacks such as false alarms arising due to water vapor or airborne particulates. As an alternative to these systems, the design of molecular probes for

the chromo-fluorogenic recognition of CO is of importance. In particular, colorimetric methods are undemanding, giving advantages such as real-time monitoring and the use of simple and inexpensive instrumentation. Moreover, certain colorimetric changes can be observed by the naked eye, thus making chromogenic protocols matchless for certain applications.


Optical detection of carbon monoxide dates back to the late twentieth century when the presence of CO was revealed by a chemically infused paper that turned brown when exposed to the gas. However, only in the past few years has the number of chromogenic probes for CO detection based on new sensing concepts increased. In this field, oxoacetato-bridged triruthenium cluster complexes,⁵ glass-immobilized rhodium complexes,⁶ iron porphyrin functionalized polypyrroles,⁷ and a phosphino diaminopyridine iron complex⁸ have been reported to display color modulations in the presence of carbon monoxide. More recently the optical detection of CO using bimetallic rhodium complexes has also been reported by some of us.⁹ Regardless of several advantages offered by chromogenic sensors, only a few probes for carbon monoxide detection using emission changes have been reported so far involving the use of iron¹⁰ and palladium¹¹ complexes. However, in some of these reported systems the modulations caused by the presence of carbon monoxide reveal particular shortcomings, typically involving poor color or emission changes, sensing in solution but not in air, and high detection limits, which hamper the use of these probes as viable sensing systems.

Within this context, we have taken advantage of the well-known ability of alkenyl ruthenium(II) complexes to react with small donor ligands such as carbon monoxide¹² to explore their use as potential colorimetric probes for CO sensing. The organometallic probe [Ru(CH=CHPyr-1)Cl(CO)(BTD)-(PPh₃)₂] (**1**) (Scheme 1) consists of a Ru(II) metal center bonded to a fluorogenic pyrenylvinyl (CH=CH-Pyr-1) ligand *trans* to a 2,1,3-benzothiadiazole (BTD) molecule that acts as an emission quencher (*vide infra*). The coordination sphere in **1** is completed by two triphenylphosphine ligands, a chloride, and a CO moiety. The probe was designed to express a dual chromo- and fluorogenic response to carbon monoxide. Thus,

Received: July 15, 2014

Published: August 4, 2014

Scheme 1. Formation of the Complexes $[\text{Ru}(\text{CH}=\text{CHPyr-1})\text{Cl}(\text{CO})(\text{BTD})(\text{PPh}_3)_2]$ (**1**) and $[\text{Ru}(\text{CH}=\text{CHPyr-1})\text{Cl}(\text{CO})_2(\text{PPh}_3)_2]$ (**2**)



the presence in the complex of a pyrene donor group and a BTD acceptor was expected to result in a ligand-to-ligand charge transfer band that would inhibit the fluorescence emission from pyrene. The interchange of the labile quencher BTD by CO to give **2** was envisaged to induce both a revival of the pyrene fluorescence and a color change due to the modification of the coordination sphere of the Ru(II) center.

Probe **1** was prepared in 96% yield following the simple and extensively utilized synthesis of alkenyl ruthenium complexes through hydorruthenation of a hydride complex by a suitable terminal alkyne.¹³ This process consists of the regio- and stereospecific insertion of a terminal alkyne into the Ru–H bond. Thus, the brightly colored σ -alkenyl 18-electron adduct **1** (see Scheme 1) was readily synthesized by reaction of the hydride $[\text{RuHCl}(\text{CO})(\text{PPh}_3)_3]$ ¹⁴ with 1-ethynylpyrene and BTD¹⁵ in dichloromethane at room temperature.

X-ray-quality crystals of **1** were obtained by recrystallization of the complex via vapor diffusion of diethyl ether onto a dichloromethane solution of **1**. Figure 1 provides a plot of the




Figure 1. Crystal structure of $[\text{Ru}(\text{CH}=\text{CHPyr-1})\text{Cl}(\text{CO})(\text{BTD})(\text{PPh}_3)_2]$ (**1**). Selected bond lengths (Å) and angles (deg): Ru–C30 1.824(3), Ru–C1 2.048(3), Ru–N21 2.238(3), Ru–P1 2.4064(7), Ru–P2 2.4099(6), Ru–Cl 2.4663(7); O30–C30–Ru 178.2(3), C30–Ru–P1 92.25(8), N21–Ru–P2 92.35(6), C30–Ru–Cl 178.60(8), C1–Ru–N21 171.45(11).

complex showing the atomic numbering scheme. In the structure, the Ru(II) center adopts a distorted octahedral coordination environment with two triphenylphosphine ligands in axial positions and four ligands (Cl, CO, BTD, and vinylpyrene) occupying the equatorial sites.

The electronic spectrum of a methanol solution of complex **1** in the visible region is dominated by a low-intensity band at ca. 500 nm attributed to a π – π^* pyrenylvinyl-to-BTD ligand-to-ligand charge transfer transition and a moderately intense absorption band at 390 nm due to a pyrene π – π^* transition.¹⁶

In a preliminary test, air samples containing CO were bubbled into dichloromethane or methanol solutions of **1** and an instantaneous and remarkable color change from red to yellow was observed. This color modulation is concomitant with significant reduction in intensity of the broad band at 390 nm, which additionally reveals the typical absorption bands of the pyrene group at 339 and 355 nm.¹⁷

All of these changes are consistent with displacement of the BTD ligand by CO and the formation of the complex $[\text{Ru}(\text{CH}=\text{CHPyr-1})\text{Cl}(\text{CO})_2(\text{PPh}_3)_2]$ (**2**) (Scheme 1). Crystals of **2** suitable for X-ray single-crystal diffraction studies were obtained by diethyl ether diffusion onto a dichloromethane solution of **1** under an atmosphere of CO. Figure 2 shows a plot of **2** which reveals the presence of a CO molecule *trans* to the pyrenylvinyl ligand due to the displacement of the BTD group, supporting the proposed mechanism.




Figure 2. Crystal structure of $[\text{Ru}(\text{CH}=\text{CHPyr-1})\text{Cl}(\text{CO})_2(\text{PPh}_3)_2]$ (**2**). Selected bond lengths (Å) and angles (deg): Ru–C19 1.857(2), Ru–C20 1.957(2), Ru–C1 2.104(2), Ru–P1 2.4126(5), Ru–P2 2.4060(5), Ru–Cl 2.4472(5); O19–C19–Ru 177.95(19), O20–C20–Ru 175.49(19), C19–Ru–P1 92.49(6), C19–Ru–Cl 175.94(6), C20–Ru–C1 178.37(8), C20–Ru–P2 93.09(6).

Moreover, the displacement of the BTD ligand by CO also results in the recovery of the fluorescence emission of the pyrene group. Thus, whereas **1** is very weakly fluorescent in methanol (λ_{exc} 355 nm, λ_{em} 458 nm), formation of **2** results in a remarkable 36-fold increase in emission. This can be seen in Figure 3, which shows the effect of bubbling increasing volumes of CO through a methanolic solution of compound **1**. Further experiments demonstrated that a similar chromo-fluorogenic response from **1** in the presence of CO was observed in methanol/water 1/1 v/v solutions (complex **1** is not soluble in higher proportions of water). Titration experiments with




Figure 3. Turn-on fluorescence response (λ_{ex} 355 nm) of a 1×10^{-4} mol dm^{-3} methanol solution of **1** upon addition of increasing volumes of CO (0, 0.03, 0.08, 0.2, 0.5, 0.7, 1 mL).

carbon monoxide in methanol/water 1/1 v/v allowed the determination of a limit of detection (LOD) as low as 1 ppb. The turn-on fluorescence is tentatively explained by the fact that BTD displacement eliminates the electron transfer between pyrenylvinyl and BTD ligands in **1**, resulting in a revival of the emission.

Motivated by the favorable chromo-fluorogenic sensing features of **1** observed in solution, the system was developed with a view to exploiting the potential use of this compound for CO detection in air. With this aim in mind, compound **1** was adsorbed on silica gel, resulting in an orange solid. In a typical test, this colored silica support containing the ruthenium probe was exposed to air containing different concentrations of carbon monoxide. An instantaneous and remarkable modulation of color from orange to yellow was observed, consistent with the formation of **2** (Figure 4). From further titration studies a detection limit as low as 0.6 ppb of CO in air was obtained. Furthermore, a visual color change to the naked eye from orange to yellow was observed for CO concentrations in air as low as 5 ppb. At concentrations of ca. 100 ppm, which is the concentration at which CO becomes toxic for short-term exposure to humans, the color change is extremely clear (see Figure 4).

Furthermore, the chromogenic response to CO in air using **1** adsorbed on silica gel was found to be highly selective over other gases tested. For instance, no color changes were found in the presence of Ar, N₂, O₂, CO₂, H₂S, and coordinating gases such as SO₂ and NO_x or in the presence of common volatile organic compounds such as chloroform, formaldehyde, acetone, ethanol, toluene, and hexane. Only acetonitrile induced some evident color change to yellow, but only at concentrations of ca. 5000 ppm. Importantly, in terms of potential applications (where steam may be present), exposure of **1** to water-saturated air did not trigger any chromogenic response.

These data show silica to be a simple and effective support for the chromogenic detection of CO in air using complex **1**. However, the fluorogenic response observed for **1** in the presence of carbon monoxide was relatively poor when using silica as the support. In contrast, **1** was found to display a clear turn-on emission enhancement at 495 nm in the presence of carbon monoxide when adsorbed on strips of cellulose paper. This support also offers many practical benefits, as it avoids the




Figure 4. Summary of the visual color and/or fluorescence changes observed by the naked eye in the absence (left) and in the presence of 100 ppm of CO (right) for compound **1** (a) dissolved in dichloromethane, (b) adsorbed on silica gel, (c) deposited on cellulose paper, and (d) deposited on cellulose paper and detected by fluorescence emission under UV irradiation at 355 nm.

need for silica and is easier to handle. It is also compatible with simple optoelectronic devices which can convert color changes into numerical readings.^{9c} Using cellulose paper the response of the probe was studied by monitoring the emission changes upon exposure to increasing concentrations of CO, and a remarkable LOD of 0.7 ppb was calculated from these experiments. A clear optical response to the naked eye can also be observed for concentrations of ca. 90 ppm (see Figure 4) using a conventional UV lamp, or simply the readily apparent color change. Complex **1** is also highly selective to CO on this support, and it was confirmed that no changes in color or in emission were observed in the presence of Ar, N₂, O₂, CO₂, SO₂, NO_x, H₂S, and common volatile organic compounds (e.g., chloroform, formaldehyde, acetone, ethanol, toluene, and hexane).

In summary, we report the design and use of a ruthenium(II) complex (**1**) for the simple, selective chromo-fluorogenic detection of carbon monoxide. To our knowledge, this is the first complex capable of both chromogenic and fluorogenic sensing of CO in air. In addition, the combination of two sensing modalities allows both simple visible detection as well as greater sensitivity when desired. The probe shows a color change visible to the naked eye at CO concentrations of 5 ppb and a remarkably clear change from orange to yellow at CO concentrations of 100 ppm in air. This serves as a clear warning at levels which become toxic on prolonged exposure to the gas. Moreover, **1** also displays a turn-on fluorescence emission in the presence of carbon monoxide. Both the color and turn-on emission modulations observed are highly selective and are due to a displacement of the BTD ligand in **1** by CO to yield complex **2**. This exceptionally high selectivity for CO over water vapor is of crucial importance in the potential application of this system in a domestic setting where steam is present (a

shortcoming of commercial devices). The high-yielding and straightforward synthetic procedure used to prepare **1** in air, coupled with the commercial availability and relatively low cost of ruthenium and other reagents, render compound **1** both accessible and inexpensive. These attributes make this system suitable for implementation in an easy-to-use portable optoelectronic device,^{9c} allowing **1** to be applied as an efficient chemosensor for the simple chromo-fluorogenic detection of this colorless and odorless killer.

■ ASSOCIATED CONTENT

■ Supporting Information

Text, figures, tables, and CIF files giving general considerations, instrumentation, synthesis of [Ru(CH=CHPyr-1)Cl(CO)-(BTD)(PPh₃)₂] (**1**) and [Ru(CH=CHPyr-1)Cl(CO)₂(PPh₃)₂] (**2**), silica gel immobilization of complex **1**, preparation of the cellulose probe of **1**, reactivity of **1** with CO, crystallographic data and structures of **1** and **2**, selected geometric data for compounds **1** and **2** at 173 K, fluorescence calibration curve for **1** in aqueous methanol solution and response upon addition of CO, sensitivity studies with **1** adsorbed on silica upon addition of CO, selectivity studies with **1** adsorbed on silica, and sensitivity studies with **1** deposited on cellulose strips upon addition of CO. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail for R.M.-M.: rmaez@qim.upv.es.

*E-mail for J.D.E.T.W.-E.: j.wilton-ely@imperial.ac.uk.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support from the Spanish Government (project MAT2012-38429-C04) and the Generalitat Valenciana (project PROMETEO/2009/016) is gratefully acknowledged. M.E.M. is grateful to the Spanish Ministerio de Ciencia e Innovación for an FPU grant. J.D.E.T.W.-E. and A.T. acknowledge the Leverhulme Trust for a grant (RPG-2012-634).

■ REFERENCES

- (1) (a) Wilkinson, L. J.; Waring, R. H.; Steventon, G. B.; Mitchell, S. C. *Molecules of Death. Carbon Monoxide: The Silent Killer*, 2nd ed.; Imperial College Press: London, 2007; p 37. (b) Rajiah, K.; Mathew, E. M. *Afr. J. Pharm. Pharmacol.* **2011**, *5*, 259. (c) Normand, J. C.; Durand, C.; Delafosse, B. *Arch. Mal. Prof. Environ.* **2011**, *72*, 240.
- (2) (a) <http://www.testo.com>. (b) <http://www.figarosensor.com>. (c) Gutmacher, D.; Foelml, C.; Vollenweider, W.; Hoefler, U.; Wöllenstein, J. *Procedia Eng.* **2011**, *25*, 1121. (d) <http://www.intlsensor.com>. (e) Kaminski, C.; Poll, A. *Electrochemical or Solid State H₂S Sensors: Which is Right For You?*; InTech: USA, 1985; p 55.
- (3) (a) Wolfrum, J. *Proc. Combust. Inst.* **1998**, *27*, 1. (b) Cho, K. H.; Lee, S. W.; Lee, J. H.; Choi, K. S. *Anal. Sci. Technol.* **2000**, *13*, 222. (c) <http://www.draeger.com>.
- (4) (a) Wang, B.; Zhao, Y.; Yu, D.; Hu, L. M.; Cao, J. S.; Gao, F. L.; Liu, Y.; Wang, L. J. *Chin. Sci. Bull.* **2010**, *55*, 228. (b) Ge, H.; Liu, J. *Sens. Actuators B* **2006**, *117*, 408.
- (5) Itou, M.; Araki, Y.; Ito, O.; Kido, H. *Inorg. Chem.* **2006**, *45*, 6114.
- (6) Gulino, A.; Gupta, T.; Altman, M.; Lo Schiavo, S.; Mineo, P. G.; Fragalà, I. L.; Evmenenko, G.; Dutta, P.; Van der Boom, M. E. *Chem. Commun.* **2008**, 2900.
- (7) Paul, S.; Amalraj, F.; Radhakrishnana, S. *Synth. Met.* **2009**, *159*, 1019.

(8) Benito-Garagorri, D.; Puchberger, M.; Mereiter, K.; Kirchner, K. *Angew. Chem., Int. Ed.* **2008**, *47*, 9142.

(9) (a) Esteban, J.; Ros-Lis, J. V.; Martínez-Mañez, R.; Marcos, M. D.; Moragues, M. E.; Soto, J.; Sancenón, F. *Angew. Chem., Int. Ed.* **2010**, *49*, 4934. (b) Moragues, M. E.; Esteban, J.; Ros-Lis, J. V.; Martínez-Mañez, R.; Marcos, M. D.; Martínez, M.; Soto, J.; Sancenón, F. *J. Am. Chem. Soc.* **2011**, *133*, 1576. (c) Moragues, M. E.; Montes-Robles, R.; Ros-Lis, J. V.; Alcañiz, M.; Ibañez, J.; Pardo, T.; Martínez-Mañez, R. *Sens. Actuators B* **2014**, *191*, 257. (d) Alba, M.; Pazos-Perez, N.; Vaz, B.; Formentin, P.; Tebbe, M.; Correa-Duarte, M. A.; Granero, P.; Ferré-Borrull, J.; Alvarez, R.; Pallares, J.; Fery, A.; de Lera, A. R.; Marsal, L. F.; Alvarez-Puebla, R. A. *Angew. Chem., Int. Ed.* **2013**, *52*, 6459. (e) Courbat, J.; Pascu, M.; Gutmacher, D.; Briand, D.; Wöllenstein, J.; Hoefler, U.; Severin, K.; de Rooij, N. F. *Procedia Eng.* **2011**, *25*, 1329.

(10) Wang, J.; Karpus, J.; Zhao, B. S.; Luo, Z.; Chen, P. R.; He, C. *Angew. Chem., Int. Ed.* **2012**, *51*, 9652.

(11) Michel, B. W.; Lippert, A. R.; Chang, C. J. *J. Am. Chem. Soc.* **2012**, *134*, 15668.

(12) (a) Loumrhari, H.; Ros, J.; Torres, M. R.; Santos, A.; Echavarren, A. M. *J. Organomet. Chem.* **1991**, *411*, 255. (b) Harris, M. C. J.; Hill, A. F. *Organometallics* **1991**, *10*, 3903.

(13) (a) Torres, M. R.; Vegas, A.; Santos, A.; Ros, J. *J. Organomet. Chem.* **1986**, *309*, 169. (b) Alcock, N. W.; Cartwright, J.; Hill, A. F.; Marcellin, M.; Rawles, H. M. *J. Chem. Soc., Chem. Commun.* **1995**, 369.

(14) (a) Vaska, L. *J. Am. Chem. Soc.* **1964**, *86*, 1943. (b) Schunn, R. A.; Wonchoba, E. R.; Wilkinson, G. *Inorg. Synth.* **1972**, *13*, 131.

(15) (a) Harris, M. C. J.; Hill, A. F. *J. Organomet. Chem.* **1992**, *438*, 209. (b) Herberhold, M.; Hill, A. F. *J. Organomet. Chem.* **1989**, *377*, 151. (c) Alcock, N. W.; Hill, A. F.; Roe, M. S. *J. Chem. Soc., Dalton Trans.* **1990**, 1737.

(16) Maurer, J.; Linseis, M.; Sarkar, B.; Schwederski, B.; Niemeyer, M.; Kaim, W.; Zális, S.; Anson, C.; Zabel, M.; Winter, R. F. *J. Am. Chem. Soc.* **2008**, *130*, 259.

(17) Marsh, N. D.; Mikolajczak, C. J.; Wornat, M. J. *Spectrochim. Acta, Part A* **2000**, *56*, 1499.