

Three-Component Coupling Based on Flash Chemistry. Carbolithiation of Benzyne with Functionalized Aryllithiums Followed by Reactions with Electrophiles

Aiichiro Nagaki, Daisuke Ichinari, and Jun-ichi Yoshida*

Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering, Kyoto University, Nishikyo-ku, Kyoto 615-8510, Japan

S Supporting Information

ABSTRACT: A flow microreactor method for three-component coupling of benzyne was developed based on flash chemistry. *o*-Bromophenyllithium generated from 1-bromo-2-iodobenzene and a functionalized aryllithium generated from the corresponding aryl halide were mixed at $-70\text{ }^{\circ}\text{C}$. In the subsequent reactor *o*-bromophenyllithium is decomposed to generate benzyne without affecting the functionalized aryllithium at $-30\text{ }^{\circ}\text{C}$, and carbolithiation of benzyne with the aryllithium took place spontaneously. The resulting functionalized biaryllithium was reacted with an electrophile in the subsequent reactor to give the corresponding three-component coupling product. The precise optimization of reaction conditions using the temperature–residence time mapping is responsible for the success of the present transformation. The present method has been successfully applied to the synthesis of boscalid.

Arynes including benzyne are highly reactive intermediates, and extensive studies on their synthetic applications have been reported so far.^{1,2} In particular, carbometalation of benzyne serves as a powerful method for constructing organic structures containing an *o*-disubstituted benzene ring.³ Although it is well-known that *o*-halophenyllithiums serve as good precursors of benzyne, it is difficult to use them for carbolithiation because various organolithium species are generated simultaneously, which compete with each other.

Schlosser reported a smart way of controlling the carbolithiation of benzyne by adding an *o*-dihalobenzene to an aryllithium. The reaction proceeds by a chain mechanism.^{4,5} Benzyne generated from *o*-halophenyllithium reacts with an aryllithium to give a biaryllithium, which undergoes halogen/lithium exchange with *o*-dihalobenzene to give a biaryl halide as a product regenerating *o*-halophenyllithium. However, the method suffers from the limitation of the substrates because of the requirement of delicate balance of the reactivity of organolithium intermediates (*vide infra*, Scheme 1).

If the biaryllithiums formed by carbolithiation could be used for reactions with various electrophiles at will, the carbolithiation method should serve as powerful and straightforward transition metal-free three-component synthesis of *o*-disubstituted benzenes. Such transformations, however, seem to be very difficult or impossible in batch because of short lifetimes of the intermediates.




Figure 1. Carbolithiation of benzyne with aryllithiums. Consecutive three-component coupling with subsequently added electrophiles.




Figure 2. An integrated flow microreactor system for the three-component coupling of benzyne, functionalized aryllithiums, and electrophiles. T-shaped micromixers: M1, M2, M3 and M4, microtube reactors: R1, R2, R3 and R4.

Flash chemistry⁶ solves the problem. We report here carbolithiation of benzyne generated from an *o*-halophenyllithium with aryllithiums followed by reactions of the resulting biaryllithiums with subsequently added electrophiles using an integrated flow microreactor system^{7–9} (Figure 1). It should be emphasized that the present flash method enables the use of short-lived aryllithiums bearing electrophilic functional groups such as cyano and nitro groups giving three-component coupling products bearing such functionalities.

We envisaged that space integration of reactions¹⁰ using the integrated flow microreactor system shown in Figure 2 enables the present transformation involving multiple short-lived reactive intermediates.¹¹ An *o*-halophenyllithium and an

Received: July 15, 2014

Published: August 14, 2014




Figure 3. A flow microreactor system for generation and accumulation of *o*-halophenyllithiums.




Figure 4. Temperature–residence time (T^1 – t^{R1}) maps for the halogen/lithium exchange reaction of (a) 1,2-dibromobenzene with *n*-BuLi, (b) 1-bromo-2-iodobenzene with PhLi, and (c) 1,2-diiodobenzene with PhLi followed by reaction with benzaldehyde in the flow microreactor.


aryllithium are generated by halogen/lithium exchange in M1/R1 and M2/R2, respectively. They are mixed in M3. In R3, the *o*-halophenyllithium is selectively decomposed to generate benzyne without affecting the aryllithium, which adds to benzyne spontaneously. The resulting biaryllithium is allowed to react with an electrophile in M4 and R4 to give the

Table 1. Three-Component Coupling of Benzyne, Aryllithiums, and Electrophiles Using the Integrated Flow Microreactor System^a

electrophile	product	yield (%)
		73
MeOTf ^{b,c}		53
TMSOTf ^b		63
PhCNO ^b		69
PhCHO ^b		58
Bu ₃ SnCl ^b		62
(CO ₂ Et) ₂ ^b		54
HCO ₂ Me ^{b,c}		51
I ₂ ^{b,c}		63
CBr ₄ ^b		64
C ₂ Cl ₆ ^b		71
NFSI ^{b,c}		63
NC-		70
MeOTf ^d		71
- ^{c,e}		50
O ₂ N-		50
MeI ^g		50
MeOTf ^d		58

^aThe reactions were usually carried out with 0.6 mmol/min of 1-bromo-2-iodobenzene. ^bR¹Li = BuLi, t^{R2} = 0.31 s, T^2 = 0 °C. ^c T^3 = -50 °C. ^dR¹Li = BuLi, t^{R2} = 0.31 s, T^2 = -30 °C. ^eR¹Li = BuLi, t^{R2} = 0.16 s, T^2 = -20 °C. ^fR¹Li = PhLi, t^{R2} = 1.18 s, T^2 = -70 °C. ^gR¹Li = BuLi, t^{R2} = 2.36 s, T^2 = -30 °C.

Scheme 1






Figure 5. Synthesis of boscalid.

corresponding three-component coupling product. To achieve these transformations in a harmonious way, the residence time and the temperature for each step should be precisely controlled.

First, we optimized the conditions for generating and accumulating *o*-halophenyllithiums from *o*-dihalobenzenes using a flow microreactor system¹² shown in Figure 3. The reactions of 1,2-dibromobenzene, 1-bromo-2-iodobenzene, and 1,2-diiodobenzene were carried with varying the temperature (T^1) and the residence time in R1 (t^{R1}), and *o*-halophenyllithiums were trapped with benzaldehyde. The yields of the products are plotted against T^1 and t^{R1} as contour maps with scattered overlays (Figure 4). In the case of 1,2-dibromobenzene using BuLi as a lithiating agent, the yields at short residence times are low, because bromine/lithium exchange is not so fast. The yield increases with an increase in the residence time and then decreases because of the decomposition to generate benzyne. In the case of 1-bromo-2-iodobenzene using PhLi as a lithiating agent, the yields at short residence times are high because iodine/lithium exchange is faster. At longer residence times the decomposition to benzyne takes place. In the case of 1,2-diiodobenzene using PhLi as a lithiating agent, the yields at short residence times are high, but the yield decreases very quickly with an increase in the residence time, presumably because *o*-iodophenyllithium is less stable than *o*-bromophenyllithium. Based on these results, we chose to generate *o*-bromophenyllithium as a precursor of benzyne from 1-bromo-2-iodobenzene. In this case the highest yield (93%) was obtained at -70 °C with the residence time of 0.16 s, and therefore these conditions were used for the subsequent studies. Also, it should be noted that the temperature–residence time maps in Figure 4 give crucial information about the conditions for benzyne generation.

Next, we examined the generations of benzyne from *o*-bromophenyllithium in the presence of *p*-chlorophenyllithium using an integrated flow microreactor system (Figure 2). The carbolithiation of benzyne with *p*-chlorophenyllithium followed by the reaction with added electrophiles should take place. Thus, 1-bromo-2-iodobenzene was reacted with PhLi ($T^1 = -70$ °C, $t^{R1} = 0.22$ s) at M1 and R1 to give *o*-bromophenyllithium, which is stable at this temperature within 1 s as indicated in Figure 4b. 1-Bromo-4-chlorobenzene was reacted with *n*-BuLi ($T^2 = 0$ °C, $t^{R2} = 0.31$ s)¹³ at M2 and R2 to generate *p*-chlorophenyllithium, and the resulting solution was cooled to -70 °C (T^1). The two organolithium species were mixed at M3 at -70 °C (T^1), and the mixture was warmed to -30 °C (T^3) in R3 ($t^{R3} = 2.69$ s) (See the Supporting

Information for details). Because *o*-bromophenyllithium decomposes completely at -50 °C within 2.4 s (Figure 4b), such decomposition should take place very quickly at -30 °C to generate benzyne. In contrast, *p*-chlorophenyllithium, which is generated at 0 °C, should be stable at -30 °C. Thus, flash generated benzyne reacted with *p*-chlorophenyllithium in R3 to give the corresponding biaryllithium, which was transferred to M4, where an electrophile was added. The reaction in R4 ($T^4 (= T^3) = -30$ °C, $t^{R4} = 1.06$ s) gave the final three-component coupling products.

Various electrophiles such as methyl trifluoromethanesulfonate, trimethylsilyl trifluoromethanesulfonate, phenyl isocyanate, benzaldehyde, tributyltin chloride, diethyl oxalate, and methyl formate were effective (Table 1). Notably, selective reaction of biaryllithiums with diethyl oxalate using a flow microreactor was achieved to obtain α -keto esters with high selectivity by virtue of fast 1:1 micromixing.¹⁴ Moderate yields for these reactions seem to be attributed to inevitable competitive reactions of benzyne with other organolithium species that should be present in the reaction mixture, such as *o*-bromophenyllithium and biaryllithiums.

Halogenating agents such as tetrabromomethane, iodine, hexachloroethane and *N*-fluorobenzenesulfonimide (NFSI) were also effective as electrophiles, and the corresponding biaryl halides were obtained.¹⁵ It should be noted that similar products were difficult to obtain selectively by the chain reaction via *in situ* halogen transfer. For example, the reaction of *p*-chlorobenzene and *o*-dibromobenzene in a batch macro reactor led to the formation of a mixture of desired 2-bromo-4'-chlorobiphenyl and undesired 2,2'-dibromobiphenyl at -25 °C (Scheme 1). At lower temperatures such -50 °C, the yield of the desired product slightly increased, but a significant amount of undesired 4-chlorobromobenzene was also obtained. In the present flash method, however, desired 2-bromo-4'-chlorobiphenyl was obtained selectively in a good yield (Table 1).

In addition to *p*-chlorophenyllithium, various aryllithiums bearing electrophilic functional groups¹⁶ can be used for the present three-component coupling. For example, *o*-, *m*-, and *p*-cyanophenyllithiums generated from -20 to -30 °C using BuLi ($t^{R2} = 0.16$ – 0.31 s)¹⁷ and *p*-nitrophenyllithium generated at -30 °C using PhLi ($t^{R2} = 0.31$ s)¹⁸ were successfully used in this transformation. Benzyne were generated without affecting such functionalized aryllithiums. The carbolithiation followed by the subsequent reactions with electrophiles gave the three-component coupling products. Such transformations are very difficult or impossible to achieve using conventional batch reactors. Heteroaryllithiums such as 2-pyridyllithium ($T^2 = -30$ °C, $t^{R2} = 2.36$ s) and 2-thiophenyllithium ($T^2 = -30$ °C, $t^{R2} = 0.31$ s)¹⁹ can also be used for the three-component coupling, because they are also stable at -30 °C within a second or so.

To demonstrate the utility of the present flash method, we examined the application to the synthesis of boscalid,^{20,21} an important fungicide, which belongs to the class of succinate dehydrogenase inhibitors. Benzyne generation from *o*-bromophenyllithium and the carbolithiation with 4-chlorophenyllithium followed by the reaction with tosyl azide gave 4'-chloro-2-azidobiphenyl in 60% yield (Figure 5). Reduction with NaBH₄ followed by the reaction with 2-chloronicotinoyl chloride gave boscalid in 88% yield (53% overall yield from 1-bromo-2-iodobenzene).

In conclusion, flash chemistry using integrated flow microreactor systems enabled three-component coupling of benzyne generated from *o*-bromophenyllithium, functionalized aryl-

lithiums, and electrophiles. The precise control of the reaction conditions based on the relative stability of the organolithium species that are involved in the benzyne formation and the carbolithiation is responsible for the success of the transformation. Further work is in progress to explore the full scope of this useful three-component coupling and its applications in organic synthesis.

■ ASSOCIATED CONTENT

● Supporting Information

Experimental procedures, spectroscopic data of compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

yoshida@sbchem.kyoto-u.ac.jp

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was partially supported by the Grant-in-Aid for Scientific Research (S) (no. 26220804) and Scientific Research (B) (no. 26288049).

■ REFERENCES

- (1) Reviews (a) Wenk, H. H.; Winkler, M.; Sander, W. *Angew. Chem., Int. Ed.* **2003**, *42*, 502. (b) Pena, D.; Perez, D.; Guitian, E. *Angew. Chem., Int. Ed.* **2006**, *45*, 3579. (c) Yoshida, H.; Ohshita, J.; Kunai, A. *Bull. Chem. Soc. Jpn.* **2010**, *83*, 199. (d) Kitamura, K. *Aust. J. Chem.* **2010**, *63*, 987. (e) Tadross, P. M.; Stoltz, B. M. *Chem. Rev.* **2012**, *112*, 3550.
- (2) Some recent examples (a) Goetz, A. E.; Garg, N. K. *Nat. Chem.* **2013**, *5*, 54. (b) Hendrick, C. E.; McDonald, S. L.; Wang, Q. *Org. Lett.* **2013**, *15*, 3444. (c) He, Z.; Jamison, T. F. *Angew. Chem., Int. Ed.* **2014**, *53*, 3353.
- (3) (a) Gilman, H.; Gaj, B. J. *J. Org. Chem.* **1957**, *22*, 447. (b) Dougherty, T. K.; Lau, K. S. Y.; Hedberg, F. L. *J. Org. Chem.* **1983**, *48*, 5273.
- (4) Leroux, F.; Schlosser, M. *Angew. Chem., Int. Ed.* **2002**, *41*, 4272.
- (5) A similar chain reaction using organolithiums was developed (a) Leroux, F. R.; Bonnafoux, L.; Heiss, C.; Colobert, F.; Lanfranchi, D. A. *Adv. Synth. Catal.* **2007**, *349*, 2705. (b) Bonnafoux, L.; Colobert, F.; Leroux, F. R. *Synlett* **2010**, 2953. (c) Diemer, V.; Leroux, F. R.; Colobert, F. *Eur. J. Org. Chem.* **2011**, 327. (d) Diemer, V.; Begaud, M.; Leroux, F. R.; Colobert, F. *Eur. J. Org. Chem.* **2011**, 341. (e) Hamura, T.; Chuda, Y.; Nakatsuji, Y.; Suzuki, K. *Angew. Chem., Int. Ed.* **2012**, *51*, 3368.
- (6) Flash chemistry is defined as a field of chemical synthesis where extremely fast reactions are conducted in a highly controlled manner to produce the desired compounds with high selectivity (a) Yoshida, J. *Chem. Commun.* **2005**, 4509. (b) Yoshida, J.; Nagaki, A.; Yamada, T. *Chem.—Eur. J.* **2008**, *14*, 7450. (c) Nieuwland, P. J.; Koch, K.; van Harskamp, N.; Wehrens, R.; van Hest, J. C. M.; Rutjes, F. P. J. T. *Chem. Asian J.* **2010**, *5*, 799. (d) Yoshida, J. *Chem. Rec.* **2010**, *10*, 332. (e) Yoshida, J.; Takahashi, Y.; Nagaki, A. *Chem. Commun.* **2013**, 49, 9896.
- (7) Books on flow microreactor synthesis (a) Ehrfeld, W.; Hessel, V.; Löwe, H. *Microreactors*; Wiley-VCH: Weinheim, 2000. (b) Hessel, V.; Hardt, S.; Löwe, H. *Chemical Micro Process Engineering*; Wiley-VCH Verlag: Weinheim, 2004. (c) Yoshida, J. *Flash Chemistry. Fast Organic Synthesis in Microsystems*; Wiley-Blackwell, 2008. (d) *Micro Process Engineering*; Hessel, V., Renken, A., Schouten, J. C., Yoshida, J., Eds.; Wiley-VCH Verlag: Weinheim, 2009. (e) *Microreactors in Organic Chemistry and Catalysis*, 2nd ed.; Wirth, T., Ed.; Wiley-VCH Verlag: Weinheim, 2013.
- (8) Recent reviews on flow microreactor synthesis (a) Fukuyama, T.; Rahman, M. T.; Sato, M.; Ryu, I. *Synlett* **2008**, 151. (b) Hartman, R. L.; Jensen, K. F. *Lab Chip* **2009**, *9*, 2495. (c) McMullen, J. P.; Jensen, K. F. *Annu. Rev. Anal. Chem.* **2010**, *3*, 19. (d) Yoshida, J.; Kim, H.; Nagaki, A. *ChemSusChem* **2011**, *4*, 331. (e) Wiles, C.; Watts, P. *Green Chem.* **2012**, *14*, 38. (f) Kirschning, A.; Kupracz, L.; Hartwig, J. *Chem. Lett.* **2012**, *41*, 562. (g) McQuade, D. T.; Seeberger, P. H. *J. Org. Chem.* **2013**, *78*, 6384. (h) Elvira, K. S.; i Solvas, X. C.; Wootton, R. C. R.; deMello, A. J. *Nat. Chem.* **2013**, *5*, 905. (i) Pastre, J. C.; Browne, D. L.; Ley, S. V. *Chem. Soc. Rev.* **2013**, *42*, 8849. (j) Baxendale, I. R. *J. Chem. Technol. Biotechnol.* **2013**, *88*, 519.
- (9) Some selected recent examples (a) Cantillo, D.; Baghbanzadeh, M.; Kappe, C. O. *Angew. Chem., Int. Ed.* **2012**, *51*, 10190. (b) Shu, W.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2012**, *51*, 5355. (c) Nagaki, A.; Moriwaki, Y.; Yoshida, J. *Chem. Commun.* **2012**, 48, 11211. (d) Lévesque, F.; Seeberger, P. H. *Angew. Chem., Int. Ed.* **2012**, *51*, 1706. (e) Basavaraju, K. C.; Sharma, S.; Maurya, R. A.; Kim, D. P. *Angew. Chem., Int. Ed.* **2013**, *52*, 6735. (f) Brancour, C.; Fukuyama, T.; Mukai, Y.; Skrydstrup, T.; Ryu, I. *Org. Lett.* **2013**, *15*, 2794. (g) Nguyen, J. D.; Reiß, B.; Dai, C.; Stephenson, C. R. J. *Chem. Commun.* **2013**, 49, 4352. (h) Battilocchio, C.; Hawkins, J. M.; Ley, S. V. *Org. Lett.* **2013**, *15*, 2278. (i) Kleinke, A. S.; Jamison, T. F. *Org. Lett.* **2013**, *15*, 710. (j) Guetzoyan, L.; Nikbin, N.; Baxendale, I. R.; Ley, S. V. *Chem. Sci.* **2013**, *4*, 764. (k) Fuse, S.; Mifune, Y.; Takahashi, T. *Angew. Chem., Int. Ed.* **2014**, *53*, 851. (l) Nagaki, A.; Takahashi, Y.; Yoshida, J. *Chem.—Eur. J.* **2014**, in press.
- (10) (a) Suga, S.; Yamada, D.; Yoshida, J. *Chem. Lett.* **2010**, 39, 404. (b) Yoshida, J.; Saito, K.; Nokami, T.; Nagaki, A. *Synlett* **2011**, 1189.
- (11) Some examples of generation and reactions of short-lived organolithiums in flow (a) Nagaki, A.; Takizawa, E.; Yoshida, J. *J. Am. Chem. Soc.* **2009**, *131*, 1654. (b) Tomida, Y.; Nagaki, A.; Yoshida, J. *J. Am. Chem. Soc.* **2011**, *133*, 3744. (c) Kim, H.; Nagaki, A.; Yoshida, J. *Nat. Commun.* **2011**, *2*, 264. (d) Nagaki, A.; Matsuo, C.; Kim, S.; Saito, K.; Miyazaki, A.; Yoshida, J. *Angew. Chem., Int. Ed.* **2012**, *51*, 3245.
- (12) Usutani, H.; Tomida, Y.; Nagaki, A.; Okamoto, H.; Nokami, T.; Yoshida, J. *J. Am. Chem. Soc.* **2007**, *129*, 3046.
- (13) The conditions were optimized separately. See Nagaki, A.; Tomida, Y.; Usutani, H.; Kim, H.; Takabayashi, N.; Nokami, T.; Okamoto, H.; Yoshida, J. *Chem.—Asian J.* **2007**, *2*, 1513.
- (14) Nagaki, A.; Ichinari, D.; Yoshida, J. *Chem. Commun.* **2013**, 49, 3242.
- (15) Nagaki, A.; Uesugi, Y.; Kim, H.; Yoshida, J. *Chem.—Asian J.* **2013**, *8*, 705.
- (16) (a) Knochel, P. *Handbook of Functionalized Organometallics*; Wiley-VCH: Weinheim, 2005. (b) Boudier, A.; Bromm, L. O.; Lotz, M.; Knochel, P. *Angew. Chem., Int. Ed.* **2000**, *39*, 4414.
- (17) Nagaki, A.; Kim, H.; Matsuo, C.; Usutani, H.; Yoshida, J. *Org. Biomol. Chem.* **2010**, *8*, 1212.
- (18) (a) Nagaki, A.; Kim, H.; Yoshida, J. *Angew. Chem., Int. Ed.* **2009**, *48*, 8063. (b) Nagaki, A.; Kim, H.; Yoshida, J. *Angew. Chem., Int. Ed.* **2008**, *47*, 7833. (c) Nagaki, A.; Kim, H.; Moriwaki, Y.; Matsuo, C.; Yoshida, J. *Chem.—Eur. J.* **2010**, *16*, 11167.
- (19) (a) Nagaki, A.; Yamada, S.; Doi, M.; Tomida, Y.; Takabayashi, N.; Yoshida, J. *Green Chem.* **2011**, *13*, 1110. (b) Nagaki, A.; Yamada, D.; Yamada, S.; Doi, M.; Tomida, Y.; Takabayashi, N.; Yoshida, J. *Aust. J. Chem.* **2013**, *66*, 199.
- (20) (a) Kuck, K.-H.; Gisi, U. In *Modern Crop Protection Compounds*; Krämer, W., Schirmer, U., Eds.; Wiley-VCH: Weinheim, 2007; Vol. 2, pp 415–432. (b) Earley, F. In *Modern Crop Protection Compounds*; Krämer, W., Schirmer, U., Eds.; Wiley-VCH: Weinheim, 2007; Vol. 2, pp 433–538.
- (21) (a) Eicken, K.; Goetz, N.; Harreus, A.; Ammermann, E.; Lorenz, G.; Rang, H. European Patent EP0545099, November 24, 1993. (b) Engel, S.; Oberding, T. WO Patent 2006/092429, September 8, 2006. (c) Glasnov, T. N.; Kappe, C. O. *Adv. Synth. Catal.* **2010**, *352*, 3089.