

Correction to “Reversible Interconversion of CO₂ and Formate by a Molybdenum-Containing Formate Dehydrogenase”

Arnau Bassegoda, Christopher Madden, David W. Wakerley, Erwin Reisner,* and Judy Hirst*

J. Am. Chem. Soc. **2014**, *136*, 15473–15476. DOI: 10.1021/ja508647u

Page 15474. Due to a production error, the *y*-axis legend to Figure 2 was published with incorrect units. The units for *I* should be mA cm⁻², not μA cm⁻². The corrected Figure 2 is given below.




Figure 2. Cyclic voltammograms showing reversible CO₂ reduction and formate oxidation by *Ec*FDH-H adsorbed on a graphite-epoxy electrode, pH 6.0 (left), 6.8 (middle), and 8.0 (right). The points of intersection (marked with crosses) define the reduction potentials for the CO₂/formate interconversion (vertical lines). First voltammetric cycles are shown in black and subsequent cycles (2–4, 10, and 20) in gray. Voltammograms recorded in the absence of substrates are shown as dashed traces. Conditions: 10 mM CO₂,²⁵ 10 mM formate, 25 mM of each of four pH buffers (acetate, MES, HEPES, and TAPS), voltammetric scan rate 25 mV s⁻¹, electrode rotation rate 2000 rpm, 23 °C.