


Stereochemistry and Mechanism of Undecylprodigiosin Oxidative Carbocyclization to Streptorubin B by the Rieske Oxygenase RedG

David M. Withall, Stuart W. Haynes, and Gregory L. Challis*

Department of Chemistry, University of Warwick, Coventry CV4 7AL, United Kingdom

S Supporting Information

ABSTRACT: The prodiginines are a group of specialized metabolites that share a 4-methoxyppyrolyldipyrromethene core structure. Streptorubin B is a structurally remarkable member of the prodiginine group produced by *Streptomyces coelicolor* A3(2) and other actinobacteria. It is biosynthesized from undecylprodigiosin by an oxidative carbocyclization catalyzed by the Rieske oxygenase-like enzyme RedG. Undecylprodigiosin derives from the RedH-catalyzed condensation of 2-undecylpyrrole and 4-methoxy-2, 2'-bipyrrole-5-carboxaldehyde (MBC). To probe the mechanism of the RedG-catalyzed reaction, we synthesized 2-(5-pentoxypentyl)-pyrrole, an analogue of 2-undecylpyrrole with an oxygen atom next to the site of C–C bond formation, and fed it, along with synthetic MBC, to *Streptomyces albus* expressing *redH* and *redG*. This resulted in the production of the 6'-oxa analogue of undecylprodigiosin. In addition, a small amount of a derivative of this analogue lacking the *n*-pentyl group was produced, consistent with a RedG catalytic mechanism involving hydrogen abstraction from the alkyl chain of undecylprodigiosin prior to pyrrole functionalization. To investigate the stereochemistry of the RedG-catalyzed oxidative carbocyclization, [7'-²H](7'*R*)-2-undecylpyrrole and [7'-²H](7'*S*)-2-undecylpyrrole were synthesized and fed separately, along with MBC, to *S. albus* expressing *redH* and *redG*. Analysis of the extent of deuterium incorporation into the streptorubin B produced in these experiments showed that the *pro-R* hydrogen atom is abstracted from C-7' of undecylprodigiosin and that the reaction proceeds with inversion of configuration at C-7'. This contrasts sharply with oxidative heterocyclization reactions catalyzed by other nonheme iron-dependent oxygenase-like enzymes, such as isopenicillin N synthase and clavaminic synthase, which proceed with retention of configuration at the carbon center undergoing functionalization.



INTRODUCTION

The prodiginines are a group of specialized metabolites with a common 4-methoxyppyrolyldipyrromethene core produced by a variety of bacteria, including *Streptomyces* and related actinobacteria (Figure 1).¹ Members of the prodiginine family possess a range of interesting biological activities and have recently been shown to be orally effective antimalarials in a murine infection model.^{1,2}

Streptomyces coelicolor A3(2) produces undecylprodigiosin **1** and its carbocyclic derivative streptorubin B **2** (Figure 1).^{3,4} Transannular interactions in the 10-membered carbocycle of streptorubin B **2** restrict rotation around the C2–C1' and C4–C7' bonds, resulting in the formation of atropisomers.⁵ Nuclear Overhauser effect spectroscopy (NOESY) NMR experiments showed that the hydrochloride salt of **2** exists as predominantly the *anti* atropisomer in solution.^{5,6} The absolute configuration of this atropisomer was assigned as 7'*S* by ¹H and ²H NMR spectroscopic analyses of stereoselectively deuterium labeled streptorubin B.HCl produced via a mutasynthesis approach.⁵ This stereochemical assignment was confirmed by enantioselective total synthesis and X-ray crystallographic analysis.⁶ Chiral high pressure liquid chromatography (HPLC) comparisons of **2** isolated from *S. coelicolor* with chemically synthesized (7'*S*)-, (7'*R*)-, and (7'*RS*)-streptorubin B showed that the

natural material consists of the *anti*-(7'*S*), *syn*-(7'*S*), and *anti*-(7'*R*) isomers in an approximately 88:7:5 ratio.^{5,6}

The biosynthesis of undecylprodigiosin **1** and streptorubin B **2** in *S. coelicolor* has been extensively studied.⁷ Hybrid nonribosomal peptide synthetase/polyketide synthase/ α -oxoamine synthase assembly lines construct 4-methoxy-2,2'-bipyrrole-5-carboxaldehyde (MBC) **3** and 2-undecylpyrrole **4** from L-proline, malonyl-CoA, L-serine, and S-adenosyl-L-methionine, and one unit of acetyl-CoA, six units of malonyl-CoA, and a unit of glycine, respectively (Figure 1) (CoA = coenzyme A, Ad = adenosine).^{4,8} MBC **3** and 2-undecylpyrrole **4** are condensed to form undecylprodigiosin **1** by RedH (Figure 1).⁹ The broad substrate tolerance of RedH has been exploited to produce undecylprodigiosin analogues via a mutasynthesis approach.⁹ Streptorubin B **2** is formed from undecylprodigiosin **1** via an unusual oxidative carbocyclisation reaction catalyzed by RedG,^{10,11} a Rieske nonheme iron-dependent oxygenase-like enzyme (Figure 1).¹² Remarkably, in *Streptomyces longisporus* DSM 40667 McpG, which is 75% similar in sequence to RedG, catalyzes the regio- and stereodivergent carbocyclisation of undecylprodigiosin **1** to form metacycloprodigiosin **5** (Figure 1).¹⁰

Received: April 17, 2015

Published: May 29, 2015




Figure 1. Pathway for the biosynthesis of undecylprodigiosin **1** and streptorubin B **2** in *S. coelicolor* A3(2). RedH catalyzes the condensation of 2-undecylpyrrole **4** and 4-methoxy-2,2'-bipyrrole-5-carboxaldehyde **3** to form **1**, which undergoes RedG-catalyzed oxidative carbocyclization to form **2**. In *Streptomyces longisporusruiber* DSM40667, McpG catalyzes the oxidative carbocyclisation of **1** to form metacycloprodigiosin **5**.




Figure 2. Proposed mechanism for the RedG-catalyzed conversion of undecylprodigiosin **1** to streptorubin **2**. Alternative Fe(V)O(OH) and C-7'/C-5 cation intermediates are also possible.

Based on the well-studied catalytic mechanism of naphthalene dioxygenase, we recently proposed a mechanism for the RedG-catalyzed oxidative carbocyclization reaction involving abstraction of a hydrogen atom from C-7' of enzyme bound undecylprodigiosin **1** by the active site Fe(III)OOH complex **6** to form a carbon-centered radical **7** and an Fe(IV)O(OH₂) complex **8** (Figure 2).¹² Addition of the radical to C-4 of the proximal pyrrole leads to formation of the 10-membered carbocycle **9** bearing a radical at C-5, which can be stabilized by delocalization into the adjacent π -system. Hydrogen abstraction from C-4 of **9** by the Fe(IV)O(OH₂) complex **8** yields

streptorubin B **2**. It was noted that an Fe(V)O(OH) complex, a C-7' carbocation and a 10-membered carbocycle bearing a cation at C-5 are plausible alternative intermediates to **6**, **7** and **9**, respectively.¹² One of the more remarkable facets of this mechanistic proposal is that the hydrocarbon chain of undecylprodigiosin **1** is oxidized by the Fe(III)OOH complex in preference to the considerably more electron rich 4-methoxypyrrolyldipyrromethene π -system.

Here we report the synthesis of 2-(5-pentoxypentyl)-pyrrole **10**, a 2-undecylpyrrole analogue designed to probe the RedG catalytic mechanism. The products resulting from feeding of

Scheme 1. Synthesis of 2-(5-Pentoxypentyl)-pyrrole 10





Figure 3. Extracted ion chromatograms (EICs) at $m/z = 396.26; 326.18$ from LC-MS analyses of culture extracts of *S. albus* expressing *redHG* fed with MBC 3 and 2-(5-pentoxypentyl)-pyrrole 10 (top panel), and *S. albus* expressing *redH* fed with MBC 3 and 2-(5-pentoxypentyl)-pyrrole 10 (middle panel). The corresponding EIC from LC-MS analysis of a synthetic standard of 5-hydroxypentylprodigiosin 17 is shown in the bottom panel. Peak doubling is due to geometrical isomerism, resulting from rotation around the bond between C-1" and C-5 of ring C.

this 2-undecylpyrrole analogue, together with synthetic MBC 3, to *Streptomyces albus* J1704 expressing *redH* and *redG* demonstrate that hydrocarbon oxidation precedes functionalization of the 4-methoxypyrrrolyldipyrromethene moiety. We also report the synthesis of both enantiomers of $[7'-^2\text{H}]$ -2-undecylpyrrole. Feeding of these stereoselectively deuterium labeled 2-undecylpyrroles, along with synthetic MBC 3, to *S. albus* expressing *redH* and *redG* elucidated the stereochemical course of the undecylprodigiosin to streptorubin B conversion, allowing the remarkable chemoselectivity of RedG to be rationalized.

EXPERIMENTAL PROCEDURES

See the Supporting Information.

RESULTS AND DISCUSSION



Synthesis of 2-(5-Pentoxypentyl)-pyrrole 10. We reasoned that an analogue of undecylprodigiosin 1 with an oxygen atom in place of the C-6' methylene group would allow us to probe the timing of hydrocarbon oxidation relative to pyrrole functionalization in the RedG-catalyzed carbocyclization reaction, by stabilizing a radical/cationic intermediate at C-7', which may lead to the formation of shunt products. Thus, we synthesized 2-(5-pentoxypentyl)-pyrrole 10 from *t*-

butyldimethylsilyl (TBS)-protected 1,5-pentandiol 11¹³ in six steps (Scheme 1). Deprotonation of 11 with sodium hydride and reaction of the resulting alkoxide with 1-bromopentane afforded the corresponding ether 12. Deprotection of 12 with tetra-*n*-butyl ammonium fluoride (TBAF) gave alcohol 13, which was oxidized to carboxylic acid 14, via the aldehyde, by sequential treatment with iodoxybenzoic acid (IBX) and oxone.^{14,15} Coupling of 14 with pyrrol magnesium bromide, followed by reduction of the resulting 2-acylpyrrole 15, were both accomplished according to established procedures,⁴ affording 2-(5-pentoxypentyl)-pyrrole 10.

Feeding of 2-(5-Pentoxypentyl)-pyrrole 10 and MBC 3 to *S. albus* Expressing *redHG*. We previously reported that feeding of MBC 3 and 2-undecylpyrrole 4 to a strain of *S. albus* expressing *redH* under the control of the constitutive *ermE** promoter produces undecylprodigiosin 1.¹¹ We also reported that a similar experiment using a strain of *S. albus* expressing *redHG* results in the production of streptorubin B 2, in addition to undecylprodigiosin 1.¹¹

Feeding of 2-(5-pentoxypentyl)-pyrrole 10 and synthetic MBC 3⁸ to *S. albus* expressing *redH* resulted in the formation of a red pigment that gave rise to ions with $m/z = 396.26$ in liquid chromatography-mass spectrometry (LC-MS) analyses, corresponding to the $[\text{M} + \text{H}]^+$ ion for the 6'-oxa undecylprodi-

Scheme 2. Synthesis of 5-Hydroxypentylprodigiosin 17

Scheme 3. Synthesis of $[7'-^2\text{H}](7'R)$ -2-Undecylpyrrole 22^a

^aCompound 26 is the TBS ether of (*R*)-glycidol, which was prepared according to a literature procedure.¹⁸ $[7'-^2\text{H}](7'S)$ -2-undecylpyrrole 23 was synthesized via the same route using the TBS ether of (*S*)-glycidol.

giosin analogue 16 (Figure 3). No production of 16 was observed when 10 and 3 were fed to wild type *S. albus*. When 2-(5-pentoxypentyl)-pyrrole 10 and synthetic MBC 3 were fed to *S. albus* expressing *redHG*, the 6'-oxa undecylprodigiosin analogue 16 was also produced (Figure 3), but none of the corresponding streptorubin B analogue could be detected. Instead ultrahigh resolution LC-MS analyses showed that a small amount of a compound with a molecular formula corresponding to 5-hydroxypentylprodigiosin 17 (calculated for $\text{C}_{19}\text{H}_{24}\text{N}_3\text{O}_2^+$, 326.1863; found, 326.1858), a derivative of 16 lacking the *n*-pentyl group, was produced (Figure 3). Compound 17 was not produced when 10 and 3 were fed to *S. albus* expressing *redH* alone (Figure 3).

To confirm the production of 5-hydroxypentylprodigiosin 17 when 2-(5-pentoxypentyl)-pyrrole 10 and MBC 3 are fed to *S. albus* expressing *redHG*, an authentic synthetic standard was prepared in six steps from TBS-protected 1,5-pentandiol 11¹³ (Scheme 2). Swern oxidation of 11 yielded the corresponding aldehyde,¹⁶ which was further oxidized to the acid 18 using Pinnick's reagents.¹⁷ Coupling of acid 18 with pyrrole magnesium bromide and reduction of the resulting 2-acylpyrrole 19 to the corresponding alkylpyrrole 20,⁴ was followed by deprotection with TBAF to give 2-(5-hydroxypentyl)-pyrrole 21. Acid-promoted condensation of 21 with BOC-protected MBC⁶ afforded 5-hydroxypentylprodigiosin 17 as its HCl salt. LC-MS/MS analyses confirmed that the dealkylated undecylprodigiosin analogue produced when 2-(5-

pentoxypentyl)-pyrrole 10 and MBC 3 are fed to *S. albus* expressing *redHG* has the same retention time and fragmentation pattern as the authentic standard of 17 (Figure 3 and Supporting Information).

Synthesis of 2-Undecylpyrrole Stereoselectively Deuterium Labeled at C-7'. To probe the stereoselectivity of hydrogen abstraction from C-7' and the stereochemical course of the subsequent carbocyclization reaction, we synthesized $[7'-^2\text{H}](7'R)$ -2-undecylpyrrole 22 and $[7'-^2\text{H}](7'S)$ -2-undecylpyrrole 23 from commercially available *R*- and *S*-glycidol, respectively, (Scheme 3). 4-Pentyn-1-ol 24 was protected as its 4-methoxybenzyl ether 25. TBS-protected *R*-glycidol 26¹⁸ was reacted in the presence of $\text{BF}_3\cdot\text{OEt}_2$ with the acetylide derived from 25 by deprotonation with *n*-BuLi to yield homopropargylic alcohol 27. Reduction with H_2 over 10% Pd-C converted 27 to the saturated diol 28. Selective protection of the primary alcohol in 28 with pivaloyl choride in pyridine gave ester 29, which was converted to the corresponding tosylate 30 by treatment with 4-toluenesulfonyl chloride (TsCl), 4-dimethylaminopyridine (DMAP) and Et_3N . Reaction of 30 with TBAF yielded epoxide 31, which was converted to the alcohol 32 using *n*-PrLi,¹⁹ CuCN, and $\text{BF}_3\cdot\text{OEt}_2$.²⁰ The enantiopurity of 32 was assessed by converting a small portion to the corresponding Mosher's esters,²¹ which were analyzed by ^1H NMR spectroscopy (see the Supporting Information). While a precise enantiometric excess for the esters could not be determined due to signal overlap, the ^1H NMR data confirmed




Figure 4. Mass spectra of streptorubin B 2 from LC-MS analyses of mycelial extracts of *S. albus* expressing *redHG* fed with MBC 3 and [7'-²H](7'*S*)-2-undecylpyrrole 23 (top spectrum) or [7'-²H](7'*R*)-2-undecylpyrrole 22 (bottom spectrum). When 23 is fed, 92% of the label is incorporated into streptorubin B, whereas when 22 is fed, 77% of the label is calculated to be lost (after subtraction from the $m/z = 393.2750$ signal in the bottom panel of the component due to unlabeled ¹³C₁ streptorubin B isotopomer).

that 32 is predominantly the 7*S* enantiomer. Alcohol 32 was converted to the tosylate 33, which was reacted with LiAlD₄ to afford (7*R*)-7-deuterioundecan-1-ol 34.²² Swern oxidation of 34 gave the corresponding aldehyde,¹⁶ which was further oxidized to the acid 35 using oxone.¹⁵ Coupling of acid 35 with pyrrol magnesium bromide and reduction of the resulting 2-acylpyrrole 36 yielded [7'-²H](7'*R*)-2-undecylpyrrole 22.⁴ Using TBS-protected *S*-glycidol in place of 26, [7'-²H](7'*S*)-2-undecylpyrrole 23 was synthesized via an analogous sequence of transformations.

Feeding of Deuterium-Labeled Pyrroles 22 and 23, and MBC 3, to *S. albus* Expressing *redHG*. LC-MS analysis of mycelial extracts of *S. albus* expressing *redHG* that had been fed with [7'-²H](7'*S*)-2-undecylpyrrole 23 and MBC 3 showed that 92% of the resulting streptorubin B 2 is deuterium labeled (Figure 4). In contrast, 77% of the deuterium label is lost in streptorubin B 2 resulting from the feeding of [7'-²H](7'*R*)-2-undecylpyrrole 22 and MBC 3 to *S. albus* expressing *redHG* (Figure 4). These data show that the *pro-R* hydrogen atom is preferentially abstracted from C-7' in the RedG-catalyzed conversion of undecylprodigiosin 1 to streptorubin B 2.

Comparison of chromatograms resulting from monitoring of absorbance at 533 nm (corresponding to the λ_{\max} for the protonated 4-methoxypyrrrolyldipyrromethene chromophore of 1 and 2) showed that the amount of streptorubin B 2 produced relative to undecylprodigiosin 1 was much lower in the experiment utilizing [7'-²H](7'*R*)-2-undecylpyrrole 22 than in that using [7'-²H](7'*S*)-2-undecylpyrrole 23 (see the Supporting Information). This is consistent with a primary kinetic isotope effect for the abstraction of the deuterium atom from [7'-²H](7'*R*)-undecylprodigiosin, which may account for the discrepancy in the extent of deuterium retention in streptorubin B 2 when [7'-²H](7'*S*)-2-undecylpyrrole 23 was used compared with the amount of deuterium loss from streptorubin B 2 when [7'-²H](7'*R*)-2-undecylpyrrole 22 was employed.

Mechanistic Implications. The observed production of 5-hydroxypentylprodigiosin 17 when 2-(*S*-pentoxypentyl)-pyrrole 10 and MBC 3 are fed to *S. albus* expressing *redHG* is consistent with a RedG catalytic mechanism involving oxidation of C-7' prior to pyrrole functionalization. Presumably, 17 arises via abstraction of a hydrogen atom at C-7' of 10 by the Fe(III)OOH complex 6 to form a radical (or cation) that is




Figure 5. Mechanistic interpretation of the results from feeding of 2-(5-pentoxypentyl)-pyrrole **10** and stereoselectively deuterium labeled 2-undecylpyrroles **22** and **23**, along with MBC **3**, to *S. albus* expressing *redHG*. The C-7' radical formed by abstraction of the *pro-R* hydrogen atom in undecylprodigiosin **1** by the RedG ferric hydroperoxide complex **6** undergoes inversion prior to adding to C-4 of the proximal pyrrole, resulting in *S*-configured streptorubin **2**. The reduced reactivity of the radical formed via abstraction of a hydrogen atom from C-7' of 2-(5-pentoxypentyl)-pyrrole **10** by ferric hydroperoxide complex **6** prevents it from adding to C-4 of the proximal pyrrole. Instead, the radical “rebounds” onto the ferryl complex **8** to form the hemiketal complex **37**, which undergoes protonation-triggered collapse to liberate pentanal and form 5-hydroxypentylprodigiosin **17**.

unable to add to C-4 of the proximal pyrrole, either because the reactivity of the radical (cation) toward the pyrrole is reduced by the neighboring oxygen lone pair, or because of repulsion between the lone pairs of the C-6' oxygen atom and the pyrrole π -face (or both). The Fe(IV)O(OH₂) complex **8** formed by hydrogen abstraction from **10** by **6** reacts with the C-7' radical instead, forming the hemiketal complex **37**, which upon protonation can collapse with loss of *n*-pentanal to give 5-hydroxypentylprodigiosin **17** (Figure 5). Similar dealkylation reactions are known to be catalyzed by other members of the Rieske nonheme iron-dependent oxygenase family, such as Dicamba monooxygenase, which demethylates Dicamba (3, 6-dichloro-2-methoxybenzoate) via hydroxylation to form a hemiacetal that collapses with elimination of formaldehyde.²³

Analysis of deuterium incorporation into streptorubin **2** resulting from feeding of stereoselectively deuterium labeled 2-undecylpyrroles **22** and **23**, together with MBC **3**, to *S. albus* expressing *redHG* shows that RedG abstracts predominantly the *pro-R* hydrogen atom from C-7' of undecylprodigiosin **1**. Together with the previous observation that streptorubin **2** produced by *S. coelicolor* has predominantly the 7'*S* configuration,⁵ this result shows that the RedG-catalyzed C–C bond forming reaction proceeds with inversion of configuration at C-7' (Figure 5). In contrast, carbon-heteroatom bond formation in oxidative cyclizations catalyzed by other nonheme iron-dependent enzymes, such as isopenicillin N synthase and clavaminatase synthase proceeds with retention of configuration at the carbon atom undergoing functionalization.^{24,25} This stereochemical outcome is observed because the heteroatom participating in oxidative cyclization is ligated *cis* to the oxygen binding site of the nonheme iron center in these enzymes.^{26,27}

The inversion of configuration at C-7' in the RedG-catalyzed reaction suggests that the alkyl chain of the substrate may be juxtaposed between its 4-methoxypyrrrolyldipyrromethene moiety and the nonheme iron center (Figure 5), preventing the former from coordinating to the latter. Such an arrangement uses the alkyl chain to “insulate” the nonheme iron center

from 4-methoxypyrrrolyldipyrromethene moiety, which could prevent its electron rich π -system from being directly oxidized by the Fe(III)OOH complex **6**. However, this arrangement would probably place the Fe(IV)O(OH₂) complex **8** too far away from the hydrogen atom that needs to be removed from intermediate **9** to produce streptorubin **2** for this to occur via direct abstraction. This hydrogen atom could be extracted directly if positioned near one of the oxygen atoms of the hydroperoxide ligand in the Fe(III)OOH complex **6**, but this would require a change in the conformation of the alkyl chain after abstraction of the *pro-R* hydrogen atom from C-7' to permit C–C bond formation. Alternatively, the orientation of intermediate **9** in the active site could change to allow direct hydrogen abstraction by the Fe(IV)O(OH₂) complex **8**, or indirect hydrogen abstraction from intermediate **9** by complex **8** could occur via an appropriately placed active site water molecule. It is even conceivable that an electron is transferred from intermediate **9** to complex **8**. The resulting C-5 cation intermediate could then be deprotonated by a basic residue within the active site, obviating the need for the nonheme iron center to be located near H-4 of the pyrrole. This C-5 cation intermediate could also be formed via hydride abstraction from C-7' by the Fe(III)OOH complex **6** to form a transient alkyl cation that reacts with C-4 of the pyrrole. Alternatively, the transient alkyl cation could result from reaction of intermediate **7** with complex **8**, installing a hydroxyl group that undergoes subsequent protonation to promote its departure. Further experiments will be required to discriminate between these various possibilities.

CONCLUSIONS

The results reported herein not only illuminate the mechanism of the RedG-catalyzed oxidative carbocyclization of undecylprodigiosin **1** to streptorubin **2**, but also provide a starting point for understanding and investigating the mechanism and stereochemistry of similar oxidative carbocyclization reactions catalyzed by RedG homologues in the biosynthesis of other prodiginine alkaloids. For example, McpG has been shown to




Figure 6. Proposed catalytic mechanism for MarG. The key difference from the RedG catalytic mechanism is that the intermediate **41** resulting from addition of the alkyl radical to the 4-methoxypyrrrolyldipyrromethene π -system is hypothesized to react with the Fe(IV)O(OH₂) complex **8** to produce the hydroxylated product **42**.




Figure 7. Proposed mechanism for formation of the premarineosins **39** from intermediate **42**. The premarineosins are reduced to form the marineosins **40** by MarA.

catalyze the conversion of undecylprodigiosin **1** to metacycloprodigiosin **5** with opposite absolute stereocontrol, resulting in the 9'*R* absolute configuration (Figure 1).¹⁰ It seems reasonable to propose that this also proceeds with inversion of configuration, for the reasons discussed above, but involves abstraction of the *pro-S* hydrogen atom from C-9'. This hypothesis could be tested by synthesizing both enantiomers of 2-undecylpyrrole stereoselectively deuterium-labeled at C-9' and analyzing whether the deuterium atom is retained when each is fed, along with MBC, to strains of *S. albus* expressing *mcpH* and *mcpG*.

Recently, Reynolds and co-workers reported that the RedG homologue MarG catalyzes the conversion of (10'*S*)-10'-hydroxyundecylprodigiosin **38** to the premarineosins **39** (Figures 6 and 7), which undergo MarA-catalyzed reduction to marineosins A and B **40** (Figure 7).^{28,29} We hypothesize that the MarG catalytic cycle involves abstraction of a hydrogen atom from C-8' of **38** by the Fe(III)OOH complex **6** (or an Fe(V)O(OH) complex) to yield a radical (or cation) that

attacks C-1'. The resulting delocalized radical **41** can be quenched by “rebounding” onto the Fe(IV)O(OH₂) complex **8** (rather than by hydrogen atom abstraction as in the RedG proposed catalytic mechanism) to give a product **42** that has undergone both oxidative carbocyclisation and hydroxylation (Figure 6). Elimination of water from **42**, followed by addition of the C-10' hydroxyl group to the π -system would yield the premarineosins **39** (Figure 7). Addition of the C-10' hydroxyl group to the alternative delocalized cation intermediate would yield the premarineosins **39** directly.

The gene cluster believed to direct the biosynthesis of deschlororoseophilin **43** and prodigiosin R1 **44** in *Streptomyces griseoviridis* contains four RedG homologues (RphG, RphG2, RphG3 and RphG4).³⁰ One of these (RphG3) is likely nonfunctional because it lacks the conserved Cys and His residues that ligate the [2Fe-2S] cluster in the N-terminal Rieske domain. Another presumably catalyzes C–C bond formation between C-4 and C-9' in 11'-dimethyl-undecylprodigiosin **45** to form prodigiosin R1 **44** via a very similar process




Figure 8. Proposed mechanism for the biosynthesis of deschlororoseophilin **43** and prodigiosin R1 **44** in *Streptomyces griseoviridis*.

to the McpG-catalyzed conversion of undecylprodigiosin **1** to metacycloprodigiosin **5** (Figure 8).¹⁰ The remaining two RedG homologues are hypothesized to be responsible for the conversion of **45** to deschlororoseophilin **43** (Figure 8). One is proposed to catalyze C–C bond formation between C-10' and C-4 in **45** to give **46** (Figure 8). The other is proposed to catalyze the conversion of **46** to **47** via an analogous process to the MarG-catalyzed conversion of (10'S)-10'-hydroxyundecylprodigiosin **38** to **42** (Figures 6 and 8). Rearrangement of intermediate **47** via ring opening, tautomerization, ring closure and elimination of ammonia would give deschlororoseophilin **43** (Figure 8).

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and ¹H NMR spectra of the Mosher's esters of alcohol **32** and its enantiomer. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.5b03994.

■ AUTHOR INFORMATION

Corresponding Author

*g.l.challis@warwick.ac.uk

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was funded by BBSRC through a doctoral training grant to the University of Warwick, the RSC Briggs Scholarship (to D.M.W.) and the European Commission (Framework 6 integrated project, contract no. 005224). The assistance of Dr. Ivan Prokes and Dr. Lijiang Song with acquisition of NMR and mass spectra, respectively, is gratefully acknowledged. Regan Thomson and Dennis Hu (Northwestern University) are thanked for helpful discussions.

■ REFERENCES

- (1) Fürstner, A. *Angew. Chem., Int. Ed.* **2003**, *42*, 3582–3603.
- (2) Papireddy, K.; Smilkstein, M.; Kelly, J. X.; Shweta; Salem, S. M.; Alhamadsheh, M.; Haynes, S. W.; Challis, G. L.; Reynolds, K. A. *J. Med. Chem.* **2011**, *54*, 5296–5306.

- (3) Tsao, S.-W.; Rudd, B. A. M.; He, X.-G.; Chang, C.-J.; Floss, H. G. *J. Antibiot.* **1985**, *38*, 128–131.
- (4) Mo, S.; Sydor, P. K.; Corre, C.; Alhamadsheh, M. M.; Stanley, A. E.; Haynes, S. W.; Song, L.; Reynolds, K. A.; Challis, G. L. *Chem. Biol.* **2008**, *15*, 137–148.
- (5) Haynes, S. W.; Sydor, P. K.; Corre, C.; Song, L.; Challis, G. L. *J. Am. Chem. Soc.* **2011**, *133*, 1793–1798.
- (6) Hu, D. X.; Clift, M. D.; Lazarski, K. E.; Thomson, R. J. *J. Am. Chem. Soc.* **2011**, *133*, 1798–1804.
- (7) For a recent review, see: Challis, G. L. *J. Ind. Microbiol. Biotechnol.* **2014**, *41*, 219–232.
- (8) Stanley, A. E.; Walton, L. J.; Kourdi-Zerikly, M.; Corre, C.; Challis, G. L. *Chem. Commun.* **2006**, 3981–3983.
- (9) Haynes, S. W.; Sydor, P. K.; Stanley, A. E.; Song, L.; Challis, G. L. *Chem. Commun.* **2008**, 1865–1867.
- (10) Sydor, P. K.; Barry, S. M.; Odulate, O. M.; Barona-Gomez, F.; Haynes, S. W.; Corre, C.; Song, L.; Challis, G. L. *Nat. Chem.* **2011**, *3*, 388–392.
- (11) Sydor, P. K.; Challis, G. L. *Methods Enzymol.* **2012**, *516*, 195–218.
- (12) Barry, S. M.; Challis, G. L. *ACS Catal.* **2013**, *3*, 2362–2370.
- (13) McDougal, P. G.; Rico, J. G.; Oh, Y.-I.; Condon, B. D. *J. Org. Chem.* **1986**, *51*, 3388–3390.
- (14) Travis, B. R.; Sivakumar, M.; Hollist, G. O.; Borhan, B. *Org. Lett.* **2003**, *5*, 1031–1034.
- (15) More, J. D.; Finney, N. S. *Org. Lett.* **2002**, *4*, 3001–3003.
- (16) Omura, K.; Swern, D. *Tetrahedron* **1978**, *34*, 1651–1660.
- (17) Bal, B. S.; Childers, W. E.; Pinnick, H. W. *Tetrahedron* **1981**, *37*, 2091–2096.
- (18) Ichimaru, N.; Murai, M.; Kakutani, N.; Kako, J.; Ishihara, A.; Nakagawa, Y.; Nishioka, T.; Yagi, T.; Miyoshi, H. *Biochemistry* **2008**, *47*, 10816–10826.
- (19) Freeman, P. K.; Hutchinson, L. L. *J. Org. Chem.* **1980**, *45*, 1924–1930.
- (20) Lipshutz, B. H.; Wilhelm, R. S.; Kozlowski, J. A.; Parker, D. J. *Org. Chem.* **1984**, *49*, 3928–3938.
- (21) Dale, J. A.; Mosher, H. S. *J. Am. Chem. Soc.* **1973**, *95*, 512–519.
- (22) The displacement of homochiral secondary tosylates with LiAlD₄ is known to proceed with clean inversion. See: Ashby, E. C.; DePriest, R. N.; Goel, A. B.; Wenderoth, B.; Pham, T. N. *J. Org. Chem.* **1984**, *49*, 3545–3556.
- (23) Jiang, W. Z.; Wilson, M. A.; Weeks, D. P. *ACS Chem. Biol.* **2013**, *8*, 1687–1691.
- (24) Baldwin, J. E.; Wan, T. S. *Tetrahedron* **1981**, *37*, 1589–1595.

- (25) Basak, A.; Salowe, S. P.; Townsend, C. A. *J. Am. Chem. Soc.* **1990**, *112*, 1654–1656.
- (26) Roach, P. L.; Clifton, I. J.; Hensgens, C. M. H.; Shibata, N.; Schofield, C. J.; Hajdu, J.; Baldwin, J. E. *Nature* **1997**, *387*, 827–830.
- (27) Zhang, Z.; Ren, J.; Stammers, D. K.; Baldwin, J. E.; Harlos, K.; Schofield, C. *J. Nat. Struct. Biol.* **2000**, *7*, 127–133.
- (28) Salem, S. M.; Kancharla, P.; Florova, G.; Gupta, S.; Lu, W.; Reynolds, K. A. *J. Am. Chem. Soc.* **2014**, *136*, 4656–4574.
- (29) Kancharla, P.; Lu, W.; Salem, S. M.; Kelly, J. G.; Reynolds, K. A. *J. Org. Chem.* **2014**, *79*, 11674–11689.
- (30) Kawasaki, T.; Sakurai, F.; Nagatsuka, S.; Hayakawa, Y. *J. Antibiot.* **2009**, *62*, 271–276.