


Design of Modified Amine Transfer Reagents Allows the Synthesis of α -Chiral Secondary Amines via CuH-Catalyzed Hydroamination

Dawen Niu and Stephen L. Buchwald*

Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, Massachusetts 02139, United States

S Supporting Information

ABSTRACT: The CuH-catalyzed hydroamination of alkenes and alkynes using a silane and an amine transfer reagent represents a simple strategy to access chiral amine products. We have recently reported methods to prepare chiral amines with high efficiency and stereoselectivity using this approach. However, the current technology is limited to the synthesis of trialkylamines from dialkylamine transfer reagents (R_2NOBz). When monoalkylamine transfer reagents [$RN(H)OBz$] were used for the synthesis of chiral secondary amines, competitive, nonproductive consumption of these reagents by the CuH species resulted in poor yields. In this paper, we report the design of a modified type of amine transfer reagent that addresses this limitation. This effort has enabled us to develop a CuH-catalyzed synthesis of chiral secondary amines using a variety of amine coupling partners, including those derived from amino acid esters, carbohydrates, and steroids. Mechanistic investigations indicated that the modified amine transfer reagents are less susceptible to direct reaction with CuH.



INTRODUCTION

Chiral amines are ubiquitous structural motifs found in many pharmaceutical agents, natural products, and catalysts for asymmetric synthesis (see Figure 1 for selected examples).






Figure 1. Representative natural products and pharmaceutical agents that feature a chiral amine motif.

As a result, general and selective methods for their synthesis have long been pursued.¹ Methodologies using resolution² or chiral auxiliaries³ have been established as reliable ways to procure these compounds. Significant progress has also been made in the development of catalytic, asymmetric methods for their preparation. Many of these reported catalytic methods are based on asymmetric transformations of imine, enamine, or enamide intermediates.⁴ Other strategies, such as asymmetric allylation


a) Traditional hydroamination



b) CuH catalyzed hydroamination: synthesis of chiral tertiary amines



c) This work: synthesis of chiral secondary amines



$R' = H$: generally poor yields
 $R' = NMe_2$: broad substrate scope, good yields

Figure 2. Hydroamination approaches to make α -chiral amines.

of amine nucleophiles⁵ and direct C–H bond insertion by a nitrenoid species,⁶ are important alternatives.

Received: May 26, 2015

Published: July 5, 2015





Figure 3. CuH-catalyzed hydroamination of styrenes for the formation of chiral secondary amines. ^aYields are determined using GC with dodecane as an internal standard; unless otherwise noted, CuH solution used in this study was prepared in a nitrogen-filled glovebox. ^bIsolated yields on 1 mmol scale (average of two runs); enantiomeric excesses (ee) were determined by chiral HPLC analysis; see Supporting Information for experimental details. ^cThree equivalents of HSi(OEt)₂Me was used, and 10e was added over 1.5 h.

Asymmetric hydroamination,⁷ the net stereoselective addition of a hydrogen atom and an amino group directly across a double bond, represents a particularly appealing strategy to prepare chiral amines. Typically, a hydroamination reaction entails the direct union of an alkene **1** with a primary or secondary amine nucleophile **2** in the presence of a catalyst (Figure 2a).⁷ Based on catalytic copper(I) hydride chemistry⁸ and recent developments in the copper-mediated amination of carbon-based nucleophiles,⁹ our group recently reported a mechanistically distinct approach toward asymmetric hydroamination¹⁰ (Figure 2b). In this technique, an olefin first undergoes asymmetric hydrocupration to provide an alkylcopper intermediate, which is then intercepted by a suitable electrophilic amine transfer reagent. Our laboratory has applied this hydroamination strategy to the synthesis of chiral tertiary alkylamines from styrenes,^{10a} 1,1-disubstituted alkenes,^{10b} vinylsilanes,^{10c} and alkynes.^{10d} Independently, Miura and co-workers

have reported a similar approach for the hydroamination of styrenes^{11a} and strained internal alkenes.^{11b}

To date, this approach to hydroamination^{10,11} has been limited to the synthesis of tertiary alkylamines with *O*-benzoyl-*N,N*-dialkylhydroxylamines (R₂NOBz, R = alkyl) as the dialkylamine transfer reagents. The expansion of this method to monoalkylamine transfer reagents to allow the direct preparation of chiral secondary amines would be of considerable interest. Herein we report the development of a copper-catalyzed hydroamination process to directly generate chiral, branched secondary amines **8** from styrenes (Figure 2c). One key factor in the development of this method is the design and use of a modified class of amine transfer reagents **7**, which improved the efficiency and generality of the transformation. Mechanistic studies indicate that use of the modified amine transfer reagent suppresses nonproductive consumption of the reagent by the copper hydride intermediate.

Scheme 1. Hydroamination Reaction in the Synthesis and Derivatization of Drugs^a

^aReactions performed on 0.5 mmol scale. Isolated yields are reported (average of two runs). Enantioselectivities and diastereoselectivities were determined by chiral HPLC or ¹H NMR analysis. Conditions A: (1) NH₂OH·HCl, pyridine; (2) NaBH₃CN, HCl in MeOH, MeOH/THF; (3) 4-(dimethylamino)benzoic acid, CDI, CH₂Cl₂. Conditions B: Pd(OAc)₂, SPhos, potassium vinyltrifluoroborate, K₂CO₃, dioxane/H₂O.

Use of 4-fluorostyrene yielded **11k**, which resembles the core structure of PF-05105679 (Figure 1), in 92% yield and 95% ee. Lastly, this process also permits the preparation of **11l–n**, which contain synthetically versatile aryl bromide. The successful formation of **11l** was somewhat surprising since related alkyl-copper intermediates had been shown to undergo rearrangement to afford the arylcopper species.¹⁷

We next investigated the scope of the amine transfer reagents that could be employed (Table 1). The amine transfer agents used in this study were prepared from the corresponding primary hydroxylamine **15** and 4-(dimethylamino)benzoic acid via condensation effected by 1,1'-carbonyldiimidazole (CDI).¹⁸ As summarized in Table 1, amine transfer reagents with secondary or tertiary alkyl group substituents are competent substrates, delivering products **17a–d** in high yields and enantioselectivities. The use of chiral amine transfer agents afforded products with a high level of diastereoselectivity. The configuration of the newly generated stereocenter was determined by ligand employed (**17c** and **17d**). When either racemic ligand or racemic electrophile was used, a near unity ratio of diastereomers was formed

(see Supporting Information). These results are consistent with the diastereoselectivity of the hydroamination process being under catalyst control.

We found that amine transfer reagents prepared from α -amino esters¹⁹ were competent substrates as well, and gave the *N*-monoalkylated amino esters with high levels of stereocontrol (**17e–l**, Table 1). Amino esters spanning a range of steric and electronic properties could be utilized (**17e–j**). Importantly, the protecting groups on the amino esters could be methyl (**17e,f** and **17j**), benzyl (**17g**), or *tert*-butyl (**17h,i**), allowing a variety of choices for the selection of downstream deprotection methods. No epimerization of the labile stereocenter adjacent to the carbonyl group was observed, reflecting the overall mildness of the reaction system. Not surprisingly, use of different enantiomers of the ligand led to the formation of different diastereomers (**17h** and **17i**), again supporting a catalyst-controlled stereodetermining step. Furthermore, an amine transfer reagent derived from a quaternary α -amino ester could also be employed, giving **17j** in excellent yield and enantioselectivity. This method could also transform a vinylsilane^{10c} to the α -aminosilane (**17k**), a class of




Figure 4. Relative rates of the reactions between LCuH and different amine transfer agents. Si* = Si(OEt)₂Me. **Conditions A:** a 0.6 mL of a stock solution made from Cu(OAc)₂ (3.6 mg), (R)-DTBM-SEGPHOS (26 mg), PPh₃ (11.6 mg), HSi(OEt)₂Me (0.32 mL, 2.0 mmol), and THF-*d*₈ (1.0 mL) is used. The progress of these experiments was monitored by ¹H NMR.

building blocks often employed for the synthesis of peptidomimetics.²⁰ Finally, this hydroamination reaction can be integrated into a cascade sequence, delivering cyclic product **17l** after in situ alkylation of the intermediate secondary amine.

To further demonstrate the utility of this methodology, we applied it in the context of drug molecule synthesis (Scheme 1). For example, this method was applied to the synthesis of Sensipar (**21**), a drug used to treat secondary hyperparathyroidism. Commercially available aldehyde **18** was converted to amine transfer reagent **19** and then subjected to hydroamination conditions in the presence of 1-vinylnaphthalene (**20**) to give **21** in 81% yield and 89% ee (Scheme 1a). This methodology was also applied to the derivatization of commercial pharmaceuticals. For instance, chlorpromazine (**22**), an antipsychotic, could be converted to vinylarene **23**,²¹ which then underwent hydroamination with L-valine-derived amine transfer reagent **24** to yield **25** (Scheme 1b). In a similar fashion, loratadine (**26**), an antihistamine drug, could be converted to **27** and then coupled with **29**, an amine transfer reagent derived from an estrone derivative **28**, to afford the conjugated product **30** in 85% yield and >20:1 dr (Scheme 1c). Lastly, vinylarene **32** made from tufnil (**31**), a nonsteroid anti-inflammatory drug, could successfully couple with **34**, an amine transfer reagent prepared from a glucose derivative **33**, to afford **35** in 73% yield and 17:1 dr (Scheme 1d).

MECHANISTIC STUDIES

Competition experiments were performed to investigate the role of the modified amine transfer reagents used in this study. We had hypothesized that the narrow substrate scope of the CuH-catalyzed hydroamination reaction using monoalkylamine transfer agents [e.g., BnN(H)OBz] was due to the susceptibility of these reagents toward direct, nonproductive reduction by LCuH. To address this, we conducted a competition experiment by exposing a 1:1 mixture of a pair of mono- and dialkylamine transfer reagents, **10a** and **36**, to HSi(OEt)₂Me and copper catalyst in THF-*d*₈ in the absence of styrene and monitored the consumption of these two reagents by ¹H NMR spectroscopy (Figure 4a). We found that LCuH was capable of directly reacting with the amine transfer reagents, and over 80% of the monoalkylamine transfer agent **10a** was consumed within 1 h to

give BnNH₂ and the corresponding silylated benzoyl ester (**37**).²² In contrast, only a trace (<5%) of the dialkylamine transfer agent **36** was consumed during the same period of time.²³ We then subjected a 1:1 mixture of a pair of monoalkylamine transfer reagents, **10a** (parent benzoate) and **10e** [4-(dimethylamino)benzoate], to identical conditions as described above (Figure 4b). In this case, both **10a** and **10e** were gradually consumed in the reaction system, giving the corresponding silylated esters (**37** and **38**) as products. Importantly, we found that the modified amine transfer reagent **10e** was consumed at a considerably slower rate than was **10a**. The higher stability of the modified monoalkylamine transfer reagents toward direct reaction with LCuH reaction is consistent with the increased substrate scope seen using the 4-(dimethylamino)benzoate-derived amine transfer agents.

CONCLUSION

In conclusion, we have designed a new type of amine transfer reagent that possesses a 4-(dimethylamino)benzoate group. The use of these reagents enabled the development of a general method to directly convert styrenes to chiral secondary amines. This process was applicable to mono- and disubstituted styrenes and allowed the use of a variety of functionalized, structurally diverse amine transfer reagents, including those derived from carbohydrates, steroids, and amino acid esters. The utility of this reaction was highlighted by its application to the synthesis of pharmaceutically important drugs as well as the conjugation of other ones. Competition experiments have revealed that, relative to the corresponding *O*-benzoyl-*N*-alkyl hydroxylamines, the modified amine transfer reagents (*O*-[4-(dimethylamino)benzoyl]-*N*-alkyl hydroxylamines) are less susceptible to direct reaction with LCuH. The information gained from this study should prove useful in the design and development of other CuH-catalyzed processes.²⁴

ASSOCIATED CONTENT

Supporting Information

Experimental procedures and characterization data for all compounds. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.5b05446.

AUTHOR INFORMATION

Corresponding Author

*sbuchwal@mit.edu

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors acknowledge the National Institutes of Health for financial support (GM58160). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We thank Dr. M.T. Pirnot and Dr. Y. Wang for assistance in the preparation of this manuscript.

REFERENCES

(1) Nugent, T. C., Ed. *Chiral Amine Synthesis: Methods, Developments and Applications*; Wiley-VCH: Weinheim, Germany, 2010.

(2) Turner, N. J.; Carr, R. *Biocatalytic Routes to Nonracemic Chiral Amines*. In *Biocatalysis in the Pharmaceutical and Biotechnology Industries*; Patel, R. N., Ed.; CRC Press LLC: Boca Raton, FL, 2006; pp 743–755.

(3) Robak, M. T.; Herbage, M. A.; Ellman, J. A. *Chem. Rev.* **2010**, *110*, 3600.

(4) (a) Nugent, T. C.; El-Shazly, M. *Adv. Synth. Catal.* **2010**, *352*, 753. (b) Xie, J.; Zhu, S.; Zhou, Q. *Chem. Rev.* **2011**, *111*, 1713. (N) Tufvesson, P.; Lima-Ramos, J.; Jensen, J. S.; Al-Haque, N.; Neto, W.; Woodley, J. M. *Biotechnol. Bioeng.* **2011**, *108*, 1479. (c) Gopalaiiah, K. *Chem. Rev.* **2013**, *113*, 3248. (d) Feng, X.; Du, H. *Tetrahedron Lett.* **2014**, *55*, 6959. (e) Bloch, R. *Chem. Rev.* **1998**, *98*, 1407. (f) Kobayashi, S.; Ishitani, H. *Chem. Rev.* **1999**, *99*, 1069. (g) Friestad, G. K.; Mathies, A. K. *Tetrahedron* **2007**, *63*, 2541. (h) Yamada, K.; Tomioka, K. *Chem. Rev.* **2008**, *108*, 2874. (i) Kobayashi, S.; Mori, Y.; Fossey, J. S.; Salter, M. M. *Chem. Rev.* **2011**, *111*, 2626. (j) Yus, M.; González-Gómez, J. C.; Foubelo, F. *Chem. Rev.* **2011**, *111*, 7774. (k) Parmar, D.; Sugiono, E.; Raja, S.; Rueping, M. *Chem. Rev.* **2014**, *114*, 9047.

(5) (a) Johannsen, M.; Jørgensen, K. A. *Chem. Rev.* **1998**, *98*, 1689. (b) Helmchen, G. *Iridium-Catalyzed Asymmetric Allylic Substitutions*. In *Iridium Complexes in Organic Synthesis*; Oro, L. A., Claver, C., Eds.; Wiley-VCH: Weinheim, Germany, 2009; pp 211–250. (c) Hartwig, J. F.; Stanley, L. M. *Acc. Chem. Res.* **2010**, *43*, 1461. (d) Liu, W.-B.; Xia, J.-B.; You, S.-L. *Top. Organomet. Chem.* **2012**, *38*, 155. (e) Tosatti, P.; Nelson, A.; Marsden, S. P. *Org. Biomol. Chem.* **2012**, *10*, 3147.

(6) (a) Roizen, J. L.; Harvey, M. E.; Du Bois, J. *Acc. Chem. Res.* **2012**, *45*, 911. (b) Davies, H. M. L.; Manning, J. R. *Nature* **2008**, *451*, 417. (c) Müller, P.; Fruit, C. *Chem. Rev.* **2003**, *103*, 2905.

(7) (a) Muller, T. E.; Beller, M. *Chem. Rev.* **1998**, *98*, 675. (b) Hartwig, J. F. *Pure Appl. Chem.* **2004**, *76*, 507. (c) Hong, S.; Marks, T. J. *Acc. Chem. Res.* **2004**, *37*, 673. (d) Beller, M.; Seayad, J.; Tillack, A.; Jiao, H. *Angew. Chem., Int. Ed.* **2004**, *43*, 3368. (e) Hultsch, K. C. *Org. Biomol. Chem.* **2005**, *3*, 1819. (f) Muller, T. E.; Hultsch, K. C.; Yus, M.; Foubelo, F.; Tada, M. *Chem. Rev.* **2008**, *108*, 3795. (g) Hultsch, K. C. *Adv. Synth. Catal.* **2005**, *347*, 367. (h) Huang, L.; Arndt, M.; Goofen, K.; Heydt, H.; Goofen, L. J. *Chem. Rev.* **2015**, *115*, 2596.

(8) (a) Appella, D. H.; Moritani, Y.; Shintani, R.; Ferreira, E. M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1999**, *121*, 9473. (b) Hughes, G.; Kimura, M.; Buchwald, S. L. *J. Am. Chem. Soc.* **2003**, *125*, 11253. (c) Rainka, M. P.; Aye, Y.; Buchwald, S. L. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, *101*, 5821.

(9) (a) Berman, A. M.; Johnson, J. S. *J. Am. Chem. Soc.* **2004**, *126*, 5680. (b) Berman, A. M.; Johnson, J. S. *J. Org. Chem.* **2006**, *71*, 219. (c) Campbell, M. J.; Johnson, J. S. *Org. Lett.* **2007**, *9*, 1521. (d) Rucker, R. P.; Whittaker, A. M.; Dang, H.; Lalic, G. *Angew. Chem., Int. Ed.* **2012**, *51*, 3953. (e) Yan, X.; Yang, X.; Xi, C. *Catal. Sci. Technol.* **2014**, *4*, 4169. (f) Erdik, E.; Ay, M. *Chem. Rev.* **1989**, *89*, 1947. (g) Barker, T. J.; Jarvo, E. R. *Synthesis* **2011**, *24*, 3954.

(10) (a) Zhu, S.; Niljianskul, N.; Buchwald, S. L. *J. Am. Chem. Soc.* **2013**, *135*, 15746. (b) Zhu, S.; Buchwald, S. L. *J. Am. Chem. Soc.* **2014**, *136*, 15913. (c) Niljianskul, N.; Zhu, S.; Buchwald, S. L. *Angew. Chem.,*

Int. Ed. **2015**, *54*, 1638. (d) Shi, S.; Buchwald, S. L. *Nat. Chem.* **2015**, *7*, 38.


(11) (a) Miki, Y.; Hirano, K.; Satoh, T.; Miura, M. *Angew. Chem., Int. Ed.* **2013**, *52*, 10830. (b) Miki, Y.; Hirano, K.; Satoh, T.; Miura, M. *Org. Lett.* **2014**, *16*, 1498.

(12) In contrast to the large number of successful examples using dialkylamine transfer reagents (R_2NOBz , $R \neq H$) in copper-mediated amination reactions, use of the analogous monoalkylamine transfer reagents $[RN(H)OBz]$ remains underdeveloped. For representative examples, see ref 9b and (a) Yotphan, S.; Beukeaw, D.; Reutrakul, V. *Tetrahedron* **2013**, *69*, 6627. (b) McDonald, S. L.; Wang, Q. *Angew. Chem., Int. Ed.* **2014**, *53*, 1867. (c) Matsuda, N.; Hirano, K.; Satoh, T.; Miura, M. *Org. Lett.* **2011**, *13*, 2860.

(13) (a) Ascic, E.; Buchwald, S. L. *J. Am. Chem. Soc.* **2015**, *137*, 4666–4669. (b) PPH_3 is used as a secondary ligand due to the observed beneficial effects it has on CuH-catalyzed reactions. This concept was developed by Lipshutz: Lipshutz, B. H.; Noson, K.; Chrisman, W.; Lower, A. *J. Am. Chem. Soc.* **2003**, *125*, 8779.

(14) Attempts to modify the properties of dialkylamine transfer reagents by adjusting the benzoate group have been reported by Johnson.^{9c}

(15) (a) Alkyl carboxylate-based amine transfer reagents **13** and **14** were also tested. Reagent **14** containing a bulky adamantyl group provided yields of **11b** and **11c** similar to that of 4-(dimethylamino)benzoate **10e**. Due to the availability of the 4-(dimethylamino)benzoic acid and its ease of removal after reaction,^{15b} 4-(dimethylamino)benzoates were used for most of this work. (b) 4-(Dimethylamino)benzoic acid is sparingly soluble in Et_2O , $EtOAc$, or CH_2Cl_2 but readily soluble in 2 M aqueous K_2CO_3 .



(16) Attempts to use **10e** in the hydroamination of β,β -disubstituted styrenes or terminal unactivated alkenes gave low yields of secondary amine products.

(17) (a) Grigg, R. D.; Van Hoveln, R.; Schomaker, J. M. *J. Am. Chem. Soc.* **2012**, *134*, 16131. (b) Van Hoveln, R. J.; Schmid, S. C.; Schomaker, J. M. *Org. Biomol. Chem.* **2014**, *12*, 7655.

(18) Smithen, D. A.; Mathews, C. J.; Tomkinson, N. C. O. *Org. Biomol. Chem.* **2012**, *10*, 3756.

(19) Several methodologies have been developed to convert α -amino acid esters into the corresponding primary N-hydroxylamines. For references, see: (a) Tokuyama, H.; Kuboyama, H.; Fukuyama, T. *Org. Synth.* **2003**, *80*, 207. (b) Wittman, M. D.; Halcomb, R. L.; Danishefsky, S. J. *J. Org. Chem.* **1990**, *55*, 1981. (c) Fukuzumi, T.; Bode, J. W. *J. Am. Chem. Soc.* **2009**, *131*, 3864.

(20) Sun, H.; Martin, C.; Kesselring, D.; Keller, R.; Moeller, K. D. *J. Am. Chem. Soc.* **2006**, *128*, 13761.

(21) Molander, G. A.; Brown, A. R. *J. Org. Chem.* **2006**, *71*, 9681.

(22) Formation of $BnNH_2$ and the silylated ester **37** could be detected by GC/MS and 1H NMR analysis.

(23) Reduction of the dialkylamine transfer agent **36** is much slower. Approximately, 60% of **36** was consumed after 20 h under described conditions.

(24) A manuscript detailing the use of related modified amine transfer reagents to effect hydroamination of unactivated internal alkenes is in press.