

# Biochemical and Structural Basis for Controlling Chemical Modularity in Fungal Polyketide Biosynthesis

Jaclyn M. Winter,<sup>†,#</sup> Duilio Cascio,<sup>§</sup> David Dietrich,<sup>||</sup> Michio Sato,<sup>⊥</sup> Kenji Watanabe,<sup>⊥</sup> Michael R. Sawaya,<sup>§</sup> John C. Vederas,<sup>||</sup> and Yi Tang<sup>\*,†,‡</sup>

<sup>†</sup>Department of Chemical and Biomolecular Engineering, University of California, Los Angeles, California 90095, United States

<sup>‡</sup>Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90095, United States


<sup>§</sup>Department of Energy (DOE) Institute for Genomics and Proteomics, University of California, Los Angeles, California 90095, United States

<sup>||</sup>Department of Chemistry, University of Alberta, Edmonton, Alberta T6G 2G2, Canada

<sup>⊥</sup>Department of Pharmaceutical Sciences, University of Shizuoka, Shizuoka 422-8526, Japan

## Supporting Information

**ABSTRACT:** Modular collaboration between iterative fungal polyketide synthases (IPKSs) is an important mechanism for generating structural diversity of polyketide natural products. Inter-PKS communication and substrate channeling are controlled in large by the starter unit acyl carrier protein transacylase (SAT) domain found in the accepting IPKS module. Here, we reconstituted the modular biosynthesis of the benzaldehyde core of the chaetoviridin and chaetomugilin azaphilone natural products using the IPKSs CazF and CazM. Our studies revealed a critical role of CazM's SAT domain in selectively transferring a highly reduced triketide product from CazF. In contrast, a more oxidized triketide that is also produced by CazF and required in later stages of biosynthesis of the final product is not recognized by the SAT domain. The structural basis for the acyl unit selectivity was uncovered by the first X-ray structure of a fungal SAT domain, highlighted by a covalent hexanoyl thioester intermediate in the SAT active site. The crystal structure of SAT domain will enable protein engineering efforts aimed at mixing and matching different IPKS modules for the biosynthesis of new compounds.



## INTRODUCTION


The modular, assembly line biosynthesis of polyketides and nonribosomal peptides exemplifies how nature assembles complex natural products from simple precursors.<sup>1</sup> While this elegant biosynthetic logic is almost universally applicable among bacterial polyketide synthases (PKSs), the eukaryotic fungal iterative polyketide synthases (IPKSs) have evolved to be more simplified and enigmatic.<sup>2</sup> Each fungal IPKS consists of a single module, of which the catalytic domains embedded within are used iteratively to construct a finished product. Two major types of chemically distinct fungal IPKS modules have been discovered, including the nonreducing IPKSs (NR-PKSs) that produce aromatic products such as bikaverin,<sup>3</sup> and the highly reducing IPKSs (HR-PKSs) that produce highly reduced polyketides including the cholesterol-lowering lovastatin.<sup>4</sup> The HR-PKSs are particularly complex as the domains responsible for  $\alpha$ -methylation and  $\beta$ -reduction are used permutatively during each iteration with precise yet unresolved programming rules to yield structurally diverse products.

The collaboration between fungal PKS modules is an important mechanism to combine biosynthetic capabilities

from different PKSs and represents the fungal equivalent of assembly line biosynthesis.<sup>4</sup> The most frequently observed chemical modularity is the combination of an upstream HR-PKS and a downstream NR-PKS, in which a reduced polyketide synthesized by the HR-PKS is transferred to the NR-PKS as a starter unit via the starter-unit acyl carrier protein-transacylase (SAT) domain and further elongated.<sup>5</sup> Such distribution of labor leads to a polyketide product that combines two or more chemically distinct substructures. Examples include the biosynthesis of norsolorinic acid (**1**),<sup>5a</sup> in which an NR-PKS accepts a hexanoyl chain from an upstream fatty acid synthase (FAS)-like module and synthesizes the anthraquinone core that is eventually morphed into aflatoxin.<sup>6</sup> In the biosynthesis of the resorcylic acid lactones (RALs) such as hypothemycin (**2**),<sup>7</sup> radicicol (**3**)<sup>8</sup> and zearalenone (**4**)<sup>9</sup> (Figure 1A), a highly reduced polyketide precursor synthesized by an HR-PKS is transferred to an NR-PKS and further elongated and cyclized by the NR-PKS to yield the RALs. Another large family of

Received: April 30, 2015

Published: July 14, 2015



**Figure 1.** Chemical modularity in building fungal polyketides. (A) Representative molecules. The red portion indicates reduced products of upstream synthases that are transferred to downstream NR-PKSs. (B) Proposed biosynthesis of chaetoviridin A (6) and chaetomugilin A (7). Domain abbreviations: ketosynthase (KS), malonyl-CoA:acyl carrier protein acyltransferase (MAT), dehydratase (DH), methyltransferase (MT), enoylreductase (ER), ketoreductase (KR), acyl carrier protein (ACP), starter-unit:ACP-transacylase (SAT), product template (PT), and reductive domain (R). 11-SNAC was synthesized and used to directly load the KS of CazM.

fungal polyketides that are derived from the chemically modular synthesis is the azaphilones such as asperfuranone (5)<sup>10</sup> and chaetoviridin A (6).<sup>11</sup> In these pathways, the NR-PKSs employ a C-terminal reductive (R) domain to release the products as benzaldehydes, with the benzyl ring typically undergoing further oxidative transformations.


The modularity of the two PKS system is therefore an attractive feature that can be engineered to make new biosynthetic products. Indeed, successful production of analogs of 1, 5 and assorted RALs have been achieved through the mixing-and-matching of different nonreducing and reducing PKS modules, as well as using precursor-directed biosynthesis in which chemically synthesized reduced precursors are used to prime the functions of the NR-PKS module.<sup>5b,12–14</sup> Notwithstanding the conceptual simplicity of the collaboration between modules, fundamental insights into how the modules communicate through protein–protein interactions and how substrates are recognized and transferred between modules are needed to enable more rational biosynthetic engineering strategies.

The *caz* biosynthetic cluster from *Chaetomium globosum* is responsible for the production of 6 and chaetomugilin A (7), and contains the NR-PKS CazM and HR-PKS CazF.<sup>11</sup> In vitro assays confirmed CazF synthesizes the (4*S*,5*R*)-5-hydroxy-4-methyl-3-oxohexanoyltriketide 8, which is transferred to cazisochromene 9 by the acyltransferase CazE to yield 6.<sup>11</sup> The biosynthesis of 9 is proposed to derive from cazaldehyde A (10), which appears to be the product of a collaborative HR-PKS and NR-PKS interaction. However, the presence of CazF as the lone HR-PKS in the *caz* gene cluster suggests that CazF may be involved in the biosynthesis of 10 in addition to 8. In this model, CazF is proposed to also synthesize the more reduced (*S*,*E*)-4-methyl-hex-2-enoyltriketide 11. The product 11 is attached to the ACP of CazF as a thioester and is transferred to CazM by the SAT domain to serve as the starter unit in the synthesis of 10. The proposed mechanism of the SAT-mediated chain transfer is shown in Figure 1B, in which

the acyl chain 11 is first transacylated to the active site cysteine of the SAT, followed by transfer to the phosphopantetheinyl thiol on the ACP domain of CazM. If the collaborative interactions between CazF and CazM in the synthesis of 10 can be established, this would reveal an intriguing example of chemical modularity in fungal PKS programming where an HR-PKS is capable of synthesizing two different reduced portions of a natural product.<sup>15</sup> This will also point to an important role of the CazM SAT domain, which must selectively transacylate 11 over the more polar 8 from the CazF ACP to the CazM ACP during the PKS crosstalk. Here, we report the reconstitution of the CazF/CazM interaction in the biosynthesis of 6, as well as the functional and structural characterization of the CazM SAT domain. This first X-ray crystal structure of the SAT domain from a fungal PKS will be highly useful for future biosynthetic engineering efforts.


## RESULTS AND DISCUSSION

To reconstitute the CazF and CazM interactions, we first aimed to produce CazM recombinantly from *Saccharomyces cerevisiae*. The entire gene coding sequence of *cazM* was originally annotated to be three genes: CHGG\_07645, CHGG\_07646 and CHGG\_07647 (Figure S1 and S2). cDNA sequencing revealed the presence of three extra nucleotides in the sequencing data: two in CHGG\_07645 and one in CHGG\_07646. Removal of these bases yielded one continuous protein-coding sequence. GenBank accession numbers for the corrected gDNA and mRNA sequences are KP764718 and KP771866, respectively. Once the correct uninterrupted coding sequence was in hand, CazM was expressed (299 kDa) as a C-terminal hexahistidine-tagged protein from *S. cerevisiae* BJS464-NpgA<sup>16,17</sup> at 2.2 mg/L. To assay if CazM alone can turnover polyketide products using acetyl-CoA as a starter unit, the protein was incubated with 2 mM each of acetyl-CoA, malonyl-CoA, *S*-adenosyl-L-methionine (SAM) and NADPH. Because CazM contains a reductive domain at its C-terminus, NADPH was added to the reaction to aid in the reductive offloading of



**Figure 2.** Modular CazF-CazM interaction and cazaaldehyde production. HPLC analysis ( $\lambda = 290$  nm) of polyketide products when (i) CazM was incubated with acetyl-CoA, malonyl-CoA, SAM and NADPH; (ii) CazM was incubated with CazF, malonyl-CoA, SAM and NADPH; (iii) CazM was incubated with 11-SNAC, malonyl-CoA, SAM and NADPH; (iv) CazM was incubated with 11-SNAC, malonyl-CoA and SAM; and (v) CazM was coexpressed with CazF in *S. cerevisiae* BJS464-NpgA. Two benzaldehydes **10** and **12** were observed.

**Scheme 1. Synthesis of 11-SNAC Using the SNAC- HWE Reagent **13** and 2-(S)-Methyl-1-butanol **14****




any polyketide product from the megasynthase. After incubation, no products were detected in the organic extract by LC–MS analysis suggesting that CazM requires a priming acyl substrate to initiate its PKS function (Figure 2).

We next examined the modular collaboration between CazF and CazM by mixing equimolar amounts of CazM and CazF, which was previously expressed from *S. cerevisiae*,<sup>11</sup> with 2 mM each of malonyl-CoA, SAM and NADPH. LC–MS analysis of the organic extract showed the formation of two predominant products containing identical UV profiles that are indicative of benzaldehydes. While one of the products had the same mass as the expected **10**, the other compound differed in mass ( $m/z$ ) by +2 (Figure S8). To elucidate the structures of the two products, CazF and CazM were coexpressed in *S. cerevisiae* BJS464-NpgA (Figure S7) and the two compounds were purified in sufficient quantity for complete 1D and 2D NMR spectroscopy (Figures S8–S16). The two compounds were confirmed to be **10** (1 mg/L) and cazaaldehyde B (**12**, 0.5 mg/L). While the formation of **10** was expected and confirms the successful collaboration between CazF and CazM, formation of **12**, which underwent enoyl reduction in the reduced portion of the molecule, was unexpected. Compound **12** is most likely a shunt product, as no analogue of **6** containing the corresponding reduced moiety has been isolated from *C.*

*globozum*. A variety of reduction assays were performed and verified that the conversion of **10** to **12** is CazM-dependent and not spontaneous (Figure S17).

To verify that CazF synthesizes the acyl substrate 11-CazF, we prepared the *N*-acetyl cysteamine thioester (SNAC) **11** mimic, which is expected to directly prime the ketosynthase (KS) domain of CazM for further chain elongation.<sup>8</sup> 11-SNAC was prepared from a 3-step sequence using the SNAC-HWE reagent **13** and 2-(S)-methyl-1-butanol **14** (Scheme 1). Thioesterification of commercially available diethylphosphonoacetic acid **15** with SNAC provided the phosphonate thioester **13**.<sup>18</sup> Oxidation of **14** to the aldehyde **16** followed by a HWE reaction with **13** furnished the *trans* olefin 11-SNAC. Combining 2 mM 11-SNAC and 25  $\mu$ M CazM in the presence of 2 mM each of malonyl-CoA, SAM and NADPH similarly led to the synthesis of both **10** and **12** (Figure 2). When NADPH was omitted from the reaction, no production of **10** or **12** could be detected indicating that CazM requires NADPH for the reductive offloading of its polyketide product. Altogether, these results confirm the product of the CazF module to be **11**, which is transferred to CazM for four additional rounds of chain elongation, cyclization and reductive release (Figure 1B). The formation of the side product **12** using 11-SNAC also suggests



**Figure 3.** Trans SAT complementation assays. HPLC analysis ( $\lambda = 290$  nm) of **10** synthesized when (i) CazM H277A was incubated with CazF, malonyl-CoA, SAM and NADPH; (ii) CazM H277A was incubated with 11-SNAC, malonyl-CoA, SAM and NADPH; (iii–vi) CazM H277A was incubated with CazF, malonyl-CoA, SAM, NADPH and standalone SAT domains.

that the extra reduction to yield **12** must take place on CazM, most likely by a previously unknown function of its R domain.


CazF was previously confirmed to also synthesize **8** that is recognized by CazE.<sup>11</sup> However, in the presence of CazM, no product derived from **8** can be detected, hence pointing to a strict preference for the more reduced polyketide **11** by the SAT domain of CazM. The CazM SAT domain shows sequence similarity to AT domains from FASs and bacterial modular PKSs.<sup>19</sup> The active site Cys155 found within the motif GxCxG and His277 are proposed to form the catalytic dyad<sup>5,14,19–21</sup> and were targeted for mutagenesis. While CazM mutants C155A and C155S were not solubly expressed from *S. cerevisiae* BJS464-NpgA, the H277A mutant was expressed at a comparable yield to wild-type CazM. In vitro assays with CazM H277A and CazF did not produce **10** and **12** (Figure 3). Precursor feeding of the CazM H277 mutant with 11-SNAC, however, restored the biosynthesis of **10** and **12**, thereby confirming the remaining domains of CazM were fully functional. Therefore, a functional SAT domain is absolutely required for the modular synthesis of **10**.

To test if a standalone SAT domain can complement the CazM SAT<sup>0</sup> mutant, we solubly expressed and purified the 46 kDa domain containing the linker region between SAT and KS domains from *Escherichia coli* BL21 (DE3). We reasoned that a functional, free-standing SAT domain will enable facile biochemical assays of mutants, as well as mix-and-match experiments toward analogue generation. When equimolar ratios of CazM H277A, CazF and wild type SAT were combined, we detected the restored production of **10** and **12** at levels comparable to that of CazF and wild-type CazM (Figure 3). Using the *in trans* complementation system, we constructed the active site SAT mutants C155A, C155S and H277A. While it was expected that the C155A and H277A mutants would not complement the CazM SAT<sup>0</sup> mutant, the C155S mutant was also surprisingly inactive (Figure 3). Interestingly, the SAT domains of the NR-PKSs Hpm3, Rdc1 and PKS13, which are responsible for the biosynthesis of the RALs **5**, **6**, and **7**, respectively, all contain a GxSxG active site motif. Our complementation assay indicates that the active site thiol is

absolutely essential for catalysis by CazM and the labile thioester intermediate is preferred over the more stable oxyester.<sup>4</sup> Hence, the catalytic environment in the SAT domain is precisely tuned for the acyl transfer reactions shown in Figure 1B.

To gain atomic resolution into the acyl transfer step between CazF and CazM; especially the role of the cysteine and the acyl binding pocket, we solved the crystal structure of the standalone SAT domain to 1.6 Å using single anomalous dispersion (SAD) (Table S5). The SAT crystallized in the space group  $P2_12_12_1$  with one monomer per asymmetric unit. Similar to structures of malonyl-CoA:ACP transacylases (MCAT) from bacterial fatty acid synthases,<sup>22–24</sup> both *trans*-AT<sup>25</sup> and *cis*-AT domains in PKSs<sup>19,26–30</sup> (for example, see the avermectin loading AT domain in Figure 4B), the CazM SAT architecture is comprised of a large  $\alpha/\beta$ -hydrolase core and a small ferredoxin-like subdomain containing a four-stranded antiparallel  $\beta$ -sheet and two distal  $\alpha$ -helices (Figure 4A).

We next performed cocrystallization of the SAT domain with either *trans*-2-hexenoic acid or hexanoyl-CoA, which served as mimics of the triketide substrate **11**. Serendipitously, the cocrystallized structure of the SAT with hexanoyl-CoA showed well-defined electron density for a hexanoyl thioester at the active site Cys and was solved by molecular replacement to 1.4 Å (Figure 4A; Table S5). The active site cavity is located at the  $\alpha/\beta$ -hydrolase and ferredoxin interface and was calculated to have a solvent accessible area of 238.49 Å<sup>3</sup> using CASTp<sup>31</sup> (Figure 4C). Predominantly hydrophobic amino acids line the cavity consistent with selectivity for a more reduced ligand (Figure 4D). These residues most likely dictate the preference of the SAT for the triketide **11** over the more oxidized **8**, both of which can be synthesized by CazF (Figure 1B). Residues G105, L159, A184, G188, A187, I191, and I337 line the bottom of the cavity and are all located within 5 Å of the distal carbon in the hexanoyl chain. These residues therefore limit the length of the starter unit to a triketide and potentially a tetraketide. Additionally, the imidazole side chain of His22 is in close proximity to the  $\beta$ -carbon of the hexanoyl substrate and could further sterically contribute to the observed selectivity toward



**Figure 4.** Crystal structure of the CazM SAT domain. (A) The monomeric SAT domain consists of a large subdomain containing an  $\alpha/\beta$ -hydrolase-like fold and a small subdomain containing a ferredoxin-like fold. (B) Structure of the avermectin AT loading domain Ave-AT<sup>0</sup> (PDB 4RL1).<sup>19</sup> The two teal helices indicate a change in orientation compared to  $\alpha J$  and  $\alpha P$  in the SAT structure and the active site S120 is shown as a stick. (C) Electrostatic surface representation of the SAT domain showing a cross section of the cavity leading to the active site and bound hexanoyl. (D) Residues located within 5 Å of bound hexanoyl and distances of residues lining the bottom of the active site cavity to C6 of hexanoyl (yellow dashed lines). Polar contacts are shown as black dashes. (E) Active site residue comparison between the *apo* and hexanoyl-bound SAT. The *apo* SAT residues are shown in gray, whereas the hexanoyl-bound residues are shown in purple. Bound hexanoyl is shown in ball and stick form. (F) Electrostatic surface maps of the *apo*-SAT (left) and CazF ACP (right) docking interfaces. Colors range from blue (positive) to white to red (negative). The homology model for CazF ACP was generated using I-TASSER<sup>34</sup> and docking simulations were performed with PatchDock.<sup>35</sup> The ACP is rotated 180° such that A–C on the SAT and ACP interfaces should match up. On the SAT, A is R374, B is R243 and C is R366. On the ACP, A is N61, B is E58 and C is D31. (G) The homology model of CazF ACP (colored in yellow) was used as a ligand protein to dock with the *apo* SAT structure (colored in white). Helix  $\alpha P$  of the SAT is highlighted in purple and surface residues that may be responsible for protein–protein interaction are shown as sticks. The active site Ser in the ACP and Cys in the SAT are in bold. All electrostatic surface maps were generated using PyMOL.<sup>32</sup>

the reduced  $\beta$ -methylene carbon found in **11** over the more bulky  $\beta$ -ketone present in **8**. Hence, capturing the thioester intermediate in the crystal structure not only confirmed the role of the active site cysteine, but also revealed factors that contribute to the SAT acyl substrate selectivity.

Comparison of the *apo*-SAT and hexanoyl-bound enzyme reveals a subtle conformational change in the active site. After the attachment of hexanoyl, the rotamer conformation of Cys155 located on the *N*-terminus of  $\alpha H$  changes and the loop connecting secondary structural elements  $\beta 1$  and  $\alpha B$  moves toward the active site (Figure 4E). Catalytic His277 acts as a general acid/base catalyst in the transthioesterification reaction

where  $N\epsilon$  can deprotonate the thiol in Cys155. The carbonyl backbone of Asn340 helps polarize His277 and has a hydrogen bond distance of 2.7 Å to  $N\delta$  of His277 in both *apo* and hexanoyl-bound structures (Figure 4E). Based on the conserved FGDQ(S/T) sequence in PksA, it was proposed that glutamine is involved in formation of the oxyanion hole.<sup>5</sup> The corresponding sequence from the CazM SAT consists of FAPHV and the backbone amides of His22 and Ala156 are most likely responsible for stabilizing the transition state as observed in the crystal structure.

While the majority of amino acids lining the acyl binding site are hydrophobic and select for a more lipid-like ligand, the basic

residues R243, R366 and R374 located by the entrance to the channel may be involved in inter- and intramolecular docking with the acidic ACP domains from CazF and CazM, respectively. In the published MCAT and AT structures, positively charged surface residues have been suggested to facilitate ACP binding. To provide further insight into residues responsible for protein–protein interaction, a homology model of the CazF ACP was obtained using I-TASSER.<sup>34</sup> The ACP domain from module 2 of 6-deoxyerythronolide B synthase (DEBS) (PDB: 2JU1)<sup>36</sup> was selected as the top threading template. The CazF ACP homology model was used as a ligand molecule to dock with the *apo* SAT structure. The lowest energy docking models placed the ACP in the cleft between the  $\alpha/\beta$  hydrolase and ferredoxin-like subdomains (Figure 4F). This placement is similar to docking models generated for the DEBS AT3 and ACP3,<sup>37</sup> the iterative type I enediyne synthase AT<sub>Dyn10</sub> and ACP<sub>Dyn1</sub><sup>30</sup> and the *trans*-AT kirromycin synthase KirCII and ACP5<sub>Kir</sub>.<sup>38</sup> The orientation of the ACP homology model places acidic residues D31 and E58 in close proximity to SAT residues R243 and R366 (Figure 4G). R243 is positioned at the entrance to the catalytic cavity and its side chain could make a salt bridge with the carboxylate group of either an Asp or Glu residue on the ACP. Interestingly, in the SAT structure, the orientation of helix  $\alpha$ P is parallel to  $\alpha$ O and  $\alpha$ J is parallel to  $\alpha$ K. In the majority of MCAT and AT structures, such as the MCATs from *Streptomyces coelicolor* A3(2),<sup>22</sup> *Helicobacter pylori*,<sup>23</sup> and *Xanthomonas oryzae pv oryzae*,<sup>24</sup> and the ATs from the avermectin synthase,<sup>19</sup> DEBS modules 3<sup>26</sup> and 5,<sup>27</sup> the mycolic acid synthase PKS13,<sup>28</sup> and zwittermicin synthase ZmA,<sup>29</sup> these helices are perpendicular to each other (Figure 4B and Figure S19). Extensive mutagenesis studies with the KirCII AT and ACP5<sub>Kir</sub> domains performed by Williams and co-workers demonstrated that the perpendicular  $\alpha$ P helix is the primary anchoring point for the ACP substrate. In the CazM SAT structure,  $\alpha$ P contains Arg374 and the basic residue may aid in the inter- and intramolecular docking with negatively charged residues on ACP domains. The unusual parallel configuration of the helices in the SAT may be important in the recognition of two different ACP partners. Such interactions may be mapped from the covalent ACP trapping experiments that have recently found success in analyzing these weak protein–protein interactions.<sup>39</sup>

## CONCLUSIONS

We reconstituted the modular interaction between a HR-PKS and NR-PKS in the biosynthesis of **6**. Biochemical and structural insights into how the SAT domain facilitates module crosstalk and controls substrate selection were obtained. The structure of the first SAT domain can serve as a starting point for rational protein engineering efforts toward the biosynthesis of new fungal polyketides.

## EXPERIMENTAL SECTION

**Synthesis of 11-SNAC.** To a solution of 2-(*S*)-methyl-1-butanol **14** (0.039 mL, 0.34 mmol) in CDCl<sub>3</sub> (1 mL) and D<sub>2</sub>O (1 mL) was added KBr (4.0 mg, 0.04 mmol), TEMPO (~0.6 mg) and NaOCl (0.25 mL of a 1.5 M solution, 0.38 mmol). The reaction was stirred at room temperature for 1 h and the progress of the reaction was monitored by NMR. When completed, the mixture was washed with 1 M HCl (1.0 mL, containing 16.0 mg potassium iodide/mL), 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (1.0 mL) and water (1.0 mL). The organic layer was dried and filtered to provide the aldehyde **16** as the only product (based on NMR) and was used immediately in the HWE reaction. The SNAC-HWE reagent **13** was prepared as previously reported from

diethylphosphonoacetic acid **15**.<sup>18</sup> Compound **13** (81 mg, 0.30 mmol) was dissolved in dry THF (1.2 mL) and LiBr was added (46 mg, 0.50 mmol). Addition of trimethylamine (0.08 mL, 0.60 mmol) yielded the ylide as a white solid. Aldehyde **16** formed in the first step was added in CDCl<sub>3</sub>, and the reaction was stirred overnight at room temperature. The solvent was removed, and the residue was purified by silica gel chromatography (7:3, EtOAc-hexanes) to afford **11-SNAC** (21.0 mg, 27%, two steps): [ $\alpha$ ]<sub>D</sub> 3.96 ( $c = 1.45$ , CHCl<sub>3</sub>); IR (CHCl<sub>3</sub>, cast) 3290, 3077, 2963, 2875, 1663, 1632, 1552, 1459, 1438, 1404, 1286, 1052; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, ppm) 6.83 (dd, 1H,  $J = 15.6, 7.8$  Hz, =CH), 6.09 (dd, 1H,  $J = 15.6, 1.2$  Hz, =CH), 5.38 (br s, 1H, NH), 3.47 (q, 2H,  $J = 6.2$  Hz, –NHCH<sub>2</sub>), 3.09 (t, 2H,  $J = 6.3$  Hz, SCH<sub>2</sub>), 2.25–2.19 (m, 1H, CH), 1.96 (s, 3H, Ac–CH<sub>3</sub>), 1.44–1.41 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.06 (d, 3H,  $J = 6.7$  Hz, CH–CH<sub>3</sub>), 0.88 (t, 3H,  $J = 7.5$  Hz, CH<sub>2</sub>CH<sub>2</sub>) (Figure S5); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>, ppm) 191.4, 171.0, 152.5, 127.6, 40.7, 39.0, 29.6, 29.2, 24.1, 19.6, 12.5 (Figure S5); HR-ESIMS (C<sub>11</sub>H<sub>19</sub>NO<sub>2</sub>SNa)<sup>+</sup> 252.1029 (calc), 252.1024 (found) (Figure S6).

**CazM Reconstitution.** To reconstitute an intron-free *cazM*, five overlapping pieces were constructed (Figure S2). Piece two (2,150 bp) and piece four (3,018 bp) were amplified from genomic DNA using primer pairs CazM\_P2\_F with CazM\_P2\_R and CazM\_P4\_F with CazM\_P4\_R, respectively (Table S1). Pieces one, three and five were initially annotated to contain introns and were amplified from cDNA. Total RNA was extracted using the RiboPure Yeast kit (Ambion) from a three-d old malt static culture (10 g/L malt extract, 4 g/L yeast extract, 10 g/L D-glucose) incubated at 28 °C. Residual genomic DNA was digested with DNase (2 U/ $\mu$ L) (Invitrogen) at 37 °C for four h. cDNA was created for pieces one, three and five using the reverse primers CazM\_P1\_R, CazM\_P3\_R, and CazM\_P5\_R, respectively. The cDNA was used as template for PCR and pieces one (1313 bp), three (1187 bp) and five (1733 bp) were amplified using primer pairs CazM\_P1\_F with CazM\_P1\_R, CazM\_P3\_F with CazM\_P3\_R, and CazM\_P5\_F with CazM\_P5\_R, respectively. The five exons were cotransformed with the 2  $\mu$  URA3 expression plasmid into *S. cerevisiae* BJ5464-NpgA using an *S. c.* EasyComp Transformation kit (Invitrogen) yielding plasmid pJWT-37. A single transformant was used to inoculate 3 mL SD<sub>c</sub> media (0.5 g of Bacto Technical grade cassamino acids, 2 g of dextrose, and 88 mL of Milli-Q water) supplemented with 1 mL of adenine (40 mg/20 mL of Milli-Q water), 1 mL of tryptophan (40 mg/20 mL of Milli-Q water) and nitrogen base (1.7 g of nitrogen base without amino acids and without ammonium sulfate, 5 g of ammonium sulfate and 100 mL of Milli-Q water) and grown for 72 h with constant shaking at 28 °C. A 1 mL aliquot of the seed culture was used to inoculate 1 L of YPD (10 g of yeast extract, 20 g of peptone and 950 mL of Milli-Q water) supplemented with 1% dextrose and the culture was shaken at 28 °C for 72 h. The cells were harvested by centrifugation (3750 rpm at 4 °C for 10 min) and the cell pellet was resuspended in 30 mL of lysis buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>, 150 mM NaCl, 10 mM imidazole, pH 8.0). The cells were sonicated on ice in one min intervals until homogeneous. To remove cellular debris, the homogeneous mixture was centrifuged at 17 000 rpm for 1 h at 4 °C. The supernatant was passed through a 0.45  $\mu$ m filter and Ni-NTA agarose resin (2 mL) was added to the cleared lysate. The solution was incubated with gentle rotation at 4 °C for 16 h. Soluble CazM was purified by gravity-flow column chromatography using increasing concentrations of imidazole in buffer A (50 mM Tris-HCl, 500 mM NaCl, 20–250 mM imidazole, pH 7.9). Purified protein was concentrated and buffer exchanged into Buffer B (50 mM Tris-HCl, 2 mM EDTA, 100 mM NaCl, pH 7.9) using an Ultracel 100 000 MWCO centrifugal filter (Amicon Inc.) and stored in 10% glycerol. The sequence of an intron-free *cazM* (Figure S3) was confirmed using primers CazM\_seq1–CazM\_seq11 and its protein concentration was calculated to be 2.2 mg/L using the Bradford assay with BSA as a standard.

**CazM In Vitro Assays.** To assess the activity of CazM and what substrates could prime the enzyme, 25  $\mu$ M of protein was incubated at room temperature in a 100  $\mu$ L reactions containing 2 mM *S*-adenosyl-methionine chloride (SAM), 2 mM NADPH, 2 mM malonyl-CoA and either 2 mM acetyl-CoA or 2 mM 11-SNAC in 100 mM

phosphate buffer pH 7.4. After 8 h, the reactions were extracted two times with 200  $\mu$ L 99% ethyl acetate:1% acetic acid and the organic layer was dried by speedvac. The extract was then dissolved in 20  $\mu$ L of MeOH and analyzed on a Shimadzu 2010 EV LC–MS with a Phenomenex Luna 5  $\mu$  2.0  $\times$  100 mM C18 column using positive and negative mode electrospray ionization with a linear gradient of 5–95% MeCN:H<sub>2</sub>O over 30 min followed by 95% MeCN for 15 min and a flow rate of 0.1 mL/min. To assess whether the polyketide product could be released off the ACP without a reducing agent, the assay was repeated using 11-SNAC and omitting NADPH from the reaction. The reaction was incubated for 8 h at room temperature and extracted and analyzed as described above.

CazF was expressed and purified as published previously.<sup>11</sup> The activity of CazM with its partnering HR-PKS CazF was analyzed by incubating 25  $\mu$ M CazM, 25  $\mu$ M CazF, 2 mM malonyl-CoA, 2 mM NADPH, 2 mM SAM in 100 mM phosphate buffer pH 7.4 in a total volume of 100  $\mu$ L. The reaction was incubated at room temperature for 12 h and extracted two times with 200  $\mu$ L 99% ethyl acetate:1% acetic acid. The organic layer was dried using a speedvac and the extract was resuspended in 20  $\mu$ L of MeOH and analyzed by LC–MS as mentioned above. The CazF–CazM activity reaction was repeated but the wild-type CazM was replaced with the CazM H277A mutant (construction described in the Supporting Information). The reaction was incubated at room temperature for 12 h and then extracted and analyzed as described previously.

**In Vivo Production of Cezaldehyde A (10) and B (12).** Intact *cazF* was amplified using pJW07638 as template,<sup>11</sup> AccuPrimePfx DNA polymerase (Invitrogen) and primers *CazF\_Trp\_F* and *CazF\_Trp\_R* (Table S1). The 7863 amplicon was cotransformed with the 2  $\mu$  Trp expression plasmid into *S. cerevisiae* BJ5464-NpgA using an *S. c.* EasyComp Transformation kit (Invitrogen) to yield plasmid pJW41. pJW41 and pJW37 were then cotransformed into *S. cerevisiae* BJ5464-NpgA.<sup>16,17</sup> A single transformant was used to inoculate 3 mL of SD<sub>c</sub> media (0.5 g of Bacto Technical grade cassamino acids, 2 g of dextrose, and 89 mL of Milli-Q water) supplemented with 1 mL of adenine (40 mg/20 mL of Milli-Q water) and 10 mL of nitrogen base (1.7 g of nitrogen base without amino acids, 5 g of ammonium sulfate and 100 mL of Milli-Q water) and grown for 72 h with constant shaking at 28 °C. A 1 mL aliquot of the seed culture was used to inoculate 100 mL of YPD (1.0 g of yeast extract, 2.0 g of peptone and 95.0 mL of Milli-Q water) supplemented with 1% dextrose and the culture was shaken at 28 °C at 250 rpm. A 500  $\mu$ L aliquot was extracted with ethyl acetate containing 1% acetic acid at d 1, 2, 3, and 7 (Figure S7). The organic layer was dried using a speedvac and the extract was resuspended in 20  $\mu$ L of MeOH. The crude extract was analyzed in positive and negative mode on a Shimadzu 2010 EV liquid chromatography mass spectrometer (LC–MS) (gas flow set to 4.2 mL/min, drying temperature set to 200 °C, and capillary voltage set to 2000 V). A Luna C18(2) 100  $\times$  2 mm column was used with a flow rate of 0.1 mL/min and a linear solvent gradient of 5–95% MeCN:H<sub>2</sub>O containing 0.1% formic acid over a period of 30 min followed by 15 min of 95% MeCN:H<sub>2</sub>O.

**Purification of Cezaldehyde A (10) and B (12).** To isolate enough material for NMR characterization, 10  $\times$  100 mL cultures were extracted after 48 h. The mycelium and media were extracted with ethyl acetate containing 1% acetic acid, dried over anhydrous MgSO<sub>4</sub> and concentrated in vacuo. The crude extract was fractionated using reverse phase C18 flash column chromatography (Fisher Scientific, PrepSep C18 1 g/6 mL) with 20% MeCN:H<sub>2</sub>O, 40% MeCN:H<sub>2</sub>O, 60% MeCN:H<sub>2</sub>O, 80% MeCN:H<sub>2</sub>O, and 100% MeCN as the mobile phase. The fractions were analyzed in positive and negative mode on a Shimadzu 2010 EV LCMS using the same parameters as above. Compounds 10 and 12 eluted in the 40% MeCN:H<sub>2</sub>O fraction and were purified by HPLC (Beckman Coulter) using a Luna C18 column (250  $\times$  10 mm; 5  $\mu$ m particle size) and an isocratic condition of 55% MeCN:H<sub>2</sub>O containing 0.1% TFA with a flow rate of 2.0 mL/min. The *t<sub>R</sub>* of 10 and 12 were 28.5 and 29.5 min, respectively. Compounds 10 and 12 were isolated at 1 mg/L and 0.5 mg/L, respectively, and their chemical structures were elucidated from the spectroscopic data given in Tables S3 and S4, respectively.

**Expression of the SAT Domain.** The SAT domain of CazM (Figure S3) did not contain any introns and was amplified from *C. globosum* gDNA using primer pair SAT\_pHis8\_F and SAT\_pHis8\_R, which contained *EcoRI* and *NotI* restriction sites, respectively. The PCR product was digested with *EcoRI* and *NotI* and ligated into the pHis8 expression vector<sup>40</sup> yielding expression plasmid pJW49, which was transformed into *E. coli* BL21(DE3) cells. A single colony was used to inoculate 10 mL of Luria–Bertani (LB) media supplemented with 35  $\mu$ g/mL kanamycin and grown overnight with constant shaking at 37 °C. A 5 mL aliquot of the seed culture was used to inoculate a 2 L shake flask containing 500 mL of Terrific Broth (per liter: 24 g of yeast extract, 12 g of tryptone and 4 mL of glycerol) supplemented with kanamycin. The cells were shaken at 250 rpm at 37 °C to an OD<sub>600</sub> of 0.6. To induce protein expression, isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) was added to a final concentration of 0.4 mM and the culture was shaken at 16 °C for 22 h. The cells were harvested by centrifugation (3750 rpm at 4 °C for 25 min) and the cell pellet was resuspended in 25 mL of lysis buffer (50 mM Tris-HCl, 2 mM EDTA, 2 mM DTT, 500 mM NaCl, 5 mM imidazole, pH 7.9). The cells were sonicated on ice in 30 s intervals until homogeneous. To remove cellular debris, the homogeneous mixture was centrifuged at 17 000 rpm for 30 min at 4 °C. Ni-NTA agarose resin was added to the supernatant (1 mL) and the solution was incubated at 4 °C with gentle agitation for four h. Soluble octahistidyl-tagged SAT was purified by gravity-flow column chromatography using increasing concentrations of imidazole in buffer A (50 mM Tris-HCl, 500 mM NaCl, 20 mM–250 mM imidazole, pH 7.9). Purified SAT was concentrated and buffer exchanged into Buffer B (50 mM Tris-HCl, 2 mM EDTA, 100 mM NaCl, pH 7.9) using an Ultracel 30 000 MWCO centrifugal filter (Amicon Inc.) and stored in 10% glycerol. The protein concentration was calculated to be 43 mg/L by Bradford assay using BSA as a standard.

**Standalone SAT in Vitro Studies.** To determine if a standalone SAT domain could restore function to an SAT inactive CazM, 25  $\mu$ M of CazM H277A was incubated with 25  $\mu$ M of the standalone wild-type SAT domain in a 100  $\mu$ L reaction containing 2 mM SAM, 2 mM NADPH, 2 mM mal-CoA and 25  $\mu$ M CazF in 100 mM phosphate buffer pH 7.4. After 8 h, the reaction was extracted two times with 200  $\mu$ L 99% ethyl acetate:1% acetic acid and the organic layer was dried by speedvac. The extract was then dissolved in 20  $\mu$ L of MeOH and analyzed on a Shimadzu 2010 EV LC–MS with a Phenomenex Luna 5  $\mu$  2.0  $\times$  100 mM C18 column using positive and negative mode electrospray ionization with a linear gradient of 5–95% MeCN:H<sub>2</sub>O over 30 min followed by 95% MeCN for 15 min and a flow rate of 0.1 mL/min. The assay was repeated using the standalone SAT C155S, C155A and H277A mutants.

**Crystallization of the SAT Domain.** To obtain selenomethionine-substituted SAT, a single BL21(DE3) colony transformed with pJW49 was used to inoculate 5 mL of LB media supplemented with kanamycin and grown for nine h with constant shaking at 37 °C. A 2 mL aliquot was used to inoculate 50 mL of M9 minimal media in a 250 mL flask. The 50 mL culture was supplemented with kanamycin and shaken overnight at 37 °C. A 2 L shake flask containing 500 mL of M9 minimal media was also shaken overnight at 37 °C to prewarm and aerate the media for expression. After overnight incubation, 5 mL of the 50 mL culture was used to inoculate 500 mL of prewarmed M9 minimal media supplemented with kanamycin. The cells were shaken at 250 rpm at 37 °C to an OD<sub>600</sub> of 0.5. To the 500 mL culture, 100 mg of Lys, 100 mg of Phe, 100 mg of Thr, 50 mg of Ile, 50 mg of Leu, 50 mg of Val and 60 mg of selenomethionine (Sigma-Aldrich) was added. The culture was shaken at 250 rpm for 15 min at 37 °C and then induced with 0.5 mM IPTG. After induction, the culture was shaken at 250 rpm at 16 °C for 19 h. The cells were harvested as described above and the lysate was incubated with 1 mL Ni-NTA resin for 4.5 h at 4 °C with gentle agitation. Soluble selenomethionine-enriched SAT was purified by gravity-flow column chromatography using increasing concentrations of imidazole in buffer A (50 mM Tris-HCl, 500 mM NaCl, 20 mM–250 mM imidazole, pH 7.9).

Thrombin cleavage was used to remove the N-terminal His<sub>8</sub>-tag from the selenomethionine-labeled protein. To a three inch piece of

6–8 K molecular weight cutoff Spectra/Por dialysis membrane (Spectrum Laboratories), 72 mg of protein was incubated with 144  $\mu\text{L}$  of thrombin (0.5 mg/mL, Sigma-Aldrich) and incubated overnight at 4 °C in 2 L of dialysis buffer (50 mM Tris-HCl, 500 mM NaCl, 2 mM DTT, pH 7.9). Thrombin was removed by passing the protein mixture through benzamidine sepharose 4 fast-flow (GE Healthcare). The thrombin-free extract was then passed through Ni-resin to remove any remaining His-tagged SAT. The resulting flow-through was concentrated to 8.5 mg/mL using an Ultracel 30 000 MWCO centrifugal filter (Amicon Inc.) and 100  $\mu\text{L}$  aliquots were dispensed and stored at –80 °C. Purity of the selenomethionine-enriched SAT and efficiency of the cleavage was evaluated by SDS-PAGE (Figure S4).

Crystals for Se-methionine-enriched *apo* SAT and hexanoyl-bound SAT crystals were obtained using the hanging-drop vapor-diffusion crystallization method. Mixing 1  $\mu\text{L}$  of 8.5 mg/mL Se-methionine-enriched protein with 1  $\mu\text{L}$  crystallization buffer (0.1 M Tris pH 7.0 and 20% PEG 8000) and diffusing against a reservoir containing 500  $\mu\text{L}$  of 0.1 M Tris pH 7.0 and 20% PEG 8000 produced rectangular plates after five d at 4 °C. Initial hits were obtained at the UCLA Crystallization Facility using drops dispensed by a Mosquito liquid handling device (TTP Lab Tech). The rectangular plate crystals belong to the orthorhombic space group  $P2_12_12_1$  with cell dimensions of  $a = 43.13$ ,  $b = 52.16$  and  $c = 163.04$  Å. To obtain hexanoyl-bound SAT, hexanoyl-CoA was added to a final concentration of 1.4 mM to a 1  $\mu\text{L}$ :1  $\mu\text{L}$  mixture of Se-methionine-enriched SAT protein and reservoir solution (0.1 M Tris pH 7.5 and 20% PEG 8000). After seven d at 4 °C, rectangular plate crystals were observed. The crystals belong to the orthorhombic space group  $P2_12_12_1$  with cell dimension of  $a = 45.67$ ,  $b = 52.36$  and  $c = 168.03$  Å.

**Data Collection for the SAT Domain.** All data was collected at 100 K at APS-NECAT beamline 24-ID-C on a DECTRIS-PILATUS 6 M pixel detector. Single crystals were mounted on CrystalCap HT Cryoloops (Hampton Research, Aliso Viejo, CA). Se-methionine-enriched SAT crystals were cryo-protected in mother liquor solution containing 33% glycerol and flash frozen in liquid nitrogen prior to data collection.

**Data Processing, Phasing and Refinement of the SAT Domain.** Unit cell dimensions, phasing and refinement statistics are presented in Table S5. Data sets of the Se-methionine Apo-SAT and hexanoyl-SAT crystals were collected at the Advanced Photon Source (APS) at the beamline 24-ID-C (NE-CAT). For the Se-methionine crystal a Single Anomalous Dispersion (SAD) data set was collected at 0.9792 Å. Data was indexed, integrated, and scaled using XDS/XSCALE.<sup>41</sup>

Phases for the Se-methionine *apo* SAT protein were obtained using the program HKL2MAP,<sup>42</sup> which is the graphic interface for the SHELXC/D/E programs. Seven selenium sites were found by SHELXD.<sup>43</sup> SHELXE<sup>44</sup> assigned the hand, produced the first set of phases and performed solvent flattening. The final Figure of Merit was 0.817. The new set of phases were fed into the program BUCCANEER,<sup>45</sup> which traced 95% of the protein residues. The Se-methionine *apo* SAT model was refined to 1.6 Å. Final iterative rounds of model building and refinement were carried out using Coot<sup>46</sup> and PHENIX<sup>47</sup> with TLS refinement.<sup>48</sup> More than 99.2% of the residues in the model were found in the favored regions of the Ramachandran plot by MOLPROBITY.<sup>33</sup> There is one molecule of *apo* SAT in the asymmetric unit. The *apo* SAT model has clear electron density for all residues, except for C-terminal residues 397 to 414. Additionally, some side chains with no interpretable electron density were omitted in the final model.

The hexanoyl-bound SAT structure was solved starting with the *apo* SAT model using rigid body refinement followed by 12 cycles of manual model building and refinement using PHENIX.<sup>47</sup> More than 98.7% of the residues in the model were found in the favored regions of the Ramachandran plot by MOLPROBITY.<sup>33</sup> There is one molecule of hexanoyl-bound SAT in the asymmetric unit. The hexanoyl-bound SAT model has clear electron density for all residues, except for C-terminal residues 396 to 414.

## ■ ASSOCIATED CONTENT

### ■ Supporting Information

Spectroscopic information and additional experimental details. The atomic coordinates and structure factors for *apo*SAT and hexanoyl-bound SAT have been deposited in the Protein Data Bank under accession codes 4RO5 and 4RPM, respectively. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.5b04520.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*yitang@ucla.edu

### Present Address

#Department of Medicinal Chemistry, University of Utah, Salt Lake City, Utah 84112, United States.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was supported by the US NIH (1R01GM085128 and 1DP1GM106413) to Y.T.; NSERC to J.C.V.; and JSPS (No. LS103) to K.W. We thank Mike Collazo at the UCLA-DOE X-ray Crystallization and Crystallography Core Facilities, which are supported by DOE Grant DE-FC02-02ER63421, and M. Cappel, K. Rajashankar, N. Sukumar, J. Schuermann, I. Kourinov and F. Murphy at NECAT beamlines 24-ID at APS, which are supported by grants from the NIH (5P41RR015301-10 and 8 P41 GM103403-10). Use of the APS is supported by the DOE under Contract DE-AC02-06CH11357. J.M.W. thanks L'Oréal USA for Women in Science for a postdoctoral fellowship. NMR instrumentation was supported by the NSF Equipment Grant CHE-1048804.

## ■ REFERENCES

- (1) Cane, D. E.; Walsh, C. T.; Khosla, C. *Science* **1998**, *282*, 63.
- (2) Cox, R. J. *Org. Biomol. Chem.* **2007**, *5*, 2010.
- (3) Ma, S. M.; Zhan, J.; Watanabe, K.; Xie, X.; Zhang, W.; Wang, C.; Tang, Y. *J. Am. Chem. Soc.* **2007**, *129*, 10642.
- (4) (a) Chooi, Y. H.; Tang, Y. *J. Org. Chem.* **2012**, *77*, 9933. (b) Kennedy, J.; Auclair, K.; Kendrew, S. G.; Park, C.; Vederas, J. C.; Hutchinson, C. R. *Science* **1999**, *284*, 1368.
- (5) (a) Crawford, J. M.; Dancy, B. C. R.; Hill, E. A.; Udway, D. W.; Townsend, C. A. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 16728. (b) Gao, Z.; Wang, J.; Norquay, A. K.; Qiao, K.; Tang, Y.; Vederas, J. C. *J. Am. Chem. Soc.* **2013**, *135*, 1735.
- (6) Yu, J.; Chang, P. K.; Ehrlich, K. C.; Cary, J. W.; Bhatnager, D.; Cleveland, T. E.; Payne, G. A.; Linz, J. E.; Woloshuk, C. P.; Bennett, J. W. *Appl. Environ. Microbiol.* **2004**, *70*, 1253.
- (7) Zhou, H.; Qiao, K.; Gao, Z.; Meehan, M. J.; Li, J.; Dorrestein, P. C.; Vederas, J. C.; Tang, Y. *J. Am. Chem. Soc.* **2010**, *132*, 4530.
- (8) Zhou, H.; Qiao, K.; Gao, Z.; Vederas, J. C.; Tang, Y. *J. Biol. Chem.* **2010**, *285*, 41412.
- (9) Zhou, H.; Zhan, J.; Watanabe, K.; Xie, X.; Tang, Y. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 6249.
- (10) Chiang, Y. M.; Szweczyk, E.; Davidson, A. D.; Keller, N.; Oakley, B. R.; Wang, C. C. C. *J. Am. Chem. Soc.* **2009**, *131*, 2965.
- (11) Winter, J. M.; Sato, M.; Sugimoto, S.; Chiou, G.; Garg, N. K.; Tang, Y.; Watanabe, K. *J. Am. Chem. Soc.* **2012**, *134*, 17900.
- (12) Liu, T.; Chiang, Y. M.; Somoza, A. D.; Oakley, B. R.; Wang, C. C. C. *J. Am. Chem. Soc.* **2011**, *133*, 13314.
- (13) Liu, T.; Sanchez, J. F.; Chiang, Y.-M.; Oakley, B. R.; Wang, C. C. C. *Org. Lett.* **2014**, *16*, 1676.
- (14) Huitt-Roehl, C. R.; Hill, E. A.; Adams, M. M.; Vagstad, A. L.; Li, J. W.; Townsend, C. A. *ACS Chem. Biol.* **2015**, *10*, 1443.



- (15) (a) Antosch, J.; Schaefer, F.; Gulder, T. A. M. *Angew. Chem., Int. Ed.* **2014**, *53*, 3011. (b) Zhang, G.; Zhang, W.; Zhang, Q.; Shi, T.; Ma, L.; Zhu, Y.; Li, S.; Zhang, H.; Zhao, Y. L.; Shi, R.; Zhang, C. *Angew. Chem., Int. Ed.* **2014**, *53*, 4840. (c) Li, Y.; Chen, H.; Ding, Y.; Xie, Y.; Wang, H.; Cerny, R. L.; Shen, Y.; Du, L. *Angew. Chem., Int. Ed.* **2014**, *53*, 7524.
- (16) Lee, K. K.; Da Silva, N. A.; Kealey, J. T. *Anal. Biochem.* **2009**, *394*, 75.
- (17) Keszenman-Pereyra, D.; Lawrence, S.; Twfieg, M. E.; Price, J.; Turner, G. *Curr. Genet.* **2003**, *43*, 186.
- (18) Zhou, H.; Gao, Z. Z.; Qiao, K. J.; Wang, J. J.; Vederas, J. C.; Tang, Y. *Nat. Chem. Biol.* **2012**, *8*, 331.
- (19) Wang, F.; Wang, Y.; Ji, J.; Zhou, Z.; Yu, J.; Zhu, H.; Su, Z.; Zhang, L.; Zheng, J. *ACS Chem. Biol.* **2015**, *10*, 1017.
- (20) Crawford, J. M.; Vagstad, A. L.; Ehrlich, K. C.; Townsend, C. A. *Bioorg. Chem.* **2008**, *36*, 16.
- (21) Crawford, J. M.; Vagstad, A. L.; Whitworth, K. P.; Ehrlich, K. C.; Townsend, C. A. *ChemBioChem* **2008**, *9*, 1019.
- (22) (a) Keatinge-Clay, A. T.; Shelat, A. A.; Savage, D. F.; Tsai, S.-C.; Miercke, L. J. W.; O'Connell, J. D., III; Khosla, C.; Stroud, R. M. *Structure* **2003**, *11*, 147. (b) Arthur, C. J.; Williams, C.; Pottage, K.; Ptoskon, E.; Findlow, S. C.; Burston, S. G.; Simpson, T. J.; Crump, M. P.; Crosby, J. *ACS Chem. Biol.* **2009**, *4*, 625.
- (23) Zhang, L.; Liu, W.; Xiao, J.; Hu, T.; Chen, J.; Chen, K.; Jiang, H.; Shen, X. *Protein Sci.* **2007**, *16*, 1184.
- (24) Natarajan, S.; Kim, J.-K.; Jung, T.-K.; Doan, T. T. N.; Ngo, H.-P.-T.; Hong, M.-K.; Kim, S.; Tan, V. P.; Ahn, S. J.; Lee, S. H.; Han, Y.; Ahn, Y.-J.; Kang, L.-W. *Mol. Cells* **2012**, *33*, 19.
- (25) Wong, F. T.; Jin, X.; Mathews, I. I.; Cane, D. E.; Khosla, C. *Biochemistry* **2011**, *50*, 6539.
- (26) Tang, Y.; Chen, A. Y.; Kim, C.-Y.; Cane, D. E.; Khosla, C. *Chem. Biol.* **2007**, *14*, 931.
- (27) Tang, Y.; Kim, C.-Y.; Mathews, I. I.; Cane, D. E.; Khosla, C. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 11124.
- (28) Bergeret, F.; Gavalda, S.; Chalut, C.; Malaga, W.; Quémard, A.; Pedelacq, J.-D.; Daffé, M.; Guilhot, C.; Mourey, L.; Bon, C. *J. Biol. Chem.* **2012**, *287*, 33675.
- (29) Park, H.; Keavany, B. M.; Dyer, D. H.; Thomas, M. G.; Forest, K. T. *PLoS One* **2014**, *9*, e110965.
- (30) Liew, C. W.; Nilsson, M.; Chen, M. W.; Sun, H.; Cornvik, T.; Liang, Z.-X.; Lescar, J. *J. Biol. Chem.* **2012**, *287*, 23203.
- (31) Dundas, J.; Ouyang, Z.; Tseng, J.; Binkowski, A.; Turpaz, Y.; Lliang, J. *Nucleic Acids Res.* **2006**, *34*, W116.
- (32) DeLano, W. L. *The PyMOL User's Manual*; DeLano Scientific: San Carlos, CA, 2002.
- (33) Chen, V. B. W.; Arendall, B.; Headd, J. F.; Keedy, D. A.; Immormino, R. M.; Kapral, G. J.; Murray, L. W.; Richardson, J. S.; Richardson, D. C. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2010**, *D66*, 12.
- (34) (a) Yang, J.; Yan, R.; Roy, A.; Xu, D.; Poisson, J.; Zhang, Y. *Nat. Methods* **2015**, *12*, 7. (b) Roy, A.; Kucukural, A.; Zhang, Y. *Nat. Protoc.* **2010**, *5*, 725. (c) Zhang, Y. *BMC Bioinf.* **2008**, *9*, 40.
- (35) Schneidman-Duhovny, D.; Inbar, Y.; Nussinov, R.; Wolfson, H. *J. Nucleic Acids Res.* **2005**, *33*, W363.
- (36) Alekseyev, V. Y.; Liu, C. W.; Cane, D. E.; Puglisi, J. D.; Khosla, C. *Protein Sci.* **2007**, *16*, 2093.
- (37) Wong, F. T.; Chen, A. Y.; Cane, D. E.; Khosla, C. *Biochemistry* **2010**, *49*, 95.
- (38) Ye, Z.; Musiol, E. M.; Weber, T.; Williams, G. J. *Chem. Biol.* **2014**, *21*, 636.
- (39) Nguyen, C.; Haushalter, R. W.; Lee, D. J.; Markwick, P. R.; Bruegger, J.; Caldara-Festin, G.; Finzel, K.; Jackson, D. R.; Ishikawa, F.; O'Dowd, B.; McGammon, A.; Opella, S. J.; Tsai, S.-C.; Burkart, M. D. *Nature* **2013**, *505*, 427.
- (40) Jez, J. M.; Ferrer, J. L.; Bowman, M. E.; Dixon, R. A.; Noel, J. P. *Biochemistry* **2000**, *39*, 890.
- (41) Kabsch, W. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2010**, *66*, 125.
- (42) Pape, T.; Schneider, T. R. *J. Appl. Crystallogr.* **2004**, *37*, 844.
- (43) Uson, I.; Sheldrick, G. *Curr. Opin. Struct. Biol.* **1999**, *9*, 643.
- (44) Sheldrick, G. M. *Z. Kristallogr.* **2002**, *217*, 644.
- (45) Winn, M. D.; Ballard, C. C.; Cowtan, K. D.; Dodson, E. J.; Emsley, P.; Evans, P. R.; Keegan, R. M.; Krissinel, E. B.; Leslie, A. G. W.; McCoy, A.; McNicholas, S. J.; Murshudov, G. N.; Pannu, N. S.; Potterton, E. A.; Powell, H. R.; Read, R. J.; Vagin, A.; Wilson, K. S. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2011**, *67*, 235.
- (46) Emsley, P.; Cowtan, K. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2004**, *60*, 2126.
- (47) Adams, P. D.; Grosse-Kunstleve, R. W.; Hung, L. W.; Loerger, T. R.; McCoy, A. J.; Moriarty, N. W.; Read, R. J.; Sacchettini, J. C.; Sauter, N. K.; Terwilliger, T. C. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2002**, *58*, 1948.
- (48) Painter, J.; Merritt, E. A. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2006**, *62*, 439.