


A Photostable Silicon Rhodamine Platform for Optical Voltage Sensing

Yi-Lin Huang,^{†,||} Alison S. Walker,^{†,||} and Evan W. Miller^{*,†,‡,§}

[†]Departments of Chemistry, [‡]Molecular & Cell Biology, and [§]Helen Wills Neuroscience Institute, University of California, Berkeley, California 94720, United States

S Supporting Information

ABSTRACT: This paper describes the design and synthesis of a photostable, far-red to near-infrared (NIR) platform for optical voltage sensing. We developed a new, sulfonated silicon rhodamine fluorophore and integrated it with a phenyl-enevinylene molecular wire to create a Berkeley Red Sensor of Transmembrane potential, or BeRST 1 (“burst”). BeRST 1 is the first member of a class of far-red to NIR voltage sensitive dyes that make use of a photoinduced electron transfer (PeT) trigger for optical interrogation of membrane voltage. We show that BeRST 1 displays bright, membrane-localized fluorescence in living cells, high photostability, and excellent voltage sensitivity in neurons. Depolarization of the plasma membrane results in rapid fluorescence increases (24% $\Delta F/F$ per 100 mV). BeRST 1 can be used in conjunction with fluorescent stains for organelles, Ca^{2+} indicators, and voltage-sensitive fluorescent proteins. In addition, the red-shifted spectral profile of BeRST 1, relative to commonly employed optogenetic actuators like ChannelRhodopsin2 (ChR2), which require blue light, enables optical electrophysiology in neurons. The high speed, sensitivity, photostability and long-wavelength fluorescence profiles of BeRST 1 make it a useful platform for the noninvasive, optical dissection of neuronal activity.



INTRODUCTION

Rapid changes in the membrane potential (V_{mem}) of excitable cells like neurons and cardiomyocytes play a central role in defining cellular signaling and physiological profiles of these specialized cells. Classically, V_{mem} is monitored and measured via patch clamp electrophysiology, in which a microelectrode attached to the cell of interest enables highly sensitive recordings of membrane voltage with exquisite temporal resolution.¹ However, the use of electrodes is highly invasive, limits records to the soma of a single cell (space clamp error),² and is extremely low throughput. Optical techniques for recording voltage represent a feasible solution to these problems, because they are minimally invasive, requiring delivery of a sensor and photons, and can be high throughput.^{3,4} Ca^{2+} imaging has long been used as a surrogate for direct optical measurement of voltage, in part because the robust imaging agents, whether based on small molecules^{5–7} or on fluorescent proteins,⁸ are sensitive, applicable to a wide range of biological contexts, and come in a variety of colors.⁹ The use of Ca^{2+} imaging provides only an imperfect approximation of V_{mem} changes, since intracellular $[\text{Ca}^{2+}]_i$ ($[\text{Ca}^{2+}]_i$) rises are triggered by the depolarization of V_{mem} .¹⁰ Additionally, because $[\text{Ca}^{2+}]_i$ transients last for hundreds of milliseconds and because the Ca^{2+} sensors themselves can buffer the rise and fall of $[\text{Ca}^{2+}]_i$, resolving fast spiking neuronal

events becomes impossible or difficult, requiring extensive deconvolution protocols.^{11,12}


Direct voltage imaging, therefore, is attractive, because it can provide a direct readout of V_{mem} while still achieving the spatial resolution, high throughput and minimal invasiveness of Ca^{2+} imaging.^{3,4} Recently, there has been renewed interest in the development of fluorescent voltage sensors, through the use of small molecules,^{13–17} fluorescent proteins,^{18,19} combinations of the two,^{20,21} or opsins.²²

Our lab is developing VoltageFluors,²³ or VF dyes, a small molecule platform for voltage sensing that relies on photo-induced electron transfer (PeT)²⁴ modulated by changes in V_{mem} to achieve fast, sensitive, nondisruptive voltage sensing. While the original VF dyes^{13,23} have already proved useful in cultured neurons,^{13,23} brain slices,²³ ex vivo leech nervous system²⁵ and mouse pancreatic models,²⁶ a significant limitation is an excitation and emission profile that lies in the blue/green region of the electromagnetic spectrum. Because VF dyes are excited in the 480 to 515 nm range and emit around 530 nm, this significantly overlaps with a number of useful optical tools, such as GFP, robust Ca^{2+} sensors like Oregon Green BAPTA²⁷ and the GCaMP family,^{8,28,29} and optogenetic tools like ChannelRhodopsin2 (ChR2).^{30,31} Additionally, for

Received: June 26, 2015

Published: August 3, 2015

Scheme 1. Synthesis of Berkeley Red Sensor of Transmembrane Potential 1 (BeRST 1)



imaging in thicker biological samples like tissue slices or whole animals, excitation and emission profiles (650 to 900 nm) are desirable because lower energy photons reduce tissue damage, scatter less, are absorbed by fewer endogenous chromophores, and produce lower levels of autofluorescence.³² Voltage sensitive dyes that possess excitation and emission profiles in the near-infrared (NIR) window would enable integration with current optical tools and avoid complications associated with imaging at shorter wavelengths.

RESULTS AND DISCUSSION

Design and Synthesis of Sulfonated Silicon Rhodamine. Here, we report the design and development of Berkeley Red Sensor of Transmembrane potential 1 (BeRST 1, “burst”), a new voltage-sensitive sulfonated silicon rhodamine (Si-rhodamine) fluorophore. Si-rhodamines have gained attention recently because they display photostable, far-red to NIR fluorescence that can be readily modulated by PeT.³³ BeRST 1 features a Si-rhodamine^{33,34} core bearing a sulfonic acid on the *meso* aryl ring. This sulfonated Si-rhodamine, which we term Berkeley Red, or BR (Scheme 1), provides far-red to NIR excitation and emission profiles on account of the Si-substituted xanthene, and improved retention of the dye at the extracellular surface of cell membranes, due to the anionic sulfonate (see below).^{13,23}

BR was synthesized in three steps from intermediates **1** and **3**, which are available in 1 and 4 steps, respectively, from reported procedures.^{35,36} The key step in the synthesis of BR core with attendant sulfonate is the nucleophilic addition of an organolithium intermediate, derived from lithium-halogen exchange on **1**, to xanthone **3** that had been activated with

triflic anhydride. Deprotection of the isopropyl ester with BBr_3 provided BR in 9%, over three steps (Scheme 1).

BR displays an absorbance profile centered at 658 nm and an extinction coefficient of $150\,000\text{ M}^{-1}\text{ cm}^{-1}$ in aqueous buffer (Figure 1a, 50 mM TBS, pH 7.5, 0.1% SDS). BR features far-red to NIR fluorescence, emitting strongly at 681 nm ($\Phi_{\text{fl}} = 0.24$ in TBS/SDS buffer and 0.32 in methanol). A key requirement for the use of BR in voltage sensing applications is that BR not cyclize to the nonfluorescent state in low dielectric environments like the cell membrane. Although this spirocyclization is a useful property for superresolution microscopy,^{37,38} cyclization leads to a more hydrophobic conformation and encourages the passage of the voltage sensor through the plasma membrane. Staining of internal membranes and cytosol then sharply increases background fluorescence and limits the useful signal one can extract from an imaging experiment. As predicted, placement of a sulfonate group on the *meso* aryl ring prevented spirocyclization, with no change in absorbance observed in either acidic or basic environments (pH 2.5 to 9) (Supporting Information (SI) Figure S1c and e). BR absorbance remains unchanged (SI Figure S1b and d) through a range of solvent dielectric constants from $\epsilon = 6$ (10% water/dioxane v/v) to $\epsilon = 71$ (90% water/dioxane v/v),³⁹ confirming that BR conformation displays minimal dependence on solvent polarity. The environmental insensitivity of BR stands in stark contrast to carboxy Si-Rhodamine derivatives, which display massive absorbance fluctuations upon changes in the dielectric constant of the solvent.³⁷ The spirocyclization is effectively minimized by replacement of carboxyl with sulfonate. We further confirmed that BR’s sulfonic acid renders it cell-impermeant, as HEK cells bathed in BR showed negligible cytosolic fluorescence. (SI Figure S2)




Figure 1. In vitro and cellular characterization of BeRST 1. Absorbance (blue) and emission (red) spectra of (a) BeRST 1 in aqueous buffer (50 mM TBS, pH 7.5, 0.1% SDS). Excitation was provided at 635 nm. (b) Photostability of BeRST 1 and VF2.1.Cl dyes in HEK cells. Cells were loaded with 200 nM BeRST 1 or VF2.1.Cl and then illuminated continuously for 10 min at either 631 nm (BeRST 1) or 475 nm (VF2.1.Cl) at 162 W/cm². Images were acquired at 20 s intervals. The normalized fluorescence intensity from dye-loaded cells was plotted vs time. Magenta circles represent BeRST 1-stained cells and green circles represent VF2.1.Cl-stained cells. Error bars are standard error of the mean for $n = 6$ separate experiments. Multicolor epifluorescence imaging with BeRST 1. HEK cells were loaded with 1 μ M BeRST 1, 1 μ M Hoechst 33342, and 5 μ M Rhodamine 123. (c) BeRST 1 fluorescence localized to the cell membrane, (d) Rhodamine 123 fluorescence localized to mitochondria, (e) Hoechst 33342 fluorescence localized to the nucleus, and (f) transmitted light DIC of HEK cells. Scale bar is 10 μ m.

Design and Synthesis of BeRST 1. Having established that sulfonated Si-rhodamine BR possesses bright, long-wavelength fluorescence and absorbance that does not vary with pH or solvent dielectric, we further elaborated BR with a phenylenevinylene molecular wire to generate BeRST 1. The key brominated Berkeley Red (bromo-BR, Scheme 1) was synthesized in 15% yield via chelation-assisted lithiation of 2. Lithiation was accomplished in CH₂Cl₂ at -20 °C, with triflic anhydride used to activate 3 for addition of the organolithium species. A variety of other conditions involving different combinations of solvents (THF, CH₂Cl₂), reagents (*n*-BuLi, *i*PrMgCl-LiCl⁴⁰), additives (MgSO₄), and temperatures (-78 °C, -40 and 22 °C) all gave no reaction, insufficient yield, or complicated mixtures.

Regioselective lithiation under the optimized reaction conditions was confirmed by quenching the generated lithium species with acid to give a product consistent by ¹H NMR with

ortho-lithiation (SI Spectrum 10).⁴¹ Pd-catalyzed Heck coupling of bromo-BR with methoxy-substituted phenylenevinylene dimethyl aniline molecular wire 4—available in 3 steps according to established procedures²³—provided BeRST 1 in 62% yield, after purification on silica gel. BeRST 1 exhibits spectral properties similar to the parent BR dye, with excitation and emission centered at 658 and 683 nm (Figure 1a compare to SI Figure S1a). Compared to BR, the quantum yield of BeRST 1 is substantially reduced, at 0.017 and 0.022 in aqueous buffer and MeOH, respectively.

Cellular Characterization of BeRST 1. BeRST 1 localizes to cell membranes when bath applied to HEK cells (Figure 1c). Importantly, BeRST 1 fluorescence remained localized to the cell membrane throughout the course of imaging experiments. This is in contrast to the BR fluorophore in isolation, which displays negligible cellular fluorescence after loading under identical conditions—further establishing the ability of the sulfonate in BeRST 1 and BR to prevent dye internalization (SI Figure S2a vs c).

BeRST 1 shows excellent photostability (Figure 1b), with a bleaching half-life of approximately 5 min under intense illumination conditions ($I = 162$ W/cm², 631 nm LED). Which is approximately an order of magnitude higher than intensities used for BeRST 1 imaging in concert with other optical tools (see Figures 4, 7 and 8). By comparison, VF2.1.Cl, which has a similar voltage sensitivity, 27% $\Delta F/F$ per 100 mV, has a bleaching half-life of <60 s under identical illumination intensities ($I = 162$ W/cm², 475 nm LED). Since BeRST 1 has a larger extinction coefficient than VF2.1.Cl at the incident irradiation (150 000 vs 98 000), this comparison slightly underestimates the photostability of BeRST 1 (Si-rhodamine) relative to VF2.1.Cl (fluorescein).

To determine the voltage sensitivity of BeRST 1, we performed whole-cell patch clamp electrophysiology on HEK cells loaded with 1 μ M BeRST 1 (Figure 2a and b). Depolarization of voltage-clamped HEK cells with BeRST 1 results in prompt increases in fluorescence, while hyperpolarization gives fluorescence decreases (Figure 2b, SI Movie S1). BeRST 1 has a voltage sensitivity of approximately 24% \pm 5% $\Delta F/F$ per 100 mV—comparable to first generation blue light-excitable VoltageFluors—and is linear over a physiologically relevant range spanning ± 100 mV (Figure 2b).

Voltage Imaging with BeRST 1 in Neurons. BeRST 1 stained cell membranes of cultured hippocampal neurons equally well (Figure 3a and b). Whole-cell patch-clamp electrophysiology on cultured neurons loaded with BeRST 1 established that BeRST 1 can detect action potentials (APs) in single trials. Action potentials were evoked under current clamp mode, and the resulting optical response matched the recorded V_{mem} changes exactly (Figure 3d). Optically recorded action potentials (1.8 kHz optical sampling) were indistinguishable from a simultaneously recorded traditional electrophysiological trace. When measured electrophysiologically, the fwhm duration of action potentials were 1.65 ± 0.15 ms, while optical action potential records gave values of 1.97 ± 0.14 ms ($n = 7$ cells each condition). The difference of 0.32 ms is far less than our optical sampling interval of 0.56 ms, and therefore within the error of measurement for our current imaging apparatus. Optical responses from BeRST 1 for action potentials under these conditions were approximately 9.5% $\Delta F/F$ ($\pm 1.2\%$, $n = 7$ cells).

Using whole-cell patch clamp electrophysiology without imaging, we established that the presence of BeRST 1 has no




Figure 2. Voltage sensitivity of BeRST 1 in HEK cells. (a) Plot of fractional change in fluorescence ($\Delta F/F$) vs time from BeRST 1-stained HEK cells under whole-cell, voltage-clamp conditions. Cells were held at -60 mV and stepped to hyper- or depolarizing potentials (± 100 mV) in 20 mV increments. (b) Plot of $\Delta F/F$ vs final membrane potential. Error bars are \pm standard error of the mean for $n = 15$ cells from 3 different cultures.




Figure 3. Action potential (AP) visualization with BeRST 1. (a) Rat hippocampal neurons were (b) stained with $1 \mu\text{M}$ BeRST 1 and field stimulated to evoke action potentials that were (c) recorded optically. Optical sampling frequency is 500 Hz, acquired with an EMCCD camera under $63\times$ magnification. Scale bar is $10 \mu\text{m}$. In a separate experiment, neurons were subjected to whole-cell current clamp and stimulated to induce action potential firing. Dual electrical and optical recording (1.8 kHz frame rate) are shown in panel (d). Black trace is the electrophysiological recording and red circles represent the optical response. A small stimulus artifact is apparent before and after the action potential in the black trace.

discernible effect on the ability of neurons to fire action potentials or on the measured properties of the action potentials. We found no difference in peak action potential

amplitude, duration, rise time, decay time or cellular capacitance when comparing the electrophysiological response to stimulation in neurons with or without $1 \mu\text{M}$ BeRST 1 ($n = 12$ and 11 dye-free and dye-loaded neurons, SI Figure S3). Action potentials were readily detectable under field stimulation conditions as well, giving $17.8\% \Delta F/F$ ($\pm 1.6\%$, $n = 5$ cells) (Figure 3c). In most cases, a repolarization is visible following the action potential, further highlighting the utility of BeRST 1 to detect and report on full action potential waveforms.

Multicolor Imaging with BeRST 1. One of the benefits of voltage sensors in the far-red to NIR window is that these new tools can be interfaced with a wide variety of blue, green, and red optical tools. Of particular interest are the genetically encoded optical tools based on GFP, since on the whole, these fluorescent proteins are more widely used and exhibit more predictable cellular and photophysical behavior than their red fluorescent counterparts.^{42,43} We first established that BeRST 1 can perform multicolor imaging with some commonly used molecular probes for cellular organelles: the nucleus and mitochondria. We performed simultaneous three-color live cell imaging with BeRST 1 (Figure 1c), rhodamine 123 mitochondrial stain (Figure 1d),⁴⁴ and Hoescht 33342 nuclear stain (Figure 1e).⁴⁵ Under these conditions distinct cellular components were clearly visible in 3 colors, establishing the utility of BeRST 1 for multicolor imaging. (Figure 1c–f)

Functional information is often required from a subset of cells, which are typically identified by the targeted expression of a fluorescent protein. We next sought to establish the ability of BeRST 1 to provide multicolor functional imaging, by pairing BeRST 1 staining with GFP labeling. Neurons expressing cytosolic GFP were loaded with BeRST 1 and then action potentials were evoked via field stimulation (SI Figure S4). Under these conditions, BeRST 1 gave large changes in fluorescence in cells not expressing GFP, approximately $9.2\% \Delta F/F$ per action potential ($\pm 1.6\%$), with a SNR of 62:1. GFP-expressing cells in the same field of view gave virtually identical responses ($8.7 \pm 0.8\% \Delta F/F$, SNR = 63:1, $n = 5$ pairs of cells) establishing that the response of BeRST 1 is not diminished by the presence of GFP, nor by optical bleedthrough or cross-excitation of GFP (SI Figure S4). We envision this methodology could be used to label cells or subcellular locales of interest with an optically orthogonal fluorescent protein (GFP) and then record from regions of interest (ROIs) defined by the fluorescent protein. Although extracellular field stimulation confirmed that the presence of GFP does not attenuate the BeRST 1 optical response, it did not allow us to interrogate independent signals coming from GFP-positive and -negative cells, since all neurons respond to extracellular field stimulation.

To demonstrate that we could spatially segregate individual neuronal activity in cultures expressing GFP, we imaged spontaneous activity in the same cultures expressing GFP and labeled with BeRST 1. Using GFP to define regions of interest, we could clearly distinguish the activity of the cell of interest from the surrounding cells. In Figure 4, the ROIs defined by the GFP-positive neurons (#3 and #6) clearly display activity strikingly different from the other four cells in the field of view. Taken together, these experiments demonstrate the ability of BeRST 1 to provide multisite optical recording of neuronal activity in parallel with a GFP counterstain.

We next determined the ability of BeRST 1 to participate in two-color, multifunctional imaging, that is, combining voltage imaging with either Ca^{2+} imaging, voltage imaging, or optogenetic manipulation via ChR2. The GCaMP^{28,29} family




Figure 4. Spontaneous voltage imaging in GFP-labeled cells with BeRST 1. Epifluorescence images of rat hippocampal neurons expressing (a) GFP and (b) stained with BeRST 1. Scale bar is 20 μm . (c) Optical traces of spontaneous activity in neurons from panels (a) and (b). Numbers next to traces correspond to indicated cells in panel (b). Optical sampling rate is 500 Hz.

of Ca^{2+} sensors are one of the most widely used functional probes for systems and cellular neuroscience. However, because the excitation and emission spectrum of GCaMP lies squarely within the regime of previous generations of VoltageFluors^{13,23} (as well as other functional probes), this has precluded the use of GCaMPs with PeT-based voltage sensitive dyes. Neurons expressing the genetically encoded Ca^{2+} sensor, GCaMP6s,²⁹ which provides the largest fluorescence response to Ca^{2+} , were stained with BeRST 1 (Figure 5a and b). GCaMP6s-positive neurons (Figure 5b) displayed bright green cytosolic fluorescence, while membranes were clearly labeled with BeRST 1 (Figure 5a). We evoked action potentials via field stimulation and obtained optical recordings, first in the green channel for GCaMP6s and then in the far-red to NIR channel for BeRST 1, from a single cell expressing GCaMP6s and stained with BeRST 1 (Figure 5a and b). In sequential optical traces, both the rise in V_{mem} and rise in $[\text{Ca}^{2+}]_i$ are clearly visible (Figure 5c). Dual BeRST 1 and GCaMP6s imaging enables discrimination of voltage and Ca^{2+} transients, with the voltage transient (approximately 15% $\Delta F/F$) preceding the measured GCaMP6s signal (approximately 5% $\Delta F/F$) for a single evoked action potential (Figure 5c). One of the advantages of voltage imaging is the ability to deconvolve fast spiking that Ca^{2+} imaging, with its slower intrinsic signals and slow probe unbinding kinetics, often cannot resolve. When neurons expressing both GCaMP6s and stained with BeRST 1 were stimulated to evoke trains of action potentials at 5, 10, and 20 Hz, discrete V_{mem} responses corresponding to individually evoked action potentials were visible in the BeRST 1 channel at all frequencies (Figure 5d), while Ca^{2+} response as measured by GCaMP6s were only discernible at 5 Hz, as measured by small changes in on the rising edge of GCaMP6s fluorescence (Figure 5e). Due to the intrinsic response speed of BeRST 1, two-color functional imaging should be generalizable to many other types of Ca^{2+} indicators. These experiments highlight the ability of BeRST 1 to participate in functional imaging experiments in which multiple signals need to be interpreted in the same system.

The nonoverlapping signals emanating from BeRST 1 and GFP-based fluorophores provide a unique opportunity to perform two-color voltage imaging. We expressed the genetically encoded voltage sensor, ASAP1,¹⁹ in cultured neurons and stained these neurons, as before, with BeRST 1 (Figure 6a–d). In response to an evoked train of action potentials (Figure 6e), ASAP1 provides decreases in relative fluorescence of about $6\% \pm 1.4\%$ (S.E.M., with a signal-to-noise ratio, SNR, of 8:1, $n = 4$ cells), which is consistent with the reported value of $-4.8\% \Delta F/F$.¹⁹ BeRST 1, in the same cell,




Figure 5. Dual voltage and Ca^{2+} imaging with BeRST 1 and GCaMP6s. Epifluorescence images of a rat hippocampal neuron stained (a) with BeRST 1 and expressing (b) GCaMP6s. Scale bar is 20 μm . (c) Sequential Ca^{2+} and voltage imaging during field stimulation of the neuron in panels (a) and (b). Voltage- (upper) and Ca^{2+} - (lower) induced changes in fluorescence. (d) Voltage- (left) and Ca^{2+} - (right) induced fluorescence response to trains of action potentials at the indicated frequency. Optical sample rate is 500 Hz for BeRST 1 and 40 Hz for GCaMP6s.

under identical stimulation parameters (Figure 6e), gives a $15\% \pm 2.1\%$ increase (S.E.M., SNR of 41:1, $n = 4$ cells), highlighting the utility of BeRST 1 for measurement of fast spiking events and for two-color voltage imaging with complementary genetically encoded voltage sensors based on GFP.¹⁸

Optical Electrophysiology with BeRST 1 and Chr2.

Finally, we hoped to interface BeRST 1 with optogenetic tools like ChannelRhodopsin2 (Chr2) in order to demonstrate the




Figure 6. Two-color voltage imaging with genetically encoded voltage-sensitive fluorescent proteins and small molecule voltage-sensitive dyes. Epifluorescence images of rat hippocampal neurons (a) stained with BeRST 1 and (b) expressing the voltage-sensitive fluorescent protein ASAP1. Zoomed-in region for functional imaging of voltage via (c) BeRST 1 staining or (d) ASAP1 fluorescence. The magnified regions correspond to the white boxes in panels (a) and (b). All scale bars are 20 μm . Trains of action potentials evoked by field stimulation at 5 Hz detected by changes in (e) BeRST 1 fluorescence (upper trace) or ASAP1 fluorescence (lower trace). (f) A single action potential, magnified from panel (e). Magenta is BeRST 1 and green is ASAP1 fluorescence. The optical sampling rate was 1.25 kHz.

feasibility of using light to both actuate and record the V_{mem} of living cells in an “all-optical” electrophysiology methodology.^{4,22,46} The action spectrum of ChR2 (470 nm peak response),⁴⁷ one of the more heavily used optogenetic tools,

contains significant overlap with the first generation of VoltageFluor dyes. Although red-shifted variants of ChR have been reported, all of them contain significant response peaks in the blue region of the spectrum.^{48–51} For this reason, it is advantageous to move the voltage-imaging component into the far-red to NIR region of the spectrum to avoid cross-talk in the optical channels for activation and recording.

Before carrying out optical recording of light-induced action potentials, we investigated whether any optical cross-talk exists between BeRST 1 and ChR2. We found a small amount of cross excitation of BeRST 1. When excited at 475 nm, BeRST 1-stained cells show fluorescence that is approximately 4% the intensity achieved with excitation at 631 nm (9.7 W/cm², 475 and 631 nm, SI Figure S5a–d). Because ChR2 requires much lower light intensity relative to typical imaging light intensities for BeRST 1 (80 mW/cm², 475 nm vs 20 W/cm², 631 nm for multicellular imaging), we estimate that less than 0.1% of BeRST 1 signal comes from cross excitation of ChR2 in actual imaging experiments.

Cross excitation is also observed upon excitation of BR and BeRST 1 derivatives in vitro (SI Figure S6a and b). Excitation of BR or BeRST 1 derivatives in aqueous buffer with cyan light (475 nm) gives approximately 1% or 7% of the total emission when compared to excitation at 630 nm, respectively (SI Figure S6c). Violet excitation (390 nm) in vitro gives 16% (BR) and 35% (BeRST 1) of full emission (SI Figure S6c), which is consistent with cross excitation of BeRST 1 in cellular membranes (approximately 47% with 9.7 W/cm² 390 nm excitation) (SI Figure S5c and d). HEK cells stained with BeRST 1 are voltage sensitive when imaged under violet illumination (SI Figure S5e, 390 nm excitation, 3.3 W/cm², 680 nm emission). Studies are underway to probe the nature of this exceptionally long pseudo-Stokes shift (290 nm) which may involve energy transfer from the conjugated molecular wire.³⁶

Having established that negligible cross excitation of BeRST 1 occurs (<0.1%) under experimental conditions, we next




Figure 7. All optical electrophysiology with BeRST 1 and ChR2. Epifluorescence images of rat hippocampal neurons expressing (a) ChR2-YFP and (b) stained with BeRST 1. Inset on panel (b) is the region inscribed by the dotted white box. The inset image is single frame used to acquire data for panels (d) and (f). Scale bars are 20 μm (panel a/b) and 10 μm (panel b, inset). Simultaneous (c) electrophysiological and (d) optical recording of membrane potential changes evoked by optogenetic stimulation of the neuron expressing ChR2-YFP and stained with BeRST 1. Cyan light (475 nm LED, 80 mW/cm²) was provided in 5 ms pulses at a rate of 5 Hz. Magnified view of (e) electrophysiological and (f) optical recording of the action potential highlighted in the red, dotted box from panels (c) and (d). Optical sampling rate was 1 kHz.




Figure 8. Using BeRST 1 and Chr2 to perturb network activity. Cultured rat hippocampal (c) neurons transfected with (a) Chr2-YFP and stained with (b) BeRST 1 were stimulated with 475 nm light (80 mW/cm^2 , 5 ms, 5 Hz, cyan bars) during two separate 3 s periods to evoke activity in the Chr2-YFP-expressing cell. Scale bar is $20 \mu\text{m}$. (d) Schematic representation of neurons from DIC image in panel (c), color-coded to match the corresponding traces in (e–g). The blue Chr2(+) cell is depicted making possible connections to other neurons in the field of view. Optical records of BeRST 1 responses were acquired at 500 Hz with an sCMOS camera during (e) an optical recording session and (f) subsequent trial, separated by approximately 30 s (double hash). Numbers and colors of traces refer to specific neurons in panels (a–d). Red boxes indicate areas of the trace that have been magnified for clarity in panel (g). Dotted gray lines are provided in panel (g) to help visually estimate the spike timing of BeRST 1-stained neurons.

sought to establish that BeRST 1 and Chr2 represent optically orthogonal tools for probing and perturbing neuronal activity. Neurons transiently expressing a Chr2-YFP fusion⁵² (Figure 7a and SI Figure S7b) and loaded with BeRST 1 gave bright membrane staining, as expected (Figure 7b and SI Figure S7a). Stimulation of Chr2 with brief pulses of cyan light (5 ms, 475 nm, 80 mW/cm^2) resulted in light-induced spikes in membrane potential, as measured via patch clamp electrophysiology in either whole cell (Figure 7c and d) or cell-attached (SI Figure S7d) configurations. Control experiments in cells lacking Chr2 show no spiking (data not shown). Optical recording of light-induced spikes with BeRST 1 matched the electrophysiologically recorded activity (Figure 7c vs d and e vs f; SI Figure S7d and e). Of note is the high SNR achieved by BeRST 1 during optical recordings. Although the SNR of BeRST 1 is low compared to the highly sensitive whole-cell patch clamp method (Figure 7c–f), optical sensitivity of BeRST 1 compares favorably with records acquired by the less sensitive on-cell patch clamp technique (SI Figure S7d and e).

As a final demonstration of the efficacy of using BeRST 1 in conjunction with optogenetic tools, we sought to optically probe network behavior in cultured hippocampal neurons. Analysis of neuronal connectivity and function with cellular resolution in intact brains remains an outstanding challenge, making cultured hippocampal neurons an attractive model system for studying both connectivity and function. Indeed, experimental evidence shows that neurons in culture make specific synaptic connections,⁵³ mirroring connectivity encountered in vivo.

We imaged a large area of hippocampal neurons ($370 \times 62 \mu\text{m}$, $20\times$ magnification) stained with BeRST 1 and containing just a single neuron expressing Chr2-YFP (Figure 8a–c and SI Figure S8). Continuous optical recording over a period of several seconds demonstrated, as before (Figure 4) that most neurons remain relatively silent during the recording epoch. Activation of the Chr2-YFP-expressing neuron with cyan light (475 nm , 80 mW/cm^2 , 5 ms at 5 Hz, Figure 8e–g, cyan bars)

resulted, as before (Figure 7), in bursts of spikes from the Chr2(+) neuron (Figure 8, blue trace) that exactly followed the optical stimulation pattern. Unlike previous experiments, in which the optical response of only the Chr2(+) cell was captured (Figure 7 and SI Figure S7), the larger recording area, allowed us to observe how activation of a single neuron affected the surrounding cells (Figure 8 and SI Movies S2 and S3).

During the first optical stimulation (Figure 8e and f, cyan bars; SI Movie S2), Neuron 1 (Figure 8c–e, black cell and trace)—the closest neighbor to Chr2-YFP-expressing Neuron 2 (Figure 8c–e, blue cell and trace)—fires a single AP in response to the second AP in Neuron 2 (Figure 8e, first red box and Figure 8g, first region). Neurons 3 (Figure 8c–e, purple cell and trace) and 5 (Figure 8c–e, orange cell and trace) are only weakly coupled to Neuron 2's firing and appear to initiate their own doublet of spikes that propagate into Neuron 4 (Figure 8c–e, green cell and trace) and interrupt the optically induced firing pattern of Neuron 2 (Figure 8d and e, first red box). This spike doublet is separated by approximately 12 ms and would be difficult to resolve with slower Ca^{2+} imaging.

During the second epoch of light stimulation, the response of Neuron 1 (Figure 8d and f, black cell and trace; SI Movie S3) is suppressed, and no longer appears strongly coupled to cues from optically controlled Neuron 2 (Figure 8d and f, blue cell and trace). The response of Neurons 3 and 5 are more strongly correlated to the firing of Neuron 2 (Figure 8f and g, first red box), while Neuron 4 (Figure 8d and f, green cell and trace) remains silent with the exception of several spontaneous bursts prior to optical activation of Neuron 2. Even in this “simple” cultured hippocampal model, a high degree of interconnectedness exists, which would not be observable through methods like Ca^{2+} imaging. Together, these experiments establish the ability of BeRST 1 to work in concert with classic optogenetic tools to optically interrogate the complexity of neuronal connectivity and function.

CONCLUDING REMARKS

We have described the design, synthesis, and characterization of a new sulfonated Si-rhodamine for use in voltage sensing applications. The first member of this new class of sulfonated far-red to NIR fluorophores, Berkeley Red Sensor of Transmembrane potential 1, or BeRST 1, displays bright, membrane-localized fluorescence, is highly photostable, and extremely voltage-sensitive. We show that BeRST 1 can detect action potentials in cultured hippocampal neurons and that BeRST 1 interfaces readily with a number of other optical tools, making BeRST 1 ideally suited for multicolor imaging with GFP, Ca²⁺ indicators like GCaMP, voltage sensors based on cpGFP such as ASAP1, as well as optogenetic tools like ChR2. Use of BeRST 1 alongside ChR2 permits noninvasive recording and control of membrane potential in single cells and in neuronal microcircuits.

There has been a resurgence, of late, in methods to optically monitor membrane potential, with many promising candidates arriving in the form of new genetically encoded voltage sensitive fluorescent proteins and opsins. Small molecule probes of membrane potential fill a vital role in this regard, as molecular wire PeT voltage sensors require no genetic manipulation, can be delivered to all cell types, respond rapidly to fast voltage changes, and as we show here, can be tuned across a wide color range. We have established that molecular wire PeT voltage sensing based on a phenylenevinylene molecular wire platform accommodates multiple types of fluorophores and that new far-red to NIR voltage sensors provide a convenient interface with well-established optical tools. We anticipate that BeRST 1 will prove to be a valuable tool in efforts to map membrane potential dynamics in a variety of systems.

EXPERIMENTAL SECTION

General Methods for Chemical Synthesis. Chemical reagents and solvents (dry) were purchased from commercial suppliers and used without further purification. All reactions were carried out in oven-dried flasks under N₂. Thin layer chromatography (TLC) (Silicycle, F254, 250 μm) and preparative thin layer chromatography (PTLC) (Silicycle, F254, 1000 μm) were performed on glass backed plates precoated with silica gel and were visualized by fluorescence quenching under UV light. Flash column chromatography was performed on Silicycle Silica Flash F60 (230–400 Mesh) using a forced flow of air at 0.5–1.0 bar. NMR spectra were measured on Bruker AVB-400 MHz, 100 MHz, AVQ-400 MHz, 100 MHz, Bruker AV-600 MHz, 150 MHz. NMR spectra measured on Bruker AVII-900 MHz, 225 MHz, equipped with a TCI cryoprobe accessory, were performed by Dr. Jeffrey Pelton (QB3). Chemical shifts are expressed in parts per million (ppm) and are referenced to CDCl₃ 7.26 ppm, 77.0 ppm. Coupling constants are reported as Hertz (Hz). Splitting patterns are indicated as follows: s, singlet; d, doublet; sep, septet; dd, doublet of doublet; ddd, doublet of double of doublet; dt, doublet of triplet; td, triplet of doublet; m, multiplet. High-resolution mass spectra (ESI EI) were measured by the QB3/Chemistry mass spectrometry service at University of California, Berkeley.

Isopropyl 2,5-dibromobenzenesulfonate (2). 1,4-Diazabicyclo-[2.2.2]octane (0.81 g, 7.2 mmol, 1.2 equiv) was added to 2,5-dibromobenzenesulfonyl chloride⁵⁴ (2.0 g, 6.0 mmol, 1.0 equiv) in 3:2 CH₂Cl₂/PrOH (v/v) (5.0 mL) slowly at 0 °C. White precipitate started to form. The reaction mixture was warmed to RT and stirred for 30 min. After the precipitate was removed by filtration, the filtrate was concentrated under reduced pressure and purified by flash column chromatography (5:1 hexanes/EtOAc) to afford the product as a white solid (1.3 g, 3.7 mmol, 62%). ¹H NMR (400 MHz, CDCl₃) δ 8.25 (d, J = 2.3 Hz, 1H), 7.63 (d, J = 8.4 Hz, 1H), 7.57 (dd, J = 8.4, 2.3 Hz, 1H), 4.86 (sep, J = 6.3 Hz, 1H), 1.38 (d, J = 6.3, 6H); ¹³C NMR (150 MHz,

CDCl₃) δ 138.8, 137.2, 136.8, 134.3, 121.4, 119.5, 79.3, 22.8; HRMS (EI) calcd for C₉H₁₀Br₂O₃S⁺ [M]⁺ 355.8712, found, 355.8717.

5-Bromo-2-(7-(dimethylamino)-3-(dimethyliminio)-5,5-dimethyl-3,5-dihydrodibenzo[b,e]silin-10-yl) benzenesulfonate (bromo-Berkeley Red). To a CH₂Cl₂ (1.0 mL) solution of 3³⁶ (0.10 g, 0.31 mmol, 1.0 equiv), trifluoromethanesulfonic anhydride (0.10 mL, 0.59 mmol, 1.9 equiv) was added. This solution was stirred at RT for 30 min (Solution A). In another flask, 2 (0.33 g, 0.93 mmol, 3.0 equiv) was dissolved in CH₂Cl₂ (2.0 mL). The solution was cooled to -20 °C and *n*-BuLi (1.6 M in hexane, 0.58 mL, 0.93 mmol, 3.0 equiv) was added slowly. After 10 min, Solution A was added to the reaction mixture and stirred for 1 h. The crude isopropyl ester protected-product was obtained by PTLC (5:1:1 hexanes/EtOAc/CH₂Cl₂) and subjected to 1.0 M BBr₃ in CH₂Cl₂ (2.0 mL). The reaction mixture was stirred at RT overnight and purified by PTLC (10:1 CH₂Cl₂/CH₃OH) to afford the desired product as a dark blue solid (25 mg, 0.046 mmol, 15%). ¹H NMR (600 MHz, CDCl₃) δ 8.51 (d, J = 2.0 Hz, 1H), 7.53 (dd, J = 8.1, 2.0 Hz, 1H), 7.32 (d, J = 9.5 Hz, 2H), 6.99 (d, J = 2.8 Hz, 2H), 6.89 (d, J = 8.1 Hz, 1H), 6.58 (dd, J = 9.5, 2.8 Hz, 2H), 3.25 (s, 12H), 0.55 (apparent s, 3H), 0.53 (apparent s, 3H); ¹³C NMR (225 MHz, CDCl₃) δ 173.6, 153.9, 148.3, 148.2, 143.5, 134.3, 132.2, 131.1, 130.2, 129.0, 122.9, 119.4, 113.3, 40.6, -0.9, -1.2; HRMS (ESI) calcd for C₂₅H₂₇BrN₂NaO₃SSi⁺ [M + Na]⁺ 565.0587, found, 565.0590.

2-(7-(Dimethylamino)-3-(dimethyliminio)-5,5-dimethyl-3,5-dihydrodibenzo[b,e]silin-10-yl) benzenesulfonate (Berkeley Red). Berkeley Red was prepared according to the same procedure described above from 3 (50 mg, 0.15 mmol, 1.0 equiv), trifluoromethanesulfonic anhydride (26 μL, 0.15 mmol, 1.0 equiv), I⁵⁵ (83 mg, 0.30 mmol, 1.5 equiv) and *n*-BuLi (1.6 M in hexane, 0.19 mL, 0.30 mmol, 1.5 equiv). The product was obtained as a dark blue solid (6.3 mg, 0.014 mmol, 9%). ¹H NMR (600 MHz, CDCl₃) δ 8.35 (dd, J = 8.4, 1.3 Hz, 1H), 7.54 (ddd, J = 8.4, 7.5, 1.3 Hz, 1H), 7.39 (apparent td, J = 7.5, 1.3 Hz, 1H), 7.36 (d, J = 9.6 Hz, 2H), 6.99 (dd, J = 7.5, 1.3 Hz, 1H), 6.98 (d, J = 2.9 Hz, 2H), 6.56 (dd, J = 9.6, 2.9 Hz, 2H), 3.23 (s, 12H), 0.56 (apparent s, 3H), 0.53 (apparent s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ 175.8, 153.9, 148.2, 146.8, 143.8, 135.3, 129.4, 129.4, 128.9, 128.6, 127.9, 119.2, 113.3, 40.5, -0.9, -1.2; HRMS (ESI) calcd for C₂₅H₂₈N₂NaO₃SSi⁺ [M + Na]⁺ 487.1482, found, 487.1478.

5-((E)-4-((E)-4-(Dimethylamino)-2-methoxystyryl)styryl)-2-(7-(dimethylamino)-3-(dimethyliminio)-5,5-dimethyl-3,5-dihydrodibenzo[b,e]silin-10-yl)benzenesulfonate (BeRST 1). Bromo-Berkeley Red (5.0 mg, 9.2 μmol, 1.0 equiv), 4 (2.6 mg, 9.3 μmol, 1.0 equiv), Pd(OAc)₂ (1.1 mg, 4.5 μmol, 0.50 equiv), P(*o*-tol)₃ (2.8 mg, 9.2 μmol, 1.0 equiv) and NEt₃ (50 μL) were dissolved in DMF (0.10 mL). The solution was stirred at 75 °C and the progress of the reaction was monitored by TLC. After the reaction was complete, the crude mixture was purified by PTLC (10:1 CH₂Cl₂/CH₃OH) to afford the product as a dark green solid (4.2 mg, 5.7 μmol, 62%). ¹H NMR (900 MHz, 10:1 CDCl₃/CD₃OD (v/v)) δ 8.34 (d, J = 1.8 Hz, 1H), 7.53 (dd, J = 7.8, 1.8 Hz, 1H), 7.46 (apparent s, 4H), 7.43 (d, J = 8.5 Hz, 1H), 7.39 (d, J = 16.4 Hz, 1H), 7.27–7.21 (m, 3H), 7.14 (d, J = 16.4 Hz, 1H), 6.99–6.94 (m, 3H), 6.91 (d, J = 16.4 Hz, 1H), 6.55 (dd, J = 9.7, 2.8 Hz, 2H), 6.31 (dd, J = 8.5, 2.4 Hz, 1H), 6.19 (d, J = 2.4 Hz, 1H), 3.85 (s, 3H), 3.21 (s, 12H), 2.95 (s, 6H), 0.50 (apparent s, 3H) 0.48 (apparent s, 3H); ¹³C NMR (225 MHz, 10:1 CDCl₃/CD₃OD (v/v)) δ 173.3, 158.0, 153.9, 151.4, 148.1, 145.2, 143.3, 138.7, 138.5, 134.9, 134.4, 130.4, 129.6, 128.8, 127.1, 126.9, 126.2, 126.2, 126.2, 125.9, 124.1, 123.7, 119.3, 115.0, 113.2, 105.2, 95.6, 55.3, 40.4, 40.3, -1.2, -1.4; HRMS (ESI) calcd for C₄₄H₄₈N₃O₄SSi⁺ [M + H]⁺ 742.3129, found, 742.3118.

Spectroscopic Measurements. Stock solutions of BR and BeRST 1 were prepared in DMSO (1.0–10 mM) and diluted with TBS (50 mM Tris-HCl, pH 7.5, 0.15 M NaCl) solution containing 0.10% (w/w) SDS (1:100–1:1000 dilution). UV absorbance spectra were acquired on a Shimadzu 2501 Spectrophotometer (Shimadzu). Fluorescence spectra were recorded on a Quantamaster Master 4 L-format scanning spectrofluorometer (Photon Technologies International) equipped with an LPS-220B 75-W xenon lamp and power supply, A-1010B lamp housing with integrated igniter, switchable 814

photon-counting/analog photomultiplier detection unit, and MD5020 motor driver. Samples for absorption and fluorescence measurements were contained in 1 cm path length quartz cuvettes (Starna Cells). Relative quantum yields were measured in TBS solution containing 0.10% (w/w) SDS or in CH₃OH and referenced to cresyl violet in CH₃OH, which has a quantum yield of 0.54.⁵⁶

Cell Culture and Imaging. All animal procedures were approved by the UC Berkeley Animal Care and Use Committees and conformed to the NIH Guide for the Care and Use of Laboratory Animals and the Public Health Policy.

Human embryonic kidney 293T (HEK) cells were passaged and plated onto 12 mm glass coverslips precoated with Poly-D-Lysine (PDL; 1 mg/mL; Sigma-Aldrich) to provide a confluency of ~15% and 50% for electrophysiology and imaging, respectively. HEK cells were plated and maintained in Dulbecco's modified eagle medium (DMEM) supplemented with 4.5 g/L D-glucose, 10% FBS and 1% Glutamax.

Hippocampi were dissected from embryonic day 18 Sprague–Dawley rats (Charles River Laboratory) in cold sterile HBSS (zero Ca²⁺, zero Mg²⁺). All dissection products were supplied by Invitrogen, unless otherwise stated. Hippocampal tissue was treated with trypsin (2.5%) for 15 min at 37 °C. The tissue was triturated using fire polished Pasteur pipettes, in minimum essential media (MEM) supplemented with 5% fetal bovine serum (FBS; Thermo Scientific), 2% B-27, 2% 1 M D-glucose (Fisher Scientific) and 1% glutamax. The dissociated cells were plated onto 12 mm diameter coverslips (Fisher Scientific) pretreated with PDL (as above) at a density of 30–40 000 cells per coverslip in MEM supplemented media (as above). Neurons were maintained at 37 °C in a humidified incubator with 5% CO₂. At 3 days in vitro (DIV) half of the MEM supplemented media was removed and replaced with Neurobasal media containing 2% B-27 supplement and 1% glutamax. Transfection of genetic tools was carried out using Lipofectamine 3000 at 7 DIV. Functional imaging was performed on mature neurons 13–20 DIV, except electrophysiological experiments which were performed on 12–15 DIV neurons.

Unless stated otherwise, for loading of HEK cells and hippocampal neurons, BeRST 1 was diluted in DMSO to 1 mM, and then diluted 1:1000 in HBS (in mM) 140 NaCl, 2.5 KCl, 10 HEPES, 10 D-glucose 1.3 MgCl₂ and 2 CaCl₂; pH 7.3 and 290 mOsmol. All imaging experiments were performed in HBS.

General Imaging Parameters. Epifluorescence imaging was performed on an Axio Examiner Z-1 (Zeiss) equipped with a Spectra-X Light engine LED light (Lumencor), controlled with Slidebook (v6, Intelligent Imaging Innovations). Coincident excitation with multiple LEDs was controlled by Lumencor software triggered through a Digidata 1332A digitizer and pCLAMP 10 software (Molecular Devices). Images were acquired with either a W-Plan-Apo 20×/1.0 water objective (20×; Zeiss) or a W-Plan-Apo 63×/1.0 water objective (63×; Zeiss). Images were focused onto either an OrcaFlash4.0 sCMOS camera (sCMOS; Hamamatsu) or an eVolve 128 EMCCD camera (EMCCD; Photometrix). More detailed imaging information for each experimental application are detailed in the Supporting Information.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.5b06644.

Experimental details, including HPLC spectra, NMR spectra, imaging parameters, electrophysiology data, and supporting figures as referenced in the main text. (PDF) SI Movie 1. Voltage sensitivity of BeRST 1 in patch-clamped HEK cells. (AVI)

SI Movie 2. ΔF of BeRST 1 signals during all-optical electrophysiology in rat hippocampal neurons with BeRST 1 and Chr2, first optical stimulation train. (AVI)

SI Movie 3. ΔF of BeRST 1 signals during all-optical electrophysiology in rat hippocampal neurons with BeRST 1 and Chr2, second optical stimulation train. (AVI)

■ AUTHOR INFORMATION

Corresponding Author

*evanwmiller@berkeley.edu

Author Contributions

[†]Y.-L.H. and A.S.W. contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was generously supported by start-up funds from the University of California and by the NIH (R00NS078561). We thank Ms. Danxun Li and Mr. Kaveh Karbasi for assistance with automated analysis routines.

■ REFERENCES

- (1) Hamill, O. P.; Marty, A.; Neher, E.; Sakmann, B.; Sigworth, F. J. *Pflügers Arch.* **1981**, *391*, 85.
- (2) Spruston, N.; Jaffe, D. B.; Williams, S. H.; Johnston, D. J. *Neurophysiol.* **1993**, *70*, 781.
- (3) Peterka, D. S.; Takahashi, H.; Yuste, R. *Neuron* **2011**, *69*, 9.
- (4) Scanziani, M.; Hausser, M. *Nature* **2009**, *461*, 930.
- (5) Paredes, R. M.; Etzler, J. C.; Watts, L. T.; Zheng, W.; Lechleiter, J. D. *Methods* **2008**, *46*, 143.
- (6) Adams, S. R. *Cold Spring Harbor Protoc.* **2010**, DOI: 10.1101/pdb.top70.
- (7) Oheim, M.; van 't Hoff, M.; Feltz, A.; Zamaleeva, A.; Mallet, J. M.; Collot, M. *Biochim. Biophys. Acta, Mol. Cell Res.* **2014**, *1843*, 2284.
- (8) Rose, T.; Goltstein, P. M.; Portugues, R.; Griesbeck, O. *Front. Mol. Neurosci.* **2014**, *7*, 88.
- (9) Zhao, Y.; Araki, S.; Wu, J.; Teramoto, T.; Chang, Y. F.; Nakano, M.; Abdelfattah, A. S.; Fujiwara, M.; Ishihara, T.; Nagai, T.; Campbell, R. E. *Science* **2011**, *333*, 1888.
- (10) Grienberger, C.; Konnerth, A. *Neuron* **2012**, *73*, 862.
- (11) Vogelstein, J. T.; Packer, A. M.; Machado, T. A.; Sippy, T.; Babadi, B.; Yuste, R.; Paninski, L. *J. Neurophysiol.* **2010**, *104*, 3691.
- (12) Grewe, B. F.; Langer, D.; Kasper, H.; Kampa, B. M.; Helmchen, F. *Nat. Methods* **2010**, *7*, 399.
- (13) Miller, E. W.; Lin, J. Y.; Frady, E. P.; Steinbach, P. A.; Kristan, W. B., Jr.; Tsien, R. Y. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 2114.
- (14) Yan, P.; Acker, C. D.; Zhou, W. L.; Lee, P.; Bollensdorff, C.; Negrean, A.; Lotti, J.; Sacconi, L.; Antic, S. D.; Kohl, P.; Mansvelder, H. D.; Pavone, F. S.; Loew, L. M. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 20443.
- (15) Matiukas, A.; Mitrea, B. G.; Qin, M.; Pertsov, A. M.; Shvedko, A. G.; Warren, M. D.; Zaitsev, A. V.; Wuskell, J. P.; Wei, M. D.; Watras, J.; Loew, L. M. *Heart Rhythm* **2007**, *4*, 1441.
- (16) Bradley, J.; Luo, R.; Otis, T. S.; DiGregorio, D. A. *J. Neurosci.* **2009**, *29*, 9197.
- (17) Fromherz, P.; Hubener, G.; Kuhn, B.; Hinner, M. J. *Eur. Biophys. J.* **2008**, *37*, 509.
- (18) Jin, L.; Han, Z.; Platasa, J.; Wooltorton, J. R.; Cohen, L. B.; Pieribone, V. A. *Neuron* **2012**, *75*, 779.
- (19) St-Pierre, F.; Marshall, J. D.; Yang, Y.; Gong, Y.; Schnitzer, M. J.; Lin, M. Z. *Nat. Neurosci.* **2014**, *17*, 884.
- (20) Sjulson, L.; Miesenbock, G. *J. Neurosci.* **2008**, *28*, 5582.
- (21) Ghitani, N.; Bayguinov, P. O.; Ma, Y.; Jackson, M. B. *J. Neurophysiol.* **2015**, *113*, 1249.
- (22) Hochbaum, D. R.; Zhao, Y.; Farhi, S. L.; Klapoetke, N.; Werley, C. A.; Kapoor, V.; Zou, P.; Kralj, J. M.; Maclaurin, D.; Smedemark-Margulies, N.; Saulnier, J. L.; Boulting, G. L.; Straub, C.; Cho, Y. K.; Melkonian, M.; Wong, G. K.; Harrison, D. J.; Murthy, V. N.; Sabatini,

B. L.; Boyden, E. S.; Campbell, R. E.; Cohen, A. E. *Nat. Methods* **2014**, *11*, 825.

(23) Woodford, C. R.; Frady, E. P.; Smith, R. S.; Morey, B.; Canzi, G.; Palida, S. F.; Araneda, R. C.; Kristan, W. B., Jr.; Kubiak, C. P.; Miller, E. W.; Tsien, R. Y. *J. Am. Chem. Soc.* **2015**, *137*, 1817.

(24) Li, L. S. *Nano Lett.* **2007**, *7*, 2981.

(25) Moshtagh-Khorasani, M.; Miller, E. W.; Torre, V. *Physiol. Rep.* **2013**, *1*, e00089.

(26) Dolensek, J.; Stozer, A.; Skelin Klemen, M.; Miller, E. W.; Slak Rupnik, M. *PLoS One* **2013**, *8*, e82374.

(27) Gee, K. R.; Poot, M.; Klaubert, D. H.; Sun, W.-C.; Haugland, R. P.; Mao, F. WO9739064A1, Molecular Probes, Inc.: Eugene, OR, 1997; p 79.

(28) Nakai, J.; Ohkura, M.; Imoto, K. *Nat. Biotechnol.* **2001**, *19*, 137.

(29) Chen, T. W.; Wardill, T. J.; Sun, Y.; Pulver, S. R.; Renninger, S. L.; Baohan, A.; Schreiter, E. R.; Kerr, R. A.; Orger, M. B.; Jayaraman, V.; Looger, L. L.; Svoboda, K.; Kim, D. S. *Nature* **2013**, *499*, 295.

(30) Nagel, G.; Szellas, T.; Huhn, W.; Kateriya, S.; Adeishvili, N.; Berthold, P.; Ollig, D.; Hegemann, P.; Bamberg, E. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, *100*, 13940.

(31) Zhang, F.; Vierock, J.; Yizhar, O.; Fenno, L. E.; Tsunoda, S.; Kianianmomeni, A.; Prigge, M.; Berndt, A.; Cushman, J.; Polle, J.; Magnuson, J.; Hegemann, P.; Deisseroth, K. *Cell* **2011**, *147*, 1446.

(32) Weissleder, R. *Nat. Biotechnol.* **2001**, *19*, 316.

(33) Koide, Y.; Urano, Y.; Hanaoka, K.; Terai, T.; Nagano, T. *ACS Chem. Biol.* **2011**, *6*, 600.

(34) Fu, M. Y.; Xiao, Y.; Qian, X. H.; Zhao, D. F.; Xu, Y. F. *Chem. Commun.* **2008**, 1780.

(35) Murugesan, N.; Gu, Z.; Stein, P. D.; Bisaha, S.; Spergel, S.; Girotra, R.; Lee, V. G.; Lloyd, J.; Misra, R. N.; Schmidt, J.; Mathur, A.; Stratton, L.; Kelly, Y. F.; Bird, E.; Waldron, T.; Liu, E. C. K.; Zhang, R.; Lee, H.; Serafino, R.; Abboa-Offei, B.; Mathers, P.; Giancarli, M.; Seymour, A. A.; Webb, M. L.; Moreland, S.; Barrish, J. C.; Hunt, J. T. *J. Med. Chem.* **1998**, *41*, 5198.

(36) Pastierik, T.; Sebej, P.; Medalova, J.; Stacko, P.; Klan, P. *J. Org. Chem.* **2014**, *79*, 3374.

(37) Lukinavicius, G.; Umezawa, K.; Olivier, N.; Honigmann, A.; Yang, G. Y.; Plass, T.; Mueller, V.; Reymond, L.; Correa, I. R.; Luo, Z. G.; Schultz, C.; Lemke, E. A.; Heppenstall, P.; Eggeling, C.; Manley, S.; Johnsson, K. *Nat. Chem.* **2013**, *5*, 132.

(38) Uno, S. N.; Kamiya, M.; Yoshihara, T.; Sugawara, K.; Okabe, K.; Tarhan, M. C.; Fujita, H.; Funatsu, T.; Okada, Y.; Tobita, S.; Urano, Y. *Nat. Chem.* **2014**, *6*, 681.

(39) Akerlof, G.; Short, O. A. *J. Am. Chem. Soc.* **1936**, *58*, 1241.

(40) Krasovskiy, A.; Knochel, P. *Angew. Chem., Int. Ed.* **2004**, *43*, 3333.

(41) Cho, C. H.; Kim, I. S.; Park, K. Y. *Tetrahedron* **2004**, *60*, 4589.

(42) Abe, T.; Fujimori, T. *Dev Growth Differ* **2013**, *55*, 390.

(43) Shen, Y.; Lai, T.; Campbell, R. E. *Neurophotonics* **2015**, *2*, 031203.

(44) Johnson, L. V.; Walsh, M. L.; Chen, L. B. *Proc. Natl. Acad. Sci. U. S. A.* **1980**, *77*, 990.

(45) Loewe, H.; Urbaniet, J. *Arzneim.-Forsch./Drug Res.* **1974**, *24*, 1927.

(46) Park, S. A.; Lee, S. R.; Tung, L.; Yue, D. T. *Sci. Rep.* **2014**, *4*, 6125.

(47) Lin, J. Y.; Lin, M. Z.; Steinbach, P.; Tsien, R. Y. *Biophys. J.* **2009**, *96*, 1803.

(48) Zhang, F.; Prigge, M.; Beyriere, F.; Tsunoda, S. P.; Mattis, J.; Yizhar, O.; Hegemann, P.; Deisseroth, K. *Nat. Neurosci.* **2008**, *11*, 631.

(49) Prigge, M.; Schneider, F.; Tsunoda, S. P.; Shilyansky, C.; Wietek, J.; Deisseroth, K.; Hegemann, P. *J. Biol. Chem.* **2012**, *287*, 31804.

(50) Klapoetke, N. C.; Murata, Y.; Kim, S. S.; Pulver, S. R.; Birdsey-Benson, A.; Cho, Y. K.; Morimoto, T. K.; Chuong, A. S.; Carpenter, E. J.; Tian, Z.; Wang, J.; Xie, Y.; Yan, Z.; Zhang, Y.; Chow, B. Y.; Surek, B.; Melkonian, M.; Jayaraman, V.; Constantine-Paton, M.; Wong, G. K.; Boyden, E. S. *Nat. Methods* **2014**, *11*, 338.

(51) Lin, J. Y.; Knutsen, P. M.; Muller, A.; Kleinfeld, D.; Tsien, R. Y. *Nat. Neurosci.* **2013**, *16*, 1499.

(52) Zhang, F.; Wang, L. P.; Brauner, M.; Liewald, J. F.; Kay, K.; Watzke, N.; Wood, P. G.; Bamberg, E.; Nagel, G.; Gottschalk, A.; Deisseroth, K. *Nature* **2007**, *446*, 633.

(53) Williams, M. E.; Wilke, S. A.; Daggett, A.; Davis, E.; Otto, S.; Ravi, D.; Ripley, B.; Bushong, E. A.; Ellisman, M. H.; Klein, G.; Ghosh, A. *Neuron* **2011**, *71*, 640.

(54) Nakamura, T.; Yonesu, K.; Mizuno, Y.; Suzuki, C.; Sakata, Y.; Takuwa, Y.; Nara, F.; Satoh, S. *Bioorg. Med. Chem.* **2007**, *15*, 3548.

(55) Murugesan, N.; Gu, Z.; Stein, P. D.; Bisaha, S.; Spergel, S.; Girotra, R.; Lee, V. G.; Lloyd, J.; Misra, R. N.; Schmidt, J.; Mathur, A.; Stratton, L.; Kelly, Y. F.; Bird, E.; Waldron, T.; Liu, E. C.; Zhang, R.; Lee, H.; Serafino, R.; Abboa-Offei, B.; Mathers, P.; Giancarli, M.; Seymour, A. A.; Webb, M. L.; Hunt, J. T.; et al. *J. Med. Chem.* **1998**, *41*, 5198.

(56) Magde, D.; Brannon, J. H.; Cremers, T. L.; Olmsted, J. J. *Phys. Chem.* **1979**, *83*, 696.