


# Ratiometric Imaging Using a Single Dye Enables Simultaneous Visualization of Rac1 and Cdc42 Activation

Christopher J. MacNevin,<sup>†</sup> Alexei Touthkine,<sup>‡,†</sup> Daniel J. Marston, Chia-Wen Hsu,<sup>§</sup> Denis Tsygankov,<sup>||</sup> Li Li,<sup>⊥</sup> Bei Liu, Timothy Qi, Dan-Vinh Nguyen, and Klaus M. Hahn\*

Department of Pharmacology, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, United States


W Web-Enhanced Feature S Supporting Information

**ABSTRACT:** Biosensors that report endogenous protein activity in vivo can be based on environment-sensing fluorescent dyes. The dyes can be attached to reagents that bind selectively to a specific conformation of the targeted protein, such that binding leads to a fluorescence change. Dyes that are sufficiently bright for use at low, nonperturbing intracellular concentrations typically undergo changes in intensity rather than the shifts in excitation or emission maxima that would enable precise quantitation through ratiometric imaging. We report here **mero199**, an environment-sensing dye that undergoes a 33 nm solvent-dependent shift in excitation. The dye was used to generate a ratiometric biosensor of Cdc42 (**CRIB199**) without the need for additional fluorophores. **CRIB199** was used in the same cell with a FRET sensor of Rac1 activation to simultaneously observe Cdc42 and Rac1 activity in cellular protrusions, indicating that Rac1 but not Cdc42 activity was reduced during tail retraction, and specific protrusions had reduced Cdc42 activity. A novel program (EdgeProps) used to correlate localized activation with cell edge dynamics indicated that Rac1 was specifically reduced during retraction.



## INTRODUCTION

Fluorescent biosensors elucidate the flow of information through signaling networks in living cells and animals.<sup>1</sup> To minimize intracellular biosensor concentration, biosensors ideally are bright and fluoresce at wavelengths longer than cellular autofluorescence. Biosensors based on solvent-sensitive fluorescent dyes may be constructed by attaching the dye directly to the protein of interest, where fluorescence changes are associated with changes in conformation.<sup>2,3</sup> Alternatively, solvent-sensitive dyes may be attached to “affinity reagents” that bind selectively to a given state of endogenous target proteins, leading to fluorescence change.<sup>4–7</sup> Biosensors based on solvent-sensitive dyes offer substantially enhanced sensitivity over the more common biosensors based on fluorescence resonance energy transfer (FRET), in that bright dyes can be directly excited. However, bright dyes are more likely to respond to environment changes with shifts in fluorescence intensity rather than shifts in excitation/emission maxima, as there is an inverse relationship between dye brightness and the extent of solvent-dependent wavelength shifts.<sup>8–10</sup> Intensity changes are difficult to measure in cells because intensity is subject to multiple sources of artifacts (e.g., uneven illumination and variations in cell thickness). This has been overcome using ratiometric imaging (Figure 1a), in which a second, minimally responsive fluorophore is attached to the biosensor. Ratiometric imaging is challenging because it requires site-specific attachment of two fluorophores without perturbing protein



**Figure 1.** Ratiometric imaging with a dual versus single fluorophore biosensor. (a) The affinity reagent (AR) binds to the protein of interest (POI) only when the POI is in the active state. The changing intensity of the dye (red) relative to the fixed intensity of the fluorescent protein (FP) reflects POI binding. (b) POI activation is reflected simply in the ratio of emission at two different **mero199** excitation wavelengths.

activity, and quantitation is complicated when the two fluorophores bleach at different rates. Importantly, two dyes use up more of the wavelength spectrum than would a single fluorophore, making it difficult to use multiple biosensors in the

Received: September 16, 2015

Published: February 10, 2016


same cell. We describe here **mero199**, a bright, long wavelength dye that undergoes solvent-dependent changes in its excitation maxima, enabling ratio imaging with a single dye (Figure 1b).

Among environment-sensing small molecule fluorophores, merocyanine dyes are especially well suited for live-cell imaging. They can be bright, can emit at wavelengths that minimally overlap with cellular autofluorescence, and can exhibit solvent-dependent changes in extinction coefficient, fluorescence quantum yield (QY), or excitation/emission maxima.<sup>11,12</sup> Merocyanine dyes incorporate electron-donor and -acceptor moieties linked by conjugation.<sup>13</sup> The photophysical properties of the dyes depend on the specific donor/acceptor combination and on the nature of the conjugation.<sup>8,9,14</sup> Previous studies indicate that brightness is maximized when the ground state is comprised of equal contributions from zwitterionic and nonpolar resonance forms (the so-called cyanine limit, Scheme 1a), while solvent sensitivity is maximized when the

that their ground state is more polar than their excited state.<sup>17</sup> Dyes with the indolenine donor are brighter not only because they can approach the cyanine limit but also because the indolenine imparts extended conjugation and its geminal dimethyl substitution reduces aggregation-induced fluorescence quenching. We sought to produce a dye that combined the brightness of the indolenine donor with the solvent response of the quinoline donor, leading to the design of **mero199** (Scheme 1), a dye based on a fused pyrido-indolium donor heterocycle.<sup>8</sup> Like the quinoline, aromaticity is enhanced when the dye is in the charged form, favoring solvent sensitivity, but the predominance of the charged resonance form is not as strong as in quinoline itself. This structure proved to be substantially brighter than the quinoline-containing dyes, likely because of the conjugation provided by the indolenine's six-membered ring and reduced aggregation.

Photobleaching was an important consideration in the development of **mero199**, as our survey of donor/acceptor combinations had also shown an inverse relationship between solvent sensitivity and photostability.<sup>9</sup> Photodegradation of merocyanine dyes proceeds through attack of singlet oxygen at the most nucleophilic  $\alpha$ -carbon of the central polymethine chain, and the reactivity of this position is enhanced for dyes further from the cyanine limit.<sup>18</sup> A barbituric acid acceptor produced the best compromise between solvent sensitivity, brightness, and photostability (data not shown). Substitution with an electron-withdrawing group at the nucleophilic position of the polymethine chain can significantly increase photostability, so we incorporated the cyano group in the **mero199** structure.<sup>14,18</sup> Finally, a sulfonate group was included for water solubility, and a cysteine-reactive iodoacetamide group was included for site-specific labeling of proteins or other affinity reagents.<sup>19</sup>

Scheme 1. Rational Design of Ratiometric Dye **mero199**<sup>a</sup>




<sup>a</sup>A fused pyrido-indolium donor heterocycle was included for enhanced solvatochromic properties.

zwitterionic or nonpolar ground states predominate.<sup>10,15,16</sup> Consistent with this model, our earlier studies showed that the 3,3-dimethylindolenine donor heterocycle (Scheme 1b) leads to extraordinarily bright dyes when combined with several acceptors, but these dyes all show limited solvent-dependent changes in fluorescence.<sup>9</sup> In contrast, we found that pyridine donors and quinoline donors (Scheme 1c) could produce exceptionally large solvent-dependent shifts in fluorescence maxima, but these dyes were too dim for practical live-cell imaging.<sup>9</sup> Dyes with pyridine and quinoline donors likely favor the ground-state zwitterionic resonance form rather than the cyanine limit, because the zwitterionic form increases the aromaticity of the donor heterocycle (unlike the indolenine donor). The quinoline and pyridine-containing dyes showed hypsochromic shifts with increasing solvent polarity,<sup>9</sup> indicating

## RESULTS AND DISCUSSION

Synthesis of **mero199** began with condensation of the commercially available 2,3,3-trimethylindolenine (**1**, Scheme 2) with *N*-hydroxymethyl-phthalimide, followed by preparation of the product as its perchlorate salt to give compound **2**. Conjugate addition of methyl vinyl ketone to compound **2** was followed by heating of the open chain adduct in pyridine to give the fused pyridinium salt (**3**).<sup>20</sup> To ensure protecting group compatibility with subsequent reaction conditions, the phthalimide group was removed, and the resulting primary amine was reprotected with Boc, followed by base promoted addition of *N,N*-dimethylformamide dimethyl acetal to give the enamine **4**. Treatment of compound **4** with hydroxylamine-*O*-sulfonic acid resulted in expulsion of dimethylamine to afford the nitrile-substituted product<sup>21</sup> that was subsequently reacted with a barbituric acid bearing a methyl enol ether to give the merocyanine intermediate **5**.<sup>18</sup> The Boc group was removed under mild neutral conditions using iodotrimethylsilane to give compound **6**. The free amine of compound **6** was then reacted with 1,3-propane sultone, followed by treatment with iodoacetic anhydride to provide the desired iodoacetamide-substituted dye, **mero199**, in a total of 12 linear steps.

We characterized the effects of solvent polarity on the fluorescence properties of **mero199** using previously reported methods (Table 1, Figure 2).<sup>9</sup> Increased solvent polarity led to hypsochromic shifts in both excitation and emission maxima, with a difference of 33 nm between the excitation maxima in water and butanol. Extinction coefficients in methanol and butanol were among the highest we have observed ( $2.1 \times 10^5$

Scheme 2. Synthesis of mero199<sup>a</sup>

<sup>a</sup>Reagents and conditions: (a) H<sub>2</sub>SO<sub>4</sub>, *N*-hydroxymethyl-phthalimide, 0 °C to RT, 50%; (b) HClO<sub>4</sub>, EtOH, reflux, 90%; (c) methyl vinyl ketone, 100 °C, 62%; (d) pyridine, reflux, 28%; (e) NH<sub>2</sub>NH<sub>2</sub>·H<sub>2</sub>O 3:2 CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 98%; (f) Boc<sub>2</sub>O, Et<sub>3</sub>N, 3:1 CH<sub>2</sub>Cl<sub>2</sub>/ACN; (g) (CH<sub>3</sub>O)<sub>2</sub>CHN(CH<sub>3</sub>)<sub>2</sub>, cat. DBU, DMF, 100 °C, 68% from 3; (h) NH<sub>2</sub>SO<sub>3</sub>H, 1:1 MeOH/H<sub>2</sub>O; (i) 5-(3-methoxy-allylidene)-1,3-dimethyl-pyrimidine-2,4,6-trione, NaOAc, 1:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH, reflux, 29% from 4; (j) (CH<sub>3</sub>)<sub>3</sub>SiI, ACN, 87%; (k) 1,3-propane sultone, NMP, NaOAc; (l) (ICH<sub>2</sub>CO)<sub>2</sub>O, NMP, 30% from 6.

Table 1. Photophysical Characterization of mero199

solvent	exc $\lambda_{\max}$ (nm)	em $\lambda_{\max}$ (nm)	$\epsilon$ (M <sup>-1</sup> cm <sup>-1</sup> )	QY	$\epsilon \times$ QY
water	565	606	48000	0.05	2400
methanol	588	616	211000	0.07	14700
butanol	598	622	185000	0.22	40500




Figure 2. Excitation (a) and emission (b) fluorescence spectra of mero199 in selected solvents.

and  $1.9 \times 10^5$  M<sup>-1</sup> cm<sup>-1</sup>, respectively), and the dye's relative brightness ( $\epsilon \times$  QY) compared favorably with other merocyanines that have been used in dye-based biosensors.<sup>2,5,9</sup>

In water versus butanol, the dye showed a >16-fold change in brightness. The photostability of mero199 was 5.8-fold greater than the reference dye fluorescein (Figure S1). Together, these studies demonstrated that mero199 possessed the desired combination of solvent-sensitive fluorescence, brightness, water solubility, and photostability to be used in ratiometric live-cell biosensor applications.

We next tested mero199 by using it to make an actual biosensor, based on our previous approach to visualize the nucleotide state of the small GTPase Cdc42 in live cells.<sup>4</sup> Cdc42 is a member of the rho family of GTPases, proteins that regulate diverse aspects of cell behavior including cytoskeletal dynamics and migration.<sup>22</sup> The Cdc42/Rac interactive binding (CRIB) domain derived from the Wiskott–Aldrich syndrome protein (WASP) binds only to the “activated,” GTP-bound conformation of Cdc42.<sup>23</sup> This domain was mutated to include a single cysteine at position 271, a residue known to be near a hydrophobic pocket generated upon binding of the CRIB domain to activated Cdc42.<sup>24</sup> Previous studies showed that environment-sensing fluorophores at this position exhibited significant changes in fluorescence intensity when the labeled CRIB bound to activated Cdc42 in cells.<sup>4</sup> Previously used dyes changed intensity but not excitation/emission maxima, so it had been necessary to attach a second, nonresponsive fluorophore to the biosensor for ratio imaging.<sup>4,25</sup> Our new biosensor, named CRIB199, was based on attachment of mero199 at the same position. It was assessed in vitro using the methods previously validated for the two fluorophore biosensor.<sup>9</sup> CRIB199 showed a 12 nm shift in excitation maximum upon binding Cdc42, while the emission maximum remained at 618 nm (Figure S2). Fluorescence intensity increased nearly 4-fold upon Cdc42 binding. The dissociation constant ( $K_d$ ) of the Cdc42-biosensor conjugate ( $219 \pm 87$  nM) was slightly higher than that of unlabeled CRIB binding to Cdc42 ( $77 \pm 9$  nM), as with the previously reported dye-based biosensor (Figure S3).<sup>26,4,9</sup> Cdc42 binding induced a maximum decrease in emission at 577 nm and maximum increase in emission at 605 nm (Figure S4), leading us to use these two wavelengths for ratio imaging.

To test the ability of CRIB199 to report Cdc42 activation in living cells, the biosensor was microinjected into mouse embryonic fibroblasts (MEFs). For ratio imaging, we obtained two images at each time point, one excited with a 577/10 bandpass filter and the other with a 605/15 nm filter, and used a 630/45 nm emission filter for both. Dividing the two images on a pixel by pixel basis as previously described<sup>27</sup> yielded a map of ratio values across the cell (Figure 3). As described previously, biosensor concentrations were adjusted to report Cdc42 activity while minimizing perturbation of normal Cdc42 signaling.<sup>5,27–29</sup> Consistent with earlier studies, Cdc42 was locally activated in protrusions. A control biosensor with point mutations known to reduce the CRIB domain's affinity for




Figure 3. Migrating MEF cells microinjected with CRIB199, the biosensor based on mero199 conjugated to the CRIB domain, show Cdc42 activity in protrusions at the leading edge. Cells containing a mutant biosensor with reduced affinity for Cdc42 (mutantCRIB199) showed substantially less Cdc42 activation (images are scaled identically to show this). The pseudocolor scale shows the increase in ratio relative to the lowest 5% of ratio values in the cell (see Figure S5). These low values were similar for the two biosensors. The white arrow indicates the direction of movement.




active Cdc42 (mutant CRIB199, H246D and H249D)<sup>4,30</sup> reduced the level of Cdc42 activation seen in the cells (Figures 3 and S5). These studies showed that a biosensor made with **mero199** could quantify protein activity in living cells using ratiometric imaging without the need for attachment of a second fluorophore.

One important advantage of CRIB199 is that it uses substantially less of the wavelength spectrum than biosensors based on two fluorophores (e.g., for FRET). To quantify the relative kinetics and subcellular localizations of two complex protein activities, it is valuable to visualize two protein activities in the same cell. Simultaneous imaging can be simplified using biosensors based on **mero199**. To demonstrate this, we studied activation of Cdc42 and Rac1 by combining CRIB199 with a FRET-based biosensor of Rac1<sup>29</sup> that uses wavelengths typical of the majority of genetically encoded FRET sensors (Figure 4, see also Movie 1). Rac1 and Cdc42 both regulate the dynamics of the cell edge during motility, and both are active in protrusions.<sup>29</sup> It has been shown that there are a variety of protrusions with different functions and differences in signaling,<sup>31</sup> but in many cases these cannot be distinguished based simply on inspection of protrusion morphology. We reasoned

that Cdc42 and Rac1 would not be activated similarly in all protrusions, enabling us to distinguish live-cell protrusions of different function and/or driven by different mechanisms.

We were concerned that loading two biosensors in the same cell would affect normal morphodynamics. Quantifying protrusion velocity distributions and the extent of protrusion versus retraction indicated that the two biosensors could be used simultaneously with little perturbation of protrusion behavior (Figure S6). Variation in the relative activity of Cdc42 and Rac1 among different protrusions was apparent even from visual inspection of ratio images. The retracting tails of moving cells showed more Cdc42 than Rac1 activity, while protrusions at the leading edge varied; both Cdc42 and Rac1 were elevated in some protrusions, while only Rac1 predominated in others (Figure 4a). This demonstrated the ability of simultaneous imaging to distinguish protrusions on the basis of internal signaling activity. The differences between leading edge protrusions suggested that they had different functions or behaviors despite similar morphology. Activation of Cdc42 in only some protrusions is consistent with its role in controlling the direction of cell movement.<sup>22</sup>

The absence of Rac1 in retracting cell tails suggested that retraction was associated with reduced Rac1 activity. This is consistent with Rac1's known role in polymerizing dendritic actin to drive protrusion. To probe this, we developed a software tool (EdgeProps, see Supporting Information) that correlated biosensor activity with the velocity of neighboring regions of the cell edge. For a series of equally spaced points covering the entire edge, the program determined activation level and velocity at each point over time. This analysis revealed that both Cdc42 and Rac1 activity increased with velocity during protrusion, but that Rac1 activity was reduced relative to Cdc42 during retraction (Figure 4b). Cdc42 has been shown to be required for actomyosin contractility at the cell edge and to be involved in stimulated cell repulsion events, in accordance with these observations.<sup>32,33</sup>



**Figure 4.** Migrating MEF cells containing both a FRET-based Rac1 biosensor and CRIB199, a Cdc42 biosensor based on **mero199** conjugated to the CRIB domain of WASP. (a) Ratiometric images of Cdc42 and Rac1 activity (dotted box = cell tail undergoing retraction, solid box = protrusion along the leading edge). Note that the leading edge protrusion in the box shows high Cdc42 and Rac1 activity, while the protrusion at upper right has similar Rac1 activity but reduced Cdc42 activity. (b) Plot of activation versus cell edge velocity for Rac1 versus Cdc42 ( $n = 3$  cells). The readouts of the two biosensors were normalized such that when edge velocity was zero, both biosensors were set to zero, and the biosensors' maximum values (which in both cases occurred for protrusions at maximum velocity) were set equal to 1.

## CONCLUSIONS

In summary, we have developed **mero199**, a novel dye that enables construction of live-cell biosensors using a single dye, yet is capable of ratio imaging. Building a biosensor by using only a single fluorophore facilitates combination with other molecular imaging tools; the excitation and emission wavelengths of **mero199** are compatible with pairs of fluorescent proteins used in FRET biosensors, and **mero199** excitation does not overlap that of photoresponsive proteins used for optogenetics. GTPases, and many other signaling proteins, show heterogeneous and complex responses in living cells. Simultaneous imaging will facilitate the identification of dynamic cellular structures on the basis of multiple signaling activities and can precisely define the relative localizations and kinetics of activation for two proteins. Dye-based biosensors have seen less use than biosensors based on FRET, likely because they are more cumbersome to introduce into cells. However, they have important advantages in brightness, sensitivity, and their ability to probe different aspects of the intracellular environment. We hope that dye-based biosensors will soon be more accessible given recent efforts to covalently derivatize proteins with great specificity in living cells.<sup>34–36</sup>

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.5b09764.

Supporting figures, molecular biology, imaging and image analysis methods, experimental procedures and spectroscopic data for new compounds (PDF)

### Web-Enhanced Feature

A movie of simultaneous Cdc42 and Rac1 activity in AVI format is available in the HTML version of the paper.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*khahn@med.unc.edu

### Present Addresses

<sup>‡</sup>Sigma-Aldrich Co., Natick, Massachusetts 01760.

<sup>§</sup>Division of Pre-Clinical Innovation, National Center for Advancing Translational Sciences, National Institutes of Health, Bethesda, Maryland 20892.

<sup>||</sup>The Wallace H. Coulter Department of Biomedical Engineering at Georgia Institute of Technology and Emory University, Atlanta, Georgia 30332.

<sup>⊥</sup>Departments of Drug Discovery & Biomedical Sciences, Biochemistry & Molecular Biology, and Medicine, Medical University of South Carolina, Charleston, South Carolina 29425.

### Author Contributions

<sup>†</sup>These authors contributed equally.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was supported by funding from the American Cancer Society (CJM 119169-PF-10-183-01-TBE) and the National Institutes of Health (P01-GM103723 and P41-EB002025).

## ■ REFERENCES

- (1) Sample, V.; Mehta, S.; Zhang, J. *J. Cell Sci.* **2014**, *127*, 1151.
- (2) Hahn, K.; Debiasio, R.; Taylor, D. L. *Nature* **1992**, *359*, 736.
- (3) Garrett, S. C.; Hodgson, L.; Rybin, A.; Touthkine, A.; Hahn, K. M.; Lawrence, D. S.; Bresnick, A. R. *Biochemistry* **2008**, *47*, 986.
- (4) Nalbant, P.; Hodgson, L.; Kraynov, V.; Touthkine, A.; Hahn, K. M. *Science* **2004**, *305*, 1615.
- (5) Gulyani, A.; Vitriol, E.; Allen, R.; Wu, J.; Gremyachinskiy, D.; Lewis, S.; Dewar, B.; Graves, L. M.; Kay, B. K.; Kuhlman, B.; Elston, T.; Hahn, K. M. *Nat. Chem. Biol.* **2011**, *7*, 437.
- (6) Szent-Gyorgyi, C.; Schmidt, B. A.; Creeger, Y.; Fisher, G. W.; Zakel, K. L.; Adler, S.; Fitzpatrick, J. A. J.; Woolford, C. A.; Yan, Q.; Vasilev, K. V.; Berget, P. B.; Bruchez, M. P.; Jarvik, J. W.; Waggoner, A. *Nat. Biotechnol.* **2008**, *26*, 235.
- (7) Chen, C. A.; Yeh, R. H.; Yan, X.; Lawrence, D. S. *Biochim. Biophys. Acta, Proteins Proteomics* **2004**, *1697*, 39.
- (8) Touthkine, A.; Han, W. G.; Ullmann, M.; Liu, T. Q.; Bashford, D.; Noodleman, L.; Hahn, K. M. *J. Phys. Chem. A* **2007**, *111*, 10849.
- (9) MacNevin, C. J.; Gremyachinskiy, D.; Hsu, C. W.; Li, L.; Rougie, M.; Davis, T. T.; Hahn, K. M. *Bioconjugate Chem.* **2013**, *24*, 215.
- (10) Bublitz, G. U.; Ortiz, R.; Marder, S. R.; Boxer, S. G. *J. Am. Chem. Soc.* **1997**, *119*, 3365.
- (11) Kulinich, A. V.; Derevyanko, N. A.; Ishchenko, A. A. *Russ. J. Gen. Chem.* **2006**, *76*, 1441.
- (12) Kulinich, A. V.; Ishchenko, A. A. *Russ. Chem. Rev.* **2009**, *78*, 141.
- (13) Shirinian, V. Z.; Shimkin, A. A. *Top. Het. Chem.* **2008**, *14*, 75.
- (14) Touthkine, A.; Kraynov, V.; Hahn, K. *J. Am. Chem. Soc.* **2003**, *125*, 4132.
- (15) Ortiz, R.; Marder, S. R.; Cheng, L. T.; Tiemann, B. G.; Cavagnero, S.; Ziller, J. W. *J. Chem. Soc., Chem. Commun.* **1994**, 2263.
- (16) Wurthner, F.; Archetti, G.; Schmidt, R.; Kuball, H. G. *Angew. Chem., Int. Ed.* **2008**, *47*, 4529.
- (17) Buncel, E.; Rajagopal, S. *Acc. Chem. Res.* **1990**, *23*, 226.
- (18) Touthkine, A.; Nguyen, D. V.; Hahn, K. M. *Org. Lett.* **2007**, *9*, 2775.
- (19) Touthkine, A.; Nguyen, D. V.; Hahn, K. M. *Bioconjugate Chem.* **2007**, *18*, 1344.
- (20) Chapman, D. D.; Elwood, J. K.; Heseltine, D. W.; Hess, H. M.; Kurtz, D. W. *J. Org. Chem.* **1977**, *42*, 2474.
- (21) Biere, H.; Russe, R. *Tetrahedron Lett.* **1979**, *20*, 1361.
- (22) Jaffe, A. B.; Hall, A. *Annu. Rev. Cell Dev. Biol.* **2005**, *21*, 247.
- (23) Symons, M.; Derry, J. M. J.; Karlak, B.; Jiang, S.; Lemahieu, V.; McCormick, F.; Francke, U.; Abo, A. *Cell* **1996**, *84*, 723.
- (24) Abdul-Manan, N.; Aghazadeh, B.; Liu, G. A.; Majumdar, A.; Ouerfelli, O.; Siminovitch, K. A.; Rosen, M. K. *Nature* **1999**, *399*, 379.
- (25) Goguen, B. N.; Loving, G. S.; Imperiali, B. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 5058.
- (26) Rudolph, M. G.; Bayer, P.; Abo, A.; Kuhlmann, J.; Vetter, I. R.; Wittinghofer, A. *J. Biol. Chem.* **1998**, *273*, 18067.
- (27) Hodgson, L.; Nalbant, P.; Shen, F.; Hahn, K. *Methods Enzymol.* **2006**, *406*, 140.
- (28) El-Sibai, M.; Nalbant, P.; Pang, H.; Flinn, R. J.; Sarmiento, C.; Macaluso, F.; Cammer, M.; Condeelis, J. S.; Hahn, K. M.; Backer, J. M. *J. Cell Sci.* **2007**, *120*, 3465.
- (29) Machacek, M.; Hodgson, L.; Welch, C.; Elliott, H.; Pertz, O.; Nalbant, P.; Abell, A.; Johnson, G. L.; Hahn, K. M.; Danuser, G. *Nature* **2009**, *461*, 99.
- (30) Miki, H.; Sasaki, T.; Takai, Y.; Takenawa, T. *Nature* **1998**, *391*, 93.
- (31) Condeelis, J. *Annu. Rev. Cell Biol.* **1993**, *9*, 411.
- (32) Wilkinson, S.; Paterson, H. F.; Marshall, C. J. *Nat. Cell Biol.* **2005**, *7*, 255.
- (33) Groeger, G.; Nobes, C. D. *Biochem. J.* **2007**, *404*, 23.
- (34) Plass, T.; Schultz, C. *Springer Series on Fluorescence*; Springer: Heidelberg, 2011; Vol. 10, p 225 10.1007/978-3-642-18035-4\_7.
- (35) Uttamapinant, C.; Howe, J. D.; Lang, K.; Beranek, V.; Davis, L.; Mahesh, M.; Barry, N. P.; Chin, J. W. *J. Am. Chem. Soc.* **2015**, *137*, 4602.
- (36) Schmied, W. H.; Elsasser, S. J.; Uttamapinant, C.; Chin, J. W. *J. Am. Chem. Soc.* **2014**, *136*, 15577.