

Preparing (Multi)Fluoroarenes as Building Blocks for Synthesis: Nickel-Catalyzed Borylation of Polyfluoroarenes via C–F Bond Cleavage

Jing Zhou,^{†,‡} Maximilian W. Kuntze-Fechner,[‡] Rüdiger Bertermann,[‡] Ursula S. D. Paul,[‡] Johannes H. J. Berthel,[‡] Alexandra Friedrich,[‡] Zhenting Du,[†] Todd B. Marder,^{*,‡} and Udo Radius^{*,‡}

[†]College of Science, Northwest A&F University, Yangling, Shaanxi 712100, P. R. China

[‡]Institut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg, Am Hubland, 97074 Würzburg, Germany


S Supporting Information

ABSTRACT: The $[\text{Ni}(\text{IMes})_2]$ -catalyzed transformation of fluoroarenes into arylboronic acid pinacol esters via C–F bond activation and transmetalation with bis-(pinacolato)diboron (B_2pin_2) is reported. Various partially fluorinated arenes with different degrees of fluorination were converted into their corresponding boronate esters.

Arylboronic acid esters are versatile reagents in organic synthesis, especially in the widely employed Suzuki–Miyaura cross-coupling reaction.¹ Transition-metal-catalyzed direct C–H borylation of arenes² and borylation of aryl halides³ have emerged as one of the most important mild and attractive routes for the synthesis of aryl boronates in recent years. Fluoroaromatics play an important role in pharmaceuticals, agrochemicals, and materials,⁴ and there is currently much research aimed at the development of new ways to introduce fluorine or fluorinated building blocks into organic molecules.^{4,5} A smart way to achieve this goal would employ fluoroaromatic boronic acids or boronate esters. The conversion of fluoroaromatics into arylboronic esters via C–F bond activation, however, is relatively unexplored and was restricted to noble metal catalysts until recently. Smith et al. observed small amounts of C–F bond borylation products (~5%) in the iridium-catalyzed borylation of pentafluorobenzene and in the rhodium-catalyzed borylation of 1,3,5-trifluorobenzene.⁶ C–F bond borylation promoted by $[\text{Rh}(\text{SiPh}_3)(\text{PMe}_3)_3]$ was reported by Marder, Perutz et al.⁷ in stoichiometric reactions, and Braun et al. developed the Rh(I)-catalyzed borylation of pentafluoropyridine, hexafluoropropene and other fluorinated aromatics.⁸ Furthermore, Zhang et al. reported that commercially available $[\text{Rh}(\text{cod})_2]\text{BF}_4$ catalyzes *ortho*-selective C–F bond borylation of polyfluoroarenes with B_2pin_2 .⁹ All of the above protocols employ noble metal catalysts.

Although much work has been done in the field of nickel-mediated C–F bond activation,^{5,10} applications of nickel phosphine complexes in catalytic C–F bond borylation appeared just recently. Thus, Martin et al.¹¹ and Niwa and Hosoya et al.¹² reported in 2015 the borylation of monofluoroarenes using *in situ*-generated nickel phosphine complexes from the reaction of $[\text{Ni}(\text{COD})_2]$ (COD = 1,5-cyclooctadiene) and PCy_3 (>4 equiv) (Scheme 1). Martin et al. chose bis(neopentylglycolato)diboron (B_2neop_2) as the boron

Scheme 1. Nickel-Catalyzed Borylation of Fluoroaromatics



source and NaOPh as a base to promote the reaction in THF at 110 °C, whereas Niwa et al. reported borylation using bis(pinacolato)diboron (B_2pin_2) as the boron source, CsF as an additive, and a copper source as a mediator in toluene at 80 °C for 24 h. However, borylation with the $[\text{Ni}(\text{COD})_2]/\text{PCy}_3$ system was only reported for monofluoroarenes and the reactions strongly depend on the conditions employed. Martin et al. demonstrated that the use of other nickel sources or coligands (e.g., using an NHC precursor or PPh_2 instead of PCy_3) leads to a significant drop in yield and a change of the base (NaO^tBu , HCO_2Na , CsF) or the use of B_2pin_2 as a boron source essentially kills the catalytic process. The results presented by Niwa et al. on the $[\text{Ni}(\text{COD})_2]/\text{Cu}$ halide/ PCy_3 -mediated borylation of monofluoroarenes similarly reveal a strong dependence on the copper source (CuI works, whereas other copper sources do not) and the base employed (CsF works whereas other additives investigated do not).

We are currently developing convenient methodologies to generate and use suitable, partially fluorinated organic precursors. One strategy we follow employs (i) C–F borylation of polyfluoroaromatics and (ii) use of the resulting fluoroaryl boronic ester in subsequent Suzuki–Miyaura coupling reactions. The reports by Martin et al.¹¹ and Niwa and Hosoya et al.¹² prompted us to disclose our results on the NHC nickel-catalyzed borylation of polyfluorinated aromatics (Scheme 1).

We have demonstrated earlier that $[\text{Ni}_2(\text{Pr}_2\text{Im})_4(\text{COD})]$ ($\text{Pr}_2\text{Im} = 1,3$ -diisopropyl-imidazolin-2-ylidene)^{10a} reacts with fluorinated arenes with high chemo- and regioselectivity and catalyzes C–F bond transformations.¹⁰ Thus, we envisioned this complex as a precatalyst for C–F bond borylation, but our

Received: March 3, 2016

Published: April 12, 2016

initial experiments revealed that the nickel compound decomposes in the presence of B_2pin_2 or B_2cat_2 . As an alternative, the complex $[Ni(IMes)_2]$ with the sterically more demanding NHC IMes (IMes = 1,3-dimesitylimidazol-2-ylidene), which is sufficiently stable in the presence of diboron esters,¹³ was used in the current study.

We started our investigation using 1,2,3-trifluorobenzene **1a** (1 equiv) as the model substrate, B_2pin_2 (1 equiv) as the boron source, CsF (1 equiv) as an additive, $[Ni(IMes)_2]$ (10 mol %) as the catalyst, and THF as the solvent of choice. A C–F borylation product was obtained from this reaction in 39% yield (judged by ^{19}F NMR spectroscopy; Table 1, entry 1), which


Table 1. Optimization of the Borylation of 1,2,3- $C_6F_3H_3$

	1a/ B_2pin_2	base	solvent	yield (%) ^{a,b}
1	1.1:1	CsF (1 equiv)	THF	39
2	1.1:1	CsF (1 equiv)	C_9H_{12}	50
3	1.1:1	CsF (1 equiv)	C_6H_{12}	50
4	3:1	CsF (1 equiv)	C_9H_{12}	75
5	1.1:1	NMe_4F (1 equiv)	C_9H_{12}	62
6	1.1:1	NMe_4F (0.5 equiv)	C_9H_{12}	76
7	1.1:1	NMe_4F (0.5 equiv)	C_6H_{12}	79

^aReaction conditions: 29.0 mg (0.22 mmol) 1,2,3- $C_6F_3H_3$, **1a**; $[Ni(IMes)_2]$ (10 mol %), solvent (1 mL), 80 °C, 15 h. The yields were determined by *in situ* ^{19}F NMR spectroscopy. ^bTrace amounts of isomeric C–F and C–H borylation products were detected by GC-MS. C_9H_{12} = mesitylene; C_6H_{12} = methylcyclopentane. For more details see the SI.

was identified as compound **2a** (Scheme 2), i.e., the product from a defluoroborylation at the 1-position. In addition to **2a**, the regioisomer **2a'** (borylation at the 2-position), traces of the hydrodefluorination/borylation products **2g** and **2h**, and a C–

Scheme 2. Borylation Products and Side Products



H borylation product were detected by GC-MS analysis. Interestingly, trace amounts of the diborylation compound **3a**, presumably formed by subsequent C–H and C–F borylation, were detected as well. Note that the preference of NHC nickel insertion into the C–F bond at the 1-position of 1,2,3- $C_6F_3H_3$ was observed before.^{10d} With this promising first result, different solvents, bases, and different stoichiometric ratios of fluorobenzenes and B_2pin_2 were screened using standardized conditions (80 °C, 15 h) to assess the scope and limitations of this reaction (Table 1 and Tables S1–S4). First of all, our studies clearly reveal that the nature of the solvent used is very important. For hexane, mesitylene, or methylcyclopentane instead of THF, the yields of **2a** increase to 45% (hexane) and 50% (mesitylene or methylcyclopentane), respectively (Table S1 and Table 1, entries 2, 3), whereas acetonitrile shuts the reaction down completely (Table S1). Methyl *t*-butyl ether works similarly well compared to THF; in this case, however, the C–F/C–H diborylation product **3a** and the alkylation product **4a** (Scheme 2) were detected in trace amounts (Table S1). The latter compound implies C–O bond cleavage of MTBE under catalytic conditions.¹⁴ In general, yields and turnover rates are low, but some improvement with respect to the consumption of B_2pin_2 was achieved if the amount of fluoroarene was increased to 3 equiv (75% **2a**; Table S2, Table 1, entry 4). An increase in the amount of B_2pin_2 (and CsF) to 2 equiv leads to the formation of a mixture of monoborylated **2a** and bisborylated **3a**. A screening of different additives further reveals that the choice of the base also dramatically affects the reaction and that the presence of fluoride is crucial and beneficial (Table S3). Interestingly, the addition of 0.5 equiv of water-free NMe_4F instead of CsF provided a superior yield, especially, if methylcyclopentane was used as the solvent (Table S4, Table 1, entries 5–7). However, considering the fact that water-free NMe_4F is relatively expensive,¹⁵ we also provide the conditions and results for reactions using much cheaper and commercially available fluoride sources such as CsF, KF, or NBu_4F (Tables S1–S4). CsF and KF were moderately effective and provide the product in 50% yield, while the use of water-containing TBAF shuts the reaction down completely. With water-free NMe_4F in hand, the conditions as given in Table 1, entry 7 are the best for the formation of **2a** (1.1 equiv $C_6F_3H_3$, 1 equiv B_2pin_2 , 10 mol % $[Ni(IMes)_2]$, 0.5 equiv NMe_4F , methylcyclopentane, 80 °C, 15 h).

Using these optimized conditions, we then explored the scope of the nickel-catalyzed C–F bond borylation using different fluoroarenes. As shown in Table 2, the borylation of fluoroarenes containing different degrees of fluorination was achieved in moderate to good yields. For example, using 1,3,5- $C_6F_3H_3$ (**1b**), monoborylated **2b** was obtained in 93% yield (Table 2, entry 2). Borylations of 1,2,3-trifluorobenzene (**1a**) and 1,2,3,5- $C_6F_4H_2$ (**1d**) were also accomplished with good selectivity and yield (Table 2, entries 1, 4). For other substrates such as 1,2,4- $C_6F_3H_3$ (**1c**), 1,2,4,5- $C_6F_4H_2$ (**1e**), and C_6F_5H (**1f**), the temperature had to be raised to 100 °C for successful borylation (Table 2, entries 3, 5, 6). For 1,2,4- $C_6F_3H_3$ (**1c**), borylation of the 2-position (**2c**) prevails, and only minor amounts of the isomers from borylation of the 1-(**2c''**) or of the 4-positions (**2c'**) (ratio ~10:1:1, judged from ^{19}F NMR spectroscopy) were formed. For 1,2,4,5- $C_6F_4H_2$ (**1e**), diborylation to give **2e'** was observed when using the standard reaction conditions. However, the formation of monoborylation product (**2e**) and diborylation product (**2e'**) as the main product can be controlled by the ratio of fluoroaromatic and

Table 2. Scope of the Ni-Catalyzed C–F Borylation

Ar ^F -F		[Ni(IMes) ₂] (10 mol %)		Ar ^F -Bpin	
		B ₂ pin ₂ (1 eq.), solvent			
		NMe ₄ F (0.5 eq.)/CsF (1 eq.)			
substrate		product and yield			
1	1,2,3-C ₆ F ₃ H ₃	1-Bpin-2,3-C ₆ F ₂ H ₃ (2a)	79% (38%) ^b		
2	1,3,5-C ₆ F ₃ H ₃	1-Bpin-3,5-C ₆ F ₂ H ₃ (2b)	93% (75%) ^b		
3	1,2,4-C ₆ F ₃ H ₃	1-Bpin-2,4-C ₆ F ₂ H ₃ (2c)	77% (60%) ^c	2c:2c':2c'' = 10:1:1 ^c	
4	1,2,3,5-C ₆ F ₄ H ₂	1-Bpin-2,3,5-C ₆ F ₃ H ₂ (2d)	99% (71%) ^b	2d:2d' = 10:1	
5	1,2,4,5-C ₆ F ₄ H ₂	1-Bpin-2,4,5-C ₆ F ₃ H ₂ (2e)	70% (65%) ^d	1,4-bisBpin-2,5-C ₆ F ₂ H ₂ (2e')	85% ^{e,f} (73%) ^e
6	C ₆ F ₅ H	1-Bpin-2,3,4,5-C ₆ F ₄ H (2f)	85% ^f (80%) ^{b,f}		
7	1,2-C ₆ F ₂ H ₄	1-Bpin-2-C ₆ FH ₄ (2g)	48% ^g (30%) ^g		
8	1,3-C ₆ F ₂ H ₄	1-Bpin-3-C ₆ FH ₄ (2h)	50% ^g (35%) ^g		
9	C ₆ FH ₅	C ₆ H ₅ Bpin (2i)	20% ^{g,h}		

^aReaction conditions (unless specified otherwise): B₂pin₂ (0.2 mmol), fluoroarene (0.22 mmol), [Ni(IMes)₂] (10 mol %), NMe₄F (0.1 mmol), methylcyclopentane (1 mL), 80 °C, 15 h. Yields determined by ¹⁹F NMR spectroscopy based on B₂pin₂. ^bReaction conditions: B₂pin₂ (2 mmol), fluoroarene (2.2 mmol), [Ni(IMes)₂] (10 mol %), CsF (2 mmol), methylcyclopentane (10 mL), 80 °C, 15 h. Isolated yields. ^cMethylcyclohexane, 100 °C. ^d2e (4 equiv), methylcyclohexane, 100 °C. ^eB₂pin₂ (2 equiv), fluoroarene (1 equiv), CsF (2 equiv), methylcyclohexane (20 mL), 100 °C, 15 h. ^f[Ni(IPr)₂] (10 mol %) instead of [Ni(IMes)₂]. ^gFluoroarene (2 equiv, 0.4 mmol). ^hYield was determined by GC-MS using C₁₂H₂₆ as the internal standard. For more details see the SI.


diboron reagent used. One equiv of 1,2,4,5-C₆F₄H₂, 2 equiv B₂pin₂, and CsF lead to diborylation product **2e'** in 73% yield. If the amount of 1,2,4,5-C₆F₄H₂ was increased to 4 equiv, the monoborylation product **2e** was obtained in good yield with only small amounts of **2e'**. Compounds **2e** and **2e'** can be separated easily by column chromatography (see SI). The borylation of C₆F₅H occurs not at the most nucleophilic 3-position, as probably expected, but at the position *ortho* to the hydrogen substituent. However, selective *ortho*-C–F bond activation of C₆F₅H at nickel was observed before.¹⁶ We were unsuccessful in improving the reaction with [Ni(IMes)₂] from being essentially stoichiometric in nickel (12% yield at 100 °C). However, we found that the use of [Ni(IPr)₂], bearing the sterically more demanding NHC 1,3-bis(2,6-diisopropylphenyl)imidazolin-2-ylidene IPr, as a precatalyst is beneficial to this reaction, and formation of the borylation product was observed in more than 80% yield. For 1,2-C₆F₂H₄ (**1g**) and 1,3-C₆F₂H₄ (**1h**), C–F borylation products were formed in somewhat lower yields, with trace amounts of C–H bond borylation observed by GC-MS (Table 2, entries 7, 8). Borylation of monofluorobenzene leads to PhBpin **2i** in only 20% yield (Table 2, entry 9). Moreover, we observe no reaction or only traces of products for the borylation of perfluorinated substrates such as hexafluorobenzene or octafluorotoluene. We also provide yields of the isolated product using CsF as a base in Table 2. All products have been characterized by multinuclear NMR spectroscopy and high-resolution mass spectroscopy and, for **3a**, **2e'** and **2f'**, by X-ray diffraction (see SI).

In previous work we have shown that NHC nickel complexes readily insert into C–F bonds of polyfluorinated aromatics.¹⁰ Using [Ni(IMes)₂], C–F bond activation also occurs cleanly. The reaction of 1,2,3,5-C₆F₄H₂ (**1d**) with [Ni(IMes)₂] affords the nickel fluoride complex *trans*-[Ni(IMes)₂(F)(Ar^F)] (**5**).

This reaction is quantitative according to ¹H and ¹⁹F NMR spectroscopy. Complex **5** has been characterized by proton and fluorine NMR spectroscopy, mass spectrometry, elemental analysis, and X-ray diffraction. Most significantly, the Ni–F ligand of **5** gives rise to a resonance at –344.5 ppm in the ¹⁹F NMR spectrum.

The reaction of [Ni(IMes)₂] with 1,2,3,5-C₆F₄H₂ (**1d**) in the presence of B₂pin₂ and NMe₄F leads to the initial formation of the C–F bond activation product **5** (see SI). The boryl transfer reagent should be either B₂pin₂ or its fluoride adduct, i.e., [NMe₄][B₂pin₂F] **7**.¹⁷ The reaction of **5** with an excess of B₂pin₂ and NMe₄F using catalytic conditions leads to borylated fluoroarene, complex **5** and [NMe₄][F₂Bpin]. We favor at present a mechanism as outlined in Scheme 3. The reaction of

Scheme 3. Proposed Mechanism for the NHC-Nickel-Catalyzed Borylation of Fluoroarenes



[Ni(IMes)₂] **I** with the fluoroarene leads in the first step to oxidative addition of the C–F bond with formation of *trans*-[Ni(IMes)₂(F)(Ar^F)] **II**. Although an initial “kinetic” C–H bond activation step^{3d,e} would explain the regioselectivity of the borylation reaction (borylation occurs at a site adjacent to an hydrogen substituent), we can not observe such a species even at low temperatures (see Figures S68–S70). The complex *trans*-[Ni(IMes)₂(F)(Ar^F)] **II** reacts then with [NMe₄][B₂pin₂F] by boryl transfer to give *trans*-[Ni(IMes)₂(Bpin)(Ar^F)] **III** and [NMe₄][F₂Bpin]. A final reductive elimination step (e.g., after ligand elimination from a three coordinate species as indicated in Scheme 3 or after *cis/trans* isomerization of the boryl complex) delivers the borylated fluoroaromatic Ar^F-Bpin and regenerates [Ni(IMes)₂] **I**. The borylation of the nickel fluoride complex seems to be the rate-limiting process in the sequence, as we (i) observe larger quantities of the nickel fluoride as the resting state in the reaction mixtures under catalytic conditions (see SI) and (ii) failed to isolate complexes *trans*-[Ni(IMes)₂(Bpin)(Ar^F)] so far from the stoichiometric reaction of *trans*-[Ni(IMes)₂(F)(Ar^F)] with B₂pin₂/NMe₄F.

In summary, we have developed an efficient procedure for the C–F borylation of fluoroaromatic compounds using [Ni(IMes)₂] ([Ni(IPr)₂] for pentafluorobenzene) as a catalyst, NMe₄F or CsF as a fluoride source, and B₂pin₂ as the boron source. In combination with the wide versatility of borylarene transformations, this method will enable the variation of C–F bonds in polyfluoroarenes in the future, e.g., replacing selected fluorine substituents with other groups such as other halides, amides, carboxylate, or ether functionalities. On the other hand, typical C–C bond formation steps will allow the introduction of partially fluorinated arenes into larger organic molecules.

Detailed investigations concerning the applications and mechanistic details of this reaction are in progress.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b02337.

Experimental details and data (PDF)

Crystallographic data (CIF)

■ AUTHOR INFORMATION

Corresponding Authors

*u.radius@uni-wuerzburg.de

*todd.marder@uni-wuerzburg.de

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by funds from University of Würzburg and the DFG (Ra720/12-1). We are grateful to the China Scholarship Council for providing a scholarship to Jing Zhou. We thank Prof. Dr. Maik Finze and Mathias Häring for a donation of anhydrous NMe_4F and AllyChem Co. Ltd. for a gift of B_2pin_2 . Dedicated to Prof. em. Dr. Hansgeorg Schnöckel on the occasion of his 75th birthday.

■ REFERENCES

- (1) (a) *Boronic acids: preparation, applications in organic synthesis and medicine*; Hall, D. G., Ed.; Wiley-VCH: Weinheim, Germany, 2006. (b) Suzuki, A. *Angew. Chem., Int. Ed.* **2011**, *50*, 6722.
- (2) (a) Mkhaliid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F. *Chem. Rev.* **2010**, *110*, 890. (b) Hartwig, J. F. *Chem. Soc. Rev.* **2011**, *40*, 1992. (c) Hartwig, J. F. *Acc. Chem. Res.* **2012**, *45*, 864. (d) Ros, A.; Fernandez, R.; Lassaletta, J. M. *Chem. Soc. Rev.* **2014**, *43*, 3229.
- (3) (a) Ishiyama, T.; Murata, M.; Miyaura, N. *J. Org. Chem.* **1995**, *60*, 7508. (b) Murata, M. *Heterocycles* **2012**, *85*, 1795. (c) Chow, W. K.; Yuen, O. Y.; Choy, P. Y.; So, C. M.; Lau, C. P.; Wong, W. T.; Kwong, F. Y. *RSC Adv.* **2013**, *3*, 12518. (d) Kleeberg, C.; Dang, L.; Lin, Z.; Marder, T. B. *Angew. Chem., Int. Ed.* **2009**, *48*, 5350. (e) Bose, S. K.; Marder, T. B. *Org. Lett.* **2014**, *16*, 4562. (f) Bose, S. K.; Deisenberger, A.; Eichhorn, A.; Steel, P. G.; Lin, Z.; Marder, T. B. *Angew. Chem. Int. Ed.* **2015**, *54*, 11843–11847.
- (4) (a) Dolbier, W. R. *J. Fluorine Chem.* **2005**, *126*, 157. (b) Uneyama, K. *Organofluorine Chemistry*; Wiley-Blackwell: Oxford, U.K., 2006. (c) Wang, J.; Sanchez-Rosello, M.; Acena, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. *Chem. Rev.* **2014**, *114*, 2432.
- (5) (a) Doherty, N. M.; Hoffmann, N. W. *Chem. Rev.* **1991**, *91*, 553. (b) Amii, H.; Uneyama, K. *Chem. Rev.* **2009**, *109*, 2119. (c) Sun, A. D.; Love, J. A. *Dalton Trans.* **2010**, *39*, 10362. (d) Clot, E.; Eisenstein, O.; Jasim, N.; Macgregor, S. A.; McGrady, J. E.; Perutz, R. N. *Acc. Chem. Res.* **2011**, *44*, 333. (e) Johnson, S. A.; Hatnean, J. A.; Doster, M. E. *Progress in Inorganic Chemistry*; John Wiley & Sons, Inc.: Hoboken, NJ, 2012; Vol 57, p 255. (f) Kuehnel, M. F.; Lentz, D.; Braun, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 3328. (g) Keyes, L.; Love, J. A. *Aromatic C–F Activation: Converting Fluoroarenes to Useful Building Blocks*. In *C–H and C–X Bond Functionalization*; Ribas, X., Ed.; RSC: Cambridge, U.K., 2013. (h) Weaver, J.; Senaweera, S. *Tetrahedron* **2014**, *70*, 7413. (i) Ahrens, T.; Kohlmann, J.; Ahrens, M.; Braun, T. *Chem. Rev.* **2015**, *115*, 931.
- (6) Cho, J.-Y.; Iverson, C. N.; Smith, M. R. *J. Am. Chem. Soc.* **2000**, *122*, 12868.
- (7) Lindup, R. J.; Marder, T. B.; Perutz, R. N.; Whitwood, A. C. *Chem. Commun.* **2007**, 3664.

- (8) (a) Braun, T.; Ahijado Salomon, M.; Altenhöner, K.; Teltewskoi, M.; Hinze, S. *Angew. Chem., Int. Ed.* **2009**, *48*, 1818. (b) Teltewskoi, M.; Panetier, J. A.; Macgregor, S. A.; Braun, T. *Angew. Chem., Int. Ed.* **2010**, *49*, 3947. (c) Kalläne, S. I.; Teltewskoi, M.; Braun, T.; Braun, B. *Organometallics* **2015**, *34*, 1156.
- (9) Guo, W. H.; Min, Q. Q.; Gu, J. W.; Zhang, X. *Angew. Chem.* **2015**, *127*, 9203.
- (10) (a) Schaub, T.; Radius, U. *Chem. - Eur. J.* **2005**, *11*, 5024. (b) Schaub, T.; Backes, M.; Radius, U. *J. Am. Chem. Soc.* **2006**, *128*, 15964. (c) Schaub, T.; Backes, M.; Radius, U. *Eur. J. Inorg. Chem.* **2008**, *2008*, 2680. (d) Schaub, T.; Fischer, P.; Steffen, A.; Braun, T.; Radius, U.; Mix, A. *J. Am. Chem. Soc.* **2008**, *130*, 9304. (e) Schaub, T.; Fischer, P.; Meins, T.; Radius, U. *Eur. J. Inorg. Chem.* **2011**, *2011*, 3122. (f) Zell, T.; Feierabend, M.; Halfter, B.; Radius, U. *J. Organomet. Chem.* **2011**, *696*, 1380. (g) Fischer, P.; Götz, K.; Eichhorn, A.; Radius, U. *Organometallics* **2012**, *31*, 1374.
- (11) Liu, X.-W.; Echavarren, J.; Zarate, C.; Martin, R. *J. Am. Chem. Soc.* **2015**, *137*, 12470.
- (12) Niwa, T.; Ochiai, H.; Watanabe, Y.; Hosoya, T. *J. Am. Chem. Soc.* **2015**, *137*, 14313.
- (13) Pietsch, S.; Paul, U.; Cade, I. A.; Ingleson, M. J.; Radius, U.; Marder, T. B. *Chem. - Eur. J.* **2015**, *21*, 9018.
- (14) C–O bond cleavage of ethers by NHC nickel complexes is a known process and may be used for the catalytic transformation of ethers, see for example: (a) Sergeev, A. G.; Hartwig, J. F. *Science* **2011**, *332*, 439. (b) Tollefson, E. J.; Hanna, L. E.; Jarvo, E. R. *Acc. Chem. Res.* **2015**, *48*, 2344. (c) Liu, C.; Peterson, C.; Wilson, A. K. *J. Phys. Chem. A* **2013**, *117*, 5140.
- (15) For example, Aldrich 459315:5g ~300 €. (16) Johnson, S. A.; Taylor, E. T.; Cruise, S. J. *Organometallics* **2009**, *28*, 3842.
- (17) The $[\text{NMe}_4][\text{B}_2\text{pin}_2\text{F}]$ salt was not detected in the reaction mixtures we have investigated in C_6D_6 and methylcyclopentane. However, it is known that it is only sparingly soluble in these solvents, but soluble in acetonitrile and in hot THF. We recently reported the preparation, isolation, and utilization of $[\text{NMe}_4][\text{B}_2\text{pin}_2\text{F}]$ for boryl transfer: Pietsch, S.; Neeve, E. C.; Apperley, D. C.; Bertermann, R.; Mo, F.; Qiu, D.; Cheung, M. S.; Dang, L.; Wang, J.; Radius, U.; Lin, Z.; Kleeberg, C.; Marder, T. B. *Chem. - Eur. J.* **2015**, *21*, 7082.