


Hydrogen Bonding to Hexafluoroisopropanol Controls the Oxidative Strength of Hypervalent Iodine Reagents

Ignacio Colomer,[†] Christopher Batchelor-McAuley,[‡] Barbara Odell,[†] Timothy J. Donohoe,^{*,†} and Richard G. Compton^{*,‡}

[†]Department of Chemistry, Chemistry Research Laboratory, University of Oxford, Mansfield Road, Oxford OX1 3TA, United Kingdom

[‡]Department of Chemistry, Physical and Theoretical Chemistry Laboratory, University of Oxford, South Parks Road, Oxford OX1 3QZ, United Kingdom

S Supporting Information




ABSTRACT: Hexafluoroisopropan-2-ol (HFIP) has been found to be an unusually beneficial solvent for undertaking hypervalent iodine-initiated [2+2] cycloaddition of styrenes. For the initiator phenyliodine(III) diacetate (PIDA), voltammetric data demonstrate that the enhanced reactivity in HFIP is due to its greater oxidizing abilities in this fluorinated solvent such that in HFIP the reactivity of PIDA is comparable if not superior to its fluorinated analog phenyliodine(III) bis(trifluoroacetate). These results contrast with the often reported view that the role of the fluoroalcohol is to stabilize a radical cation formed by single electron transfer. Moreover, combined NMR and HRMS results reveal the formation of a strong H-bonded adduct between the solvent and oxidizing reagent which is the physical origin of the observed altered synthetic reactivity.

1. INTRODUCTION

Hypervalent iodine reagents exhibit diverse reactivity and find widespread use in organic synthesis.¹ Much of their chemistry is analogous to that found for transition metal complexes. As an extension of these properties, under suitable conditions they may be utilized as less toxic and safer replacements for heavy metal oxidants. Surveying the literature reveals the common and regular usage of fluorinated solvents in combination with this class of reagent.² Specifically in the case of single electron transfer (SET) reactions the requirement and benefit of the use of solvents such as hexafluoroisopropanol (HFIP) and trifluoroethanol are often attributed to their ability to stabilize radical cations.³ As compared to many other solvents the properties of HFIP are extreme, it is highly polar, strongly hydrogen-bond donating and very weakly nucleophilic. The use of HFIP as a solvent or additive is known to kinetically enhance or alter the selectivity of a number of synthetically important reactions. Most notable is its use in combination with hydrogen peroxide for both epoxidations and Baeyer–Villiger oxidations,⁴ as well as a H-bond donor to epoxides or carbonyl compounds, therefore increasing their electrophilicity.⁵ It has been recently reported that hypervalent iodine reagents such as phenyliodine(III) diacetate (PIDA, **2**, Scheme 1) can initiate the dimerization of styrenes (see *trans*-anethole, **1**, Scheme 1) in hexafluoroisopropanol (HFIP, **3**, Scheme 1), leading to the synthesis of cyclobutanes **4**.⁶ Approaching the stereoselective

Scheme 1. Hypervalent Iodine-Initiated Homo- and Cross-Dimerization of Styrenes in Hexafluoroisopropanol



synthesis of substituted cyclobutanes is of great interest and represents a challenging topic in organic chemistry.⁷ Despite the presence of the cyclobutane moiety in different natural products with antifungal, anticancer and many other biological activities,⁸ the absence of reliable, efficient and benign methods for its synthesis and incorporation into complex molecules has hampered its widespread use.

This synthetic [2+2] cycloaddition is used in the present work as an exemplar reaction by which the reactivity of the hypervalent iodine(III) reagents may be explored. The formation of the cyclobutane is considered to initially proceed via single electron oxidation of the alkene **1**, to form a radical cation intermediate that can react with another molecule of **1**

Received: April 20, 2016

Published: July 5, 2016

(homodimerization) or with a different alkene (hetero- or cross dimerization). HFIP was found to be a unique solvent in which the hypervalent iodine reagents can act as single electron oxidants in the cyclobutane forming process.⁹ Importantly, analogous [2+2] cycloadditions have been seen in other organic solvents (typically CH₂Cl₂ and acetonitrile (ACN)) when initiated using a triaryl amine radical cation, anodic oxidation or photoredox catalysis¹⁰ (although we found that the sole use of PIDA in both dichloromethane or acetonitrile did not lead to the formation of the cyclobutane product). In light of the above-mentioned results, this work provides a systematic investigation of the redox properties of the alkene and the oxidant in both ACN and HFIP. The voltammetric oxidation of a series of electron-rich styrenes demonstrates that the fluorinated solvent HFIP only *relatively* minimally alters the oxidative response of the alkene. Hence, the influence of the fluoroalcohol is re-assessed, and the significant role hydrogen bonding¹¹ with the fluorinated solvents has upon controlling the reactivity of the hypervalent iodine reagents is illuminated.

2. RESULTS AND DISCUSSION

This work first studies the altered reactivity of PIDA toward *trans*-anethole. In HFIP, as opposed to ACN, this reaction is found to proceed and leads to the formation of the cyclobutane product. Having evidenced the physical origin of the altered reactivity the article continues by considering the reactivity of related substituted styrenes and iodines.

2.1. Electrochemical Behavior of *trans*-Anethole and Phenyliodine(III) Diacetate. The oxidation of *trans*-anethole **1** was recorded voltammetrically at a glassy carbon electrode in both HFIP (Figure 1 A) and acetonitrile (Supporting




Figure 1. Voltammetric response of (A) *trans*-anethole (**1**, oxidation) and (B) phenyliodine(III) diacetate (**2**, reduction) in HFIP (0.1 M Bu₄NPF₆) recorded as a function of scan rate (0.025–0.4 V s⁻¹) at a glassy carbon electrode (radius = 1.5 mm). In (A) the reversible voltammetric response of ferrocene is also shown; the ferrocene was used as an internal redox reference potential. In (B) a silver/silver hexafluorophosphate reference electrode ($E_{\text{ferrocene}} = -0.988$ V vs Ag/Ag⁺) was used due to reaction between the ferrocene and PIDA. The horizontal arrow indicates voltammetric starting potential and initial scan direction.

Information (SI), section 2.1). In both solvents a well-defined irreversible oxidative feature was observed (at +0.85 and +1.02 V vs ferrocene for HFIP and ACN, respectively). In acetonitrile this electrochemical oxidation is known to correspond to the one electron oxidation of the *trans*-anethole species followed by an irreversible and fast dimerization ($\log k$ (M s⁻¹) = 8.6 ± 0.4) of the formed radical cation.¹² In HFIP the oxidative voltammetric response was confirmed to correspond to the one electron oxidation of the *trans*-anethole. This assessment of the numbers of electrons transferred was achieved via recording of the voltammetric response at an ultramicroelectrode.

Comparison of the steady-state and linear diffusive response enables direct assessment of the number of electrons transferred during the course of the electrochemical oxidation (SI, section 2).

The oxidative peak potential of the *trans*-anethole at a macro electrode in HFIP and ACN (Figure 1A and SI, section 2.1) reflects both the thermodynamics and kinetics of the chemically irreversible redox process. By using ferrocene as an internal redox marker the potential difference between the oxidation of *trans*-anethole in the two solvents is measured to be -0.18 V, thus evidencing the oxidation of the alkene to be easier in HFIP. Relatively weak stabilizing effects of HFIP on extensively delocalized radical cations have been previously reported. For example the formal potential for the one-electron oxidation of durene (1,2,4,5-tetramethylbenzene) in HFIP was shifted by -0.14 V versus that recorded in dichloromethane.¹³ Hence, the observed shift of -0.18 V between the two solvents is consistent with the greater radical cation stabilizing abilities of HFIP over ACN.

The use of electrochemical techniques for studying cathodic processes in HFIP is limited; this situation has plausibly arisen due to the reduction processes of interest being obscured by concomitant hydrogen evolution. However, the use of non-metallic electrodes such as boron-doped diamond or glassy carbon can effectively extend the potential window of study due to their associated sluggish electron transfer kinetics toward proton reduction.¹⁴ The reduction of PIDA was studied in both HFIP (Figure 1B) and ACN (SI, section 2.3) using a glassy carbon macro-electrode. In HFIP the presence of both PIDA and ferrocene in solution led to rapid reaction and decomposition of the PIDA as evidenced by the change in the solution color. This is in contrast to solutions of PIDA and ferrocene in ACN which were found to be stable. Consequently, the voltammetry of PIDA in HFIP was recorded against a silver/silver hexafluorophosphate reference electrode. The ferrocene redox potential was determined to be -0.988 V vs Ag/Ag⁺ in HFIP (see SI, section 2.5), this measurement enabled direct comparison of the voltammetric data across the two solvents.

In both ACN and HFIP, the reduction of PIDA exhibits two distinct voltammetric reduction waves. In both cases the analysis of the first reduction wave indicated the transfer of up to three electrons to the PIDA (SI, section 2) over the course of the voltammetric reduction. On the basis of the molecular structure it may be anticipated for the hypervalent iodine to undergo two two-electron transfers with the formation of iodobenzene after the first reduction wave. However, importantly, for the voltammetric reduction in both solvents the transfer coefficients were measured to be 0.3 and 0.4 in ACN and HFIP respectively, indicating the first electron transfer to be rate determining in both cases. Most notably the reduction of PIDA occurs at a 0.85 V less negative potential (versus ferrocene) in HFIP as compared to ACN. The voltammetric peak potentials for both the oxidation ($E_{\text{p,a}}$) of *trans*-anethole and the reduction ($E_{\text{p,c}}$) of PIDA in both HFIP and ACN are summarized in Table 1. Also given in Table 1 are the differences in the reduction and oxidation peak potentials ($\Delta E_{\text{p-p}}$) in these two solvents. From these data, two points should be highlighted. First, in both solvents the voltammetric reduction of the iodine is at a significantly lower potential than the *trans*-anethole oxidation. This large difference in potentials likely indicates the electron transfer to PIDA from the *trans*-anethole is endergonic. This result is in agreement

Table 1. Voltammetric Peak Potentials (V) for Two Redox Species Measured versus Fc/Fc⁺, As Measured at 100 mV s⁻¹

	ACN	HFIP
<i>trans</i> -anethole (1), $E_{p,a}$	1.02	0.85
PIDA (2), $E_{p,c}$	-1.32	-0.47
ΔE_{p-p}	-2.35	-1.31

with previous work which has highlighted how for SET initiated cycloadditions the oxidant should be sufficiently weak so as to ensure only low concentrations of the radical cation are produced in situ avoiding dimerization and associated formation of the dication.¹⁵ Second, the difference between the voltammetric oxidation and reduction peak potentials is almost 1 V less in HFIP as compared to ACN! This change in the peak potentials predominantly reflects the significantly easier reduction of PIDA in HFIP. These results evidence the strong enhancement of the oxidizing abilities of PIDA in HFIP and yield insight into why the synthetic reaction is successful in HFIP but not ACN.

2.2. Characterizing the Altered Reactivity of PIDA in HFIP by MS and NMR. Although the voltammetric response reported above evidences a significant shift in the oxidizing abilities of the PIDA in HFIP, such techniques do not readily yield insight into the physical origin of this altered reactivity. Due to the irreversibility of the electrochemical process the shift in the voltammetric wave may reflect a change in the kinetics or the thermodynamics associated with first (rate-determining) electron transfer. In light of the importance of ligand exchange as a dominant reaction route for many iodine synthetic pathways one plausible explanation for the altered reactivity in HFIP is the substitution of the acetate groups by the solvent. Such a ligand exchange might be anticipated to be observed via high-resolution mass spectrometry (HRMS).

The HRMS of PIDA (2) was recorded and the pattern of ionization peaks found to be in agreement with the literature (see SI, section 3, for more details).¹⁶ The main species detected are the Na⁺ (344.9588) and K⁺ (360.9326) adducts, as well as [PhIOAcOIPh]⁺ (482.8936), [PhI]^{•+} (203.9428) and [PhIOAc]⁺ (262.9559). Similarly, the HRMS for HFIP (3) resulted in the observation of the anion derived from the loss of H⁺ [(CF₃)₂CHO]⁻ (166.9934). Importantly, the HRMS for a 1:1 mixture of PIDA:HFIP (5) gave the same mass spectra as obtained for PIDA (2), without the observation of any new formed species (within the sensitivity of the HRMS). This null result, although not conclusive, tentatively allows ligand exchange as a cause for the altered reactivity to be ruled out. This apparent lack of exchange with the HFIP is consistent with the low nucleophilicity of the solvent.

As an alternate explanation, the enhanced reactivity of PIDA in HFIP, may relate to the protic nature of the solvent, notably HFIP has a pK_a in water of 9.3 (as compared ~17 for isopropanol).^{15a} Nuclear magnetic resonance (NMR) was utilized to gain insight into this question. First, using ¹H NMR and ¹³C NMR (see SI, section 4, for full characterization and further details), the spectra of the species separately and in mixture were recorded. Figure 2 shows the ¹H NMR for PIDA (2), HFIP (3), and a 1:1 mixture of PIDA:HFIP (5) in CDCl₃.¹⁷ The ¹H NMR spectrum of the 1:1 complex of PIDA with HFIP (5, Figure 2c) is predominantly simply the sum of the two individual spectra. However, we observed a deshielded shift of the OH signal in HFIP (3, Figure 2b) from 2.88 to 5.88 ppm in the 1:1 adduct (5, Figure 2c), accompanied by




Figure 2. ¹H NMR of (a) PIDA (2), (b) HFIP (3), (c) 1:1 mixture of PIDA:HFIP (5), and (d) 1-D gradient NOE with $T_{mix} = 800$ ms for the 1:1 mixture of PIDA:HFIP (5) when irradiating 6-H₃ (Me), showing a plausible structure of an adduct, and the key NOE with arrows (see Supporting Information for a complete study of NOE and NOESY-2D). Red star indicates the position of the deshielded OH signal of interest.

broadening of the line width. Such a shift in the OH signal is consistent with the formation of a H-bonded adduct between the HFIP and PIDA.^{18,19} To further evidence the presence of a H-bonded adduct nuclear Overhauser effect (NOE) and diffusion-ordered spectroscopy (DOSY)²⁰ experiments were performed. NOE experiments provide evidence of dipolar spatial interactions between protons, nearby in space 2.5–4.5 Å. On irradiating the PIDA Me signal of the equimolar mixture of PIDA:HFIP (5) at 2.01 ppm (Figure 2d), significant NOEs were observed with the HFIP broad OH signal at 5.88 ppm as well as with the HFIP 1'-H at 4.40 ppm (see Supporting Information for a complete study of the NOE and NOESY-2D).

Using NMR, molecular association via H-bonding can be monitored via the differences in the diffusion coefficients (as partially reflecting molecular size) of the species involved. DOSY experiments allow quantitative study, via NMR, of the species diffusion coefficients as measured relative to an internal reference (here TMS, chosen for its non-interacting properties and low chemical shift).^{20b} The ratio of the diffusion between the species of study and the reference compound (D/D_{ref}) is relatively independent of solution viscosity. Consequently, any change in this ratio for the H-bonded adduct compared to the separate H-donor and/or the H-acceptor components may be attributed to the association of the components.

The results of the DOSY experiments (see SI, section 4, for the complete account of experiments) and the measured diffusion coefficients (Table 2) confirm the existence of a H-

Table 2. Diffusion Coefficients of PIDA (2), HFIP (3), and the 1:1 Mixture of PIDA:HFIP (5) in CDCl₃, As Measured via DOSY

compound	D ($\times 10^{-9}$ m ² s ⁻¹)	D/D^{TMS}
TMS	2.13	1.00
2	1.09	0.52
3	2.49	1.10
5	PIDA 1.09	0.51
5	HFIP 1.35	0.63

bonded species upon mixing both H-donor HFIP (3) and H-acceptor PIDA (2). While for PIDA (2) the change in the diffusion ratio (D/D_{TMS} from 0.52 to 0.51) upon mixing is small; however for HFIP (3) the variation in the diffusion ratio (D/D_{TMS} from 1.10 to 0.63) is considerable. This shows HFIP (3) is diffusing more slowly in the presence of PIDA (2). Therefore, both the NOE and DOSY results suggest the association of the HFIP with the PIDA, presumably via H-bonding. A similar set of NMR experiments were performed with a mixture of PIDA and 2-propanol, and these did not show a significant change in OH chemical shift upon mixing, nor significant changes in the diffusion ratios (DOSY), confirming the unique nature of HFIP. Although the PIDA has been shown to form a H-bonded adduct with the HFIP, a question remains as to whether this interaction serves to predominantly alter the thermodynamics or kinetics of the redox process.²¹

2.3. Exploring the Scope of Styrene Reactivity.

Although the use of HFIP as a solvent enables the homo- and heterodimerization of a number of styrenes, a notable structural motif in the successful reactions is the presence of a *p*-methoxy group. Figure 3 provides a survey of the voltammetric oxidation peak potentials for a variety of substituted *trans*- β -methylstyrenes in both ACN and HFIP (voltammograms from which the data were obtained are shown in the SI, section 2.6). As can be seen, there is a general





Figure 3. Voltammetric oxidative peak potentials (vs Fc/Fc⁺ measured at 100 mV s⁻¹) for a series of substituted *trans*- β -methylstyrenes measured in both ACN and HFIP. Dotted line indicates equi-potential between ACN and HFIP. Points in green represent styrenes able to undergo PIDA-initiated [2+2] cycloadditions in HFIP.

correlation between the peak potentials recorded in ACN and HFIP. For all but two of the tested alkenes, the voltammetric oxidation is found to occur at a lower potential in HFIP as compared to ACN, specifically the average shift in the oxidative peak potential was recorded to be ~ 140 mV, as measured against the ferrocene redox indicator. It is concluded that these generally lower peak potentials in HFIP likely reflect the weakly stabilizing nature of the solvent toward the formed cation.

None of the listed styrenes have been found to, in the presence of PIDA, react in ACN to form their associated cyclobutane products. However, the styrenes highlighted in green (Figure 3) are able found to undergo reaction in HFIP. Broadly the styrenes exhibiting oxidative peak potentials below $\sim +1.0$ V vs Fc/Fc⁺ in HFIP lead to formation of the [2+2] cycloaddition product. Given that, as has already been discussed, the reduction peak potential of PIDA is found to occur at -0.47 V (vs Fc/Fc⁺) in HFIP the initiation reaction leading to the formation of the aryl radical cation is likely endergonic in nature. Consequently, for the styrenes with the higher voltammetric oxidation potentials the cycloaddition reaction likely does not proceed due to the initiation being comparatively even less thermodynamically favorable for these more electron deficient styrenes. In general the presence of a *p*-methoxy group serves to very effectively activate the styrene, such that all of the most voltammetrically oxidizable styrenes studied contain this functionality. This is true to the extent that even the presence of an *o*-methoxy group (13) is found to be insufficiently activating to enable the synthetic reaction to proceed.

The general correlation between the synthetic and voltammetric data is notable. However, it should be commented that the synthetic reaction will be sensitive to not just the thermodynamics and kinetics of the initiation processes but is also predicated upon the efficiency of the following propagation reactions. Moreover, the position of the oxidative voltammetric wave is, as also previously discussed, sensitive to the involvement of possible further homogeneous chemical reactions.

2.4. Controlling the Reactivity of Hypervalent Iodines with HFIP. This section starts by highlighting the feasibility of using HFIP as an additive with PIDA. Next it looks to evidence the influence of HFIP on two alternate and commonly used iodine(III) reagents, specifically phenyliodine(III) bis(trifluoroacetate) (PIFA) and hydroxy(tosloxy)iodobenzene (HTIB, or Koser's reagent).²²



The NMR characterization presented in section 2.2 was performed in deuterated chloroform with 1:1 mixture of the HFIP and PIDA. The strong evidence for the formation of hydrogen bonded adduct leads to the question of the ability to use HFIP not as a solvent but as an additive. Indeed *trans*-anethole in the presence of 5.0 equiv of HFIP and 10 mol % of PIDA in deuterated chloroform is found to successfully result in the formation of the associated cyclobutane product albeit with a lower yield than obtained when utilizing HFIP as the solvent (35% yield). First, this observation further evidences adduct formation between the oxidant and the fluoroalcohol. Conversely, use of HFIP as an adduct in ACN does not lead

to significant product formation (<5% yield). Presumably, this arises due to the ACN solvent being a more available hydrogen bond acceptor than PIDA. NMR spectra of HFIP in d_3 -ACN exhibited a downfield shift in the alcohol proton of 2.55 ppm, which is interpreted as H-bonding between the solvent and the fluoroalcohol. Second, the lower yields experimentally found with the use of 5.0 equiv of HFIP as additive in $CDCl_3$ tentatively implies the importance of the H-bonding network in the pure fluoroalcohol in further enhancing the reactivity of the hypervalent iodine.

Beyond PIDA, PIFA and HTIB are two other commonly utilized and commercially available iodine(III) reagents. Importantly, for the present reaction of study the use of PIFA in HFIP as the oxidant does lead to some formation of the cyclobutane product, however the side reactions are extensive resulting in a significantly lower yield (20%). In contrast the use of HTIB does not result in any product formation in either ACN or HFIP. Table 3 shows the experimentally determined voltammetric reduction potentials of these two other iodine(III) reagents in both HFIP and ACN.

Table 3. Voltammetric Peak Potentials ($E_{p,c}$, V) for the Iodine(III) Compounds PIFA and HTIB versus Fc/Fc⁺, As Measured at 100 mV s⁻¹

	ACN	HFIP
PIDA	-1.32	-0.47
PIFA	-0.56	-0.45
HTIB	-0.42	-0.64

The electronegative nature of the fluoro groups on PIFA serve to significantly enhance the oxidative strength of the reagent in ACN as compared to PIDA (see Table 3 for comparison). Conversely the PIFA voltammetric reduction potential is only 110 mV (vs Fc/Fc⁺) lower in HFIP. This shift is far smaller than found for PIDA, hence it seems credible that the trifluoroacetate ligands significantly decrease the PIFAs hydrogen bond accepting abilities. This conclusion is corroborated by the associated NMR spectra where for a 1:1 mixture of PIFA and HFIP, **16**, in deuterated chloroform (Figure 4c) the HFIP alcoholic proton signal is only downfield shifted 0.6 ppm as compared to the analogous downfield shift of 3.0 ppm for PIDA.

Extra information supporting the weak interaction between PIFA and HFIP is evidenced using a heteronuclear Overhauser effect experiment, ¹H-¹⁹F HOESY (see Supporting Information for a complete study), providing evidence of dipolar spatial interactions between fluorine and protons, nearby in space. On irradiating the PIFA ¹⁹F signal of the equimolecular mixture of PIFA:HFIP, **16**, at -73.5 ppm, significant effects were observed with the HFIP broad OH signal at 3.70 ppm as well as with the HFIP 1'-H at 4.40 ppm.

These results are in stark contrast to those recorded for Koser's reagent (HTIB) where the iodine(III) species is on the basis of the voltammetric data found to be a weaker oxidant in the fluoroalcohol. Again the NMR data (Figure 4e) do not evidence the H-bonding between the HFIP and HTIB to be particularly strong. Interestingly, in this case the ¹H NMR shows a broad signal at 3.90 ppm corresponding to both HTIB-OH and HFIP-OH (Figure 4e); it seems likely that the two protons have become equivalent (a proposed and time-averaged structure is shown in Figure 4e). This would imply HFIP-OH has been downfield shifted by 1.0 ppm, whereas the




Figure 4. ¹H NMR of (a) HFIP (**3**), (b) PIFA, (c) 1:1 mixture of PIFA:HFIP (**16**), (d) HTIB, and (e) 1:1 mixture of HTIB:HFIP (a putative and time-averaged structure of an adduct **17** is shown). Red stars indicate the position of the deshielded OH signal of interest.

HTIB-OH by 2.3 ppm, suggesting that HTIB-OH has assumed an H-bond donor role in the proposed system. This type of interaction is possibly the origin of the “deactivation” of the reagent in HFIP, where the deactivation is evidenced both voltammetrically and synthetically. Finally, complementary DOSY experiments (see SI, section 4, for the complete account of experiments) and the measured diffusion coefficients (see SI Tables 2 and 3) using PIFA and HTIB do not show the existence of very strong H-bonded species upon mixing H-donor HFIP (**3**) and PIFA or HTIB, judged by the small changes on the D/D_{TMS} ratio. These NMR data further corroborate the conclusion that for these two iodine(III) reagents the H-bonding influence is less dominant than found for PIDA.

For the above two hypervalent reagents, it is interesting to comment that the influence of the solvent upon their redox behavior is comparable in magnitude to that found for the alkenes. Hence, the lack of reactivity observed with the HTIB in HFIP is due to the deactivation being significant enough to overcome any gains in the “activation” of the alkenes associated with the use of the fluorinated solvent.

3. CONCLUSIONS

On average, the voltammetric oxidation for the electron-rich styrenes studied within this work are found to occur in HFIP at a lower potential as measured against the ferrocene/ferrocenium redox couple potential as compared to ACN (average shift of ~140 mV as measured from 11 alkenes). This shift in the oxidative wave is attributed to the stabilization of the formed cation. Moreover, the recorded oxidative peak potentials of the styrenes are found to correlate well with the observed synthetic reactivity, such that those that oxidize voltammetrically at a higher potential are found to not synthetically undergo a [2+2] cycloaddition when initiated using hypervalent iodine reagents. Although not insignificant, this shift in potential is small in contrast to the reduction of PIDA in HFIP as compared to ACN, where the reduction wave is observed to shift ~850 mV. Hence, the difference between the voltammetric oxidation of *trans*-anethole and the hypervalent reagent PIDA is over 1 V less in HFIP than in ACN. NMR data strongly evidence this altered reactivity to be due to

the strong H-bonding and adduct formation between the HFIP and PIDA.

Two other iodine(III) reagents, PIFA and HTIB (Koser's reagent) are studied, and the solvent HFIP is shown to have markedly differing effects for these two species. Namely the electron-withdrawing nature of the trifluoroacetate groups present on the PIFA serves to enhance the oxidative ability of the reagent in ACN as compared to PIDA. However, the presence of the fluorine on the PIFA also serves to decrease the H-bonding abilities of the ligands, such that the influence of H-bonding is significantly less. Consequently, in HFIP, both PIFA and PIDA are very comparable in their oxidative strengths. Conversely, for HTIB, the use of HFIP as a solvent decreases the reactivity of the reagent, such that, with its reduction peak occurring at -0.64 V (vs Fc), the reagent is experimentally found to be unable to initiate the cycloaddition. Physically, this weaker H-bonding interaction of the fluoroalcohol with the PIFA and HTIB is confirmed via NMR. Hence, this work has demonstrated how the choice of hypervalent iodine reagent and solvent used can dramatically alter the reactivity of this group of highly versatile synthetic reagents. Moreover, physical insight into the common usage of fluoroalcohol solvents with this class of reagents is provided.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b04057.

Experimental procedures, further electrochemical and NMR data, and compound characterization (HRMS and NMR) (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*timothy.donohoe@chem.ox.ac.uk

*richard.compton@chem.ox.ac.uk

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the European Union and the European Commission for financial support: the research leading to these results has received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme (FP7/2007-2013) and from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC Grant Agreement no. 320403. T. D. W. Claridge is thanked for assistance with the NMR experiments. Dr. Rosimeire Coura Barcelos is thanked for helping with the synthesis of alkenes.

■ REFERENCES

(1) For recent reviews on hypervalent iodines, see: (a) Yoshimura, A.; Zhdankin, V. V. *Chem. Rev.* **2016**, *116*, 3328–3435. (b) Li, Y.; Hari, D. P.; Vita, M. V.; Waser, J. *Angew. Chem., Int. Ed.* **2016**, *55*, 4436–4454. (c) Zhdankin, V. V. *Hypervalent Iodine Chemistry*; John Wiley & Sons Ltd: Chichester, 2013. (2) (a) Kita, Y.; Tohma, H.; Inagaki, M.; Hatanaka, K.; Yakura, T. *Tetrahedron Lett.* **1991**, *32*, 4321–4324. (b) Kita, Y.; Tohma, H.; Hatanaka, K.; Takada, T.; Fujita, S.; Mitoh, S.; Sakurai, H.; Oka, S. *J. Am. Chem. Soc.* **1994**, *116*, 3684–3691. (c) Hamamoto, H.; Hata, K.; Nambu, H.; Shiozaki, Y.; Tohma, H.; Kita, Y. *Tetrahedron Lett.* **2004**, *45*, 2293–2295. (d) Kita, Y.; Morimoto, K.; Ito, M.; Ogawa, C.; Goto,

A.; Dohi, T. *J. Am. Chem. Soc.* **2009**, *131*, 1668–1669. (e) Dohi, T.; Ito, M.; Yamaoka, N.; Morimoto, K.; Fujioka, H.; Kita, Y. *Angew. Chem., Int. Ed.* **2010**, *49*, 3334–3337. (f) Ito, M.; Kubo, H.; Itani, I.; Morimoto, K.; Dohi, T.; Kita, Y. *J. Am. Chem. Soc.* **2013**, *135*, 14078–14081. (g) Morimoto, K.; Sakamoto, K.; Ohshika, T.; Dohi, T.; Kita, Y. *Angew. Chem., Int. Ed.* **2016**, *55*, 3652–3656.

(3) (a) Dohi, T.; Ito, M.; Yamaoka, N.; Morimoto, K.; Fujioka, H.; Kita, Y. *Tetrahedron* **2009**, *65*, 10797–10815. (b) Ebersson, L.; Hartshorn, M. P.; Persson, O.; Radner, F. *Chem. Commun.* **1996**, 2105–2112. (c) Dohi, T.; Yamaoka, N.; Kita, Y. *Tetrahedron* **2010**, *66*, 5775–5785. See also refs 2b and 2e.

(4) (a) Shuklov, I. A.; Dubrovina, N. V.; Börner, A. *Synthesis* **2007**, 2925–2943. (b) Berkessel, A.; Adrio, J. A.; Hüttenhain, D.; Neudörfl, J. M. *J. Am. Chem. Soc.* **2006**, *128*, 8421–8426. (c) Berkessel, A.; Adrio, J. A. *J. Am. Chem. Soc.* **2006**, *128*, 13412–13420.

(5) (a) Ratnikov, M. O.; Tumanov, V. V.; Smit, W. A. *Angew. Chem., Int. Ed.* **2008**, *47*, 9739–9742. (b) Li, G.-X.; Qu, J. *Chem. Commun.* **2010**, *46*, 2653–2655.

(6) Colomer, I.; Coura Barcelos, R.; Donohoe, T. *J. Angew. Chem., Int. Ed.* **2016**, *55*, 4748–4752.

(7) Different reviews concerning aspects of cyclobutane chemistry have been published. For catalytic enantioselective [2+2] cycloaddition, see: (a) Xu, Y.; Conner, M. L.; Brown, M. K. *Angew. Chem., Int. Ed.* **2015**, *54*, 11918–11928. For an update of thermal cyclobutane formation, see: (b) Alcaide, B.; Aragoncillo, C.; Almendros, P. In *Comprehensive Organic Synthesis II*, 2nd ed.; Knochel, P., Ed.; Elsevier: Amsterdam, 2014; pp 66–84. For [2+2] photocycloaddition using α,β -unsaturated ketones, see: (c) Margaretha, P. *Helv. Chim. Acta* **2014**, *97*, 1027–1035. Alternative methods for the synthesis of cyclobutanes recently reported involve Fe-catalyzed [2+2] cycloaddition of unactivated alkenes: (d) Hoyt, J. M.; Schmidt, V. A.; Tondreau, A. M.; Chirik, P. J. *Science* **2015**, *349*, 960–963. Catalyzed boration and cyclization of homoallylic sulfonates: (e) Ito, H.; Toyoda, T.; Sawamura, M. *J. Am. Chem. Soc.* **2010**, *132*, 5990–5992. Extensive work has been done on the metal-catalyzed intramolecular [2+2] of allenes. For nickel, see: (f) Noucti, N. N.; Alexanian, E. J. *Angew. Chem., Int. Ed.* **2015**, *54*, 5447–5450. For ruthenium, see: (g) Gulías, M.; Collado, A.; Trillo, B.; López, F.; Oñate, E.; Esteruelas, M. A.; Mascareñas, J. L. *J. Am. Chem. Soc.* **2011**, *133*, 7660–7663. For gold, see: (h) Luzung, M. R.; Mauleón, P.; Toste, F. D. *J. Am. Chem. Soc.* **2007**, *129*, 12402–12403.

(8) For recent reviews concerning cyclobutane-containing natural products, see: (a) Dembitsky, V. M. *Phytomedicine* **2014**, *21*, 1559–1581. (b) Sergeiko, A.; Poroikov, V. V.; Hanuš, L. O.; Dembitsky, V. M. *Open Med. Chem. J.* **2008**, *2*, 26–37. (c) Dembitsky, V. J. *Nat. Med.* **2008**, *62*, 1–33.

(9) Hypervalent iodine reagents have been proposed to act as one-electron oxidants. For the generation of amidyl *N*-centered radicals using IBX, see: (a) Nicolaou, K. C.; Baran, P. S.; Zhong, Y.-L.; Vega, J. A. *Angew. Chem., Int. Ed.* **2000**, *39*, 2525–2529. (b) Nicolaou, K. C.; Baran, P. S.; Zhong, Y.-L.; Barluenga, S.; Hunt, K. W.; Kranich, R.; Vega, J. A. *J. Am. Chem. Soc.* **2002**, *124*, 2233–2244. (c) Janza, B.; Studer, A. *J. Org. Chem.* **2005**, *70*, 6991–6994. (d) Nicolaou, K. C.; Baran, P. S.; Kranich, R.; Zhong, Y.-L.; Sugita, K.; Zou, N. *Angew. Chem., Int. Ed.* **2001**, *40*, 202–206.

(10) For the use of triaryl amine radical cation, see: (a) Bauld, N. L.; Pabon, R. *J. Am. Chem. Soc.* **1983**, *105*, 633–634. For electrochemical anodic oxidation, see: (b) Chiba, K.; Miura, T.; Kim, S.; Kitano, Y.; Tada, M. *J. Am. Chem. Soc.* **2001**, *123*, 11314–11315. For seminal examples using a photoredox approach, see: (c) Ischay, M. A.; Lu, Z.; Yoon, T. P. *J. Am. Chem. Soc.* **2010**, *132*, 8572–8574. (d) Ischay, M. A.; Ament, M. S.; Yoon, T. P. *Chem. Sci.* **2012**, *3*, 2807–2811. (e) Riener, M.; Nicewicz, D. A. *Chem. Sci.* **2013**, *4*, 2625–2629. For a related use of radical cations in metal-free cross- and ring-opening metathesis using electrochemistry, see: (f) Miura, T.; Kim, S.; Kitano, Y.; Tada, M.; Chiba, K. *Angew. Chem., Int. Ed.* **2006**, *45*, 1461–1463. For photoredox, see: (g) Ogawa, K. A.; Goetz, A. E.; Boydston, A. J. *J. Am. Chem. Soc.* **2015**, *137*, 1400–1403. (h) Goetz, A. E.; Boydston, A. J. *J. Am. Chem. Soc.* **2015**, *137*, 7572–7575.

(11) For seminal contributions on the field of H-bond catalysis, see: (a) Raheem, I. T.; Thiara, P. S.; Peterson, E. A.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2007**, *129*, 13404–13405. (b) Albrecht, L.; Dickmeiss, G.; Acosta, F. C.; Rodríguez-Esrich, C.; Davis, R. L.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2012**, *134*, 2543–2546. For a review focused on H-bonding in organic chemistry, see: (c) Doyle, A. G.; Jacobsen, E. N. *Chem. Rev.* **2007**, *107*, 5713–5743. For a general review on H-bonding, see: (d) Steiner, T. *Angew. Chem., Int. Ed.* **2002**, *41*, 48–76.

(12) Demaille, C.; Bard, A. J. *Acta Chem. Scand.* **1999**, *53*, 842–848.

(13) (a) Ebersson, L.; Hartshorn, M. P.; Persson, O.; Radner, F. *Chem. Commun.* **1996**, *18*, 2105–2112. (b) Ebersson, L.; Hartshorn, M. P.; McCullough, J. J.; Persson, O.; Radner, F. *Acta Chem. Scand.* **1998**, *52*, 1024–1028.

(14) Francke, R.; Cericola, D.; Kötz, R.; Weingarth, D.; Waldvogel, S. R. *Electrochim. Acta* **2012**, *62*, 372–380.

(15) (a) Chong, D.; Stewart, M. P.; Geiger, W. E. *J. Am. Chem. Soc.* **2009**, *131*, 7968–7969. (b) Stewart, M. P.; Lam, K.; Chong, D.; Geiger, W. E. *J. Electroanal. Chem.* **2015**, *743*, 68–77.

(16) Silva, L. F., Jr; Vasconcelos, R. S.; Lopes, N. P. *Int. J. Mass Spectrom.* **2008**, *276*, 24–30.

(17) A very minor set of peaks, that represents less than 5%, can be observed both in the ¹H NMR spectrum of PIDA (2) and in the 1:1 mixture of PIDA:HFIP (5), which we have tentatively assigned to an impurity derived from PIDA (2).

(18) For a correlation of the OH ¹H NMR shift in HFIP with H-bonding, see: (a) Purcell, K. F.; Stikeleather, J. A.; Brunk, S. D. *J. Mol. Spectrosc.* **1969**, *32*, 202–213. (b) Juršić, B.; Ladika, M.; Sunko, D. E. *Tetrahedron Lett.* **1985**, *26*, 5323–5324.

(19) Brønsted and Lewis acid catalyzed trifluoromethylation reactions using Togni reagents have been reported: (a) Koller, R.; Stanek, K.; Stolz, D.; Aardoom, R.; Niedermann, K.; Togni, A. *Angew. Chem., Int. Ed.* **2009**, *48*, 4332–4336. (b) Koller, R.; Huchet, Q.; Battaglia, P.; Welch, J. M.; Togni, A. *Chem. Commun.* **2009**, 5993–5995. (c) Santschi, N.; Geissbühler, P.; Togni, A. *J. Fluorine Chem.* **2012**, *135*, 83–86. For a very recent contribution on the isolation and reactivity of iodonium salts, see: (d) Brantley, J. N.; Samant, A. V.; Toste, F. D. *ACS Cent. Sci.* **2016**, *2*, 341–350.

(20) (a) Kapur, G. S.; Cabrita, E. J.; Berger, S. *Tetrahedron Lett.* **2000**, *41*, 7181–7185. (b) Cabrita, E. J.; Berger, S. *Magn. Reson. Chem.* **2001**, *39*, S142–S148. (c) Martins de Carvalho, E. M.; Velloso, M. H. R.; Tinoco, L. W.; Figueroa-Villar, J. D. *J. Magn. Reson.* **2003**, *164*, 197–204.

(21) (a) It is well recognized that H-bonding to an anionic species is often stronger than to a neutral molecule (see ref 21b, for example). On this basis it is credible that the altered reactivity of PIDA may relate to the greater stabilizing abilities of the HFIP towards the formed radical anion as compared to the neutral PIDA reactant. However, such an analysis certainly does not preclude the possibility of PIDA-HFIP adduct formation lowering the reorganization energy associated with the solution phase SET. (b) Quan, M.; Sanchez, D.; Wasylkiw, M. F.; Smith, D. K. *J. Am. Chem. Soc.* **2007**, *129*, 12847–12856.

(22) Beyond the use of PIDA, our previous work has demonstrated that the iodine(V) reagent Dess–Martin periodinane (DMP) can also be used in the synthesis of cyclobutanes in HFIP but not ACN. On the basis of the presence of the acetate groups one may anticipate that again the altered reactivity relates to the influence of H-bonding between the reagent and the fluorinated solvent. Certainly, the voltammetric data demonstrates a shift in the reduction peak ($E_{p,c}$ in ACN = –0.55 V, $E_{p,c}$ in HFIP = –0.45 V) comparable to that found for PIFA. However, initial NMR data indicate that the speciation and reaction of the DMP with HFIP is far more complex and plausibly warrants further attention.